



OPEN ACCESS

Core elements of serious illness conversations: an integrative systematic review

Rebecca Baxter ¹, Susanna Pusa ¹, Sofia Andersson ¹,
Erik K Fromme ^{2,3}, Joanna Paladino ^{2,3,4}, Anna Sandgren ¹

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/spcare-2023-004163>).

¹Center for Collaborative Palliative Care, Department of Health and Caring Sciences, Linnaeus University—Vaxjo Campus, Vaxjo, Sweden

²Ariadne Labs, Boston, Massachusetts, USA

³Harvard Medical School, Boston, Massachusetts, USA

⁴Massachusetts General Hospital, Boston, Massachusetts, USA

Correspondence to

Dr Rebecca Baxter, Center for Collaborative Palliative Care, Department of Health and Caring Sciences, Linnaeus University - Vaxjo Campus, Vaxjo 351 95, Sweden; rebecca.baxter@lnu.se

Received 9 January 2023

Accepted 5 June 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY. Published by BMJ.

To cite: Baxter R, Pusa S, Andersson S, *et al.* *BMJ Supportive & Palliative Care* Epub ahead of print: [please include Day Month Year]. doi:10.1136/spcare-2023-004163

ABSTRACT

Background Ariadne Labs' Serious Illness Care Program (SICP), inclusive of the Serious Illness Conversation Guide (SICG), has been adapted for use in a variety of settings and among diverse population groups. Explicating the core elements of serious illness conversations could support the inclusion or exclusion of certain components in future iterations of the programme and the guide.

Aim This integrative systematic review aimed to identify and describe core elements of serious illness conversations in relation to the SICP and/or SICG.

Design Literature published between 1 January 2014 and 20 March 2023 was searched in MEDLINE, PsycINFO, CINAHL and PubMed. All articles were evaluated using the Joanna Briggs Institute Critical Appraisal Guidelines. Data were analysed with thematic synthesis.

Results A total of 64 articles met the inclusion criteria. Three themes were revealed: (1) serious illness conversations serve different functions that are reflected in how they are conveyed; (2) serious illness conversations endeavour to discover what matters to patients and (3) serious illness conversations seek to align what patients want in their life and care.

Conclusions Core elements of serious illness conversations included explicating the intention, framing, expectations and directions for the conversation. This encompassed discussing current and possible trajectories with a view towards uncovering matters of importance to the patient as a person. Preferences and priorities could be used to inform future preparation and recommendations. Serious illness conversation elements could be adapted and altered depending on the intended purpose of the conversation.

BACKGROUND

To provide much-needed guidance for conversations about serious illness,

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ The Serious Illness Care Program (SICP) and Serious Illness Conversation Guide (SICG) are associated with improved patient outcomes and experiences.
- ⇒ Serious illness conversation content has been adapted for different patients, clinicians and contexts, yet the core elements of these conversations have not been explored.

WHAT THIS STUDY ADDS

- ⇒ Conversation elements were revealed to be multifaceted with nuanced content that could be altered depending on the intended purpose of the conversation.
- ⇒ Core conversation elements included having clear intentions and framing, establishing expectations and directions, exploring the current situation and possible trajectory, uncovering matters of importance, elucidating preferences and priorities and supporting preparation and recommendations.
- ⇒ While modifications have been made to the conversation guide, the same general questions and structure were relevant for most contexts.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ This integrative systematic review contributes important knowledge about core elements of serious illness conversations that can be used in developing or modifying future iterations of the SICP and SICG.
- ⇒ Informing the core elements for serious illness conversations strengthens the theory supporting the programme and guide and can be used to inform current clinical education and practice.

experts at Ariadne Labs (Boston, Massachusetts, USA) developed the Serious Illness Care Program (SICP), inclusive of the Serious Illness Conversation Guide

(SICG).¹ These conversations aim to elicit seriously ill patients' values and goals to ensure that they receive information and care that meets their needs.² Studies exploring the effect of the programme and the guide have found that timely serious illness conversations can reduce patient stress and anxiety, decrease resource utilisation, result in more goal-concordant discussions and improve healthcare professionals' experiences of care provision.^{3–6} While the original SICP and SICG were developed for the oncology context, in recent years both the programme and the guide have been adapted and implemented in myriad clinical settings and languages.^{1 2 7}

The original SICG outlined key conversation areas, including: illness understanding, decision making and information preferences, prognostic disclosure, patient goals and fears, views on acceptable function and trade-offs and desire for family involvement⁷; however, it has been acknowledged that the guide was not comprehensive⁷ and that other important conversation domains exist.⁸ As the programme and the guide continue to be developed, adapted and implemented, it is necessary to explicate the 'core elements' of serious illness conversations to ensure that these components are present—or justifiably absent. For the purpose of this study, the term 'core elements' refers to necessary and/or important parts of serious illness conversations.⁹ The aim of this integrative systematic review was to identify and describe core elements of serious illness care conversations in the context of the SICP and/or SICG.

METHODS

Search strategy

The search was conducted on 20 March 2023 in the bibliographic databases CINAHL, MEDLINE, PsycINFO and PubMed using the search strategy described in online supplemental material A. The search terms were established in collaboration with a university librarian. As the SICP was developed based on a literature review from 2014, the search was limited to articles published in English between 1 January 2014 and 20 March 2023. Ariadne Labs also provided a list of known publications related to the SICP (n=44).

Eligibility criteria

Eligibility criteria were developed a priori to ensure relevance to the study aim. Articles were eligible for inclusion if they: (a) explicitly stated a connection with the SICP, SICG and/or Ariadne Labs in the title, abstract or main text and (b) provided a meaningful description of at least one serious illness conversation element. The publication language was limited to English. No restrictions were applied regarding population or setting; however, book chapters, letters to the editor and conference abstracts were excluded.

Selection process

Known foundational articles were identified within the search results, including the original SICP development papers from Bernacki *et al*,^{2 7} which confirmed good sensitivity of the search strategy. Duplicate publications were removed. One author (SP) reviewed titles, abstracts, keywords and, when required, full-text articles against the inclusion criteria to identify eligible articles. Any uncertainty regarding initial inclusion was discussed with AS and RB. Next, full-text articles were screened for inclusion by RB and SP. Reference lists of included articles were hand searched.

Data collection process

Three authors (RB, SP, SA) independently extracted data from six articles to calibrate the data extraction and tabulation process. Thereafter, RB extracted data by going through each article line-by-line to identify data relevant to the study aim and copying this to the extraction form described below. Only unreferenced original data were considered for extraction from the methods, results, discussion and/or conclusions sections of articles (data from the abstract, key messages, introduction and/or background sections were therefore ineligible). Unreferenced data referred to text that was presented as original without direct citation to another source. Any uncertainty regarding data eligibility was discussed between RB, SP and SA.

Data items

An extraction form was used by RB to manually tabulate data regarding the authors, year of publication, article type, clinical context, clinicians/users, and if/how the SICP/SICG were implemented. In addition, data were extracted for tabulation regarding (a) descriptions of serious illness conversation elements and/or (b) descriptions of serious illness conversation content. This encompassed data pertaining to any time point (eg, past, present and theoretical), article type (eg, original research, case studies and clinical updates) and participant group (eg, patient, family, staff and researcher).

Risk of bias assessment

The risk of bias was assessed using the Joanna Briggs Institute (JBI) critical appraisal checklists.¹⁰ These 13 checklists are used to evaluate the trustworthiness, relevance and results of published research. As there is not yet a checklist for mixed methods studies, JBI provided advice via email that the completion of more than one checklist could be appropriate for studies that enlisted more than one method. If an article presented data, even in descriptive form, one of the checklists for research studies was selected (ie, the checklist for text and opinion was not selected). Articles were assessed by responding 'yes', 'no', 'unclear' or 'not applicable' to each checklist item. If the criteria for an item were only partially fulfilled, the item was

marked as 'unclear'. One author (SP) conducted the initial critical appraisal of all articles, and any questions regarding study type or checklist selection were discussed with RB and AS. Articles were not excluded based on the appraisal responses, instead the checklists were used to inform article characteristics and comparability to support a complete discussion of the current literature. To minimise bias, JP and EKF, who authored several articles included in this review, were not involved in the article selection, data extraction or critical appraisal process.

Synthesis methods

Thematic synthesis was selected as it provides a set of established methods for the identification of patterns and development of analytic themes in textual data.¹¹⁻¹³ This consisted of three stages: free line-by-line coding, organisation of codes into descriptive themes and development of analytical themes.¹¹ First, data were inductively interrogated for descriptions of conversation elements and coded based on the content of these descriptions. Following this, data were examined and coded for descriptions of the SICG and its content. Similar codes were compared and grouped into descriptive subthemes that remained close to the data. Lastly, the findings were synthesised and analytic themes were constructed to provide novel interpretations. The author group comprised of nurses (RB, SP, SA, AS) and physicians (EKF, JP) with experience in research and clinical practice, and extensive expertise in development and implementation of the SICP. The results were discussed and refined among the author group.

RESULTS

Study selection

The search retrieved 698 articles and a further 44 articles were provided by Ariadne Labs (figure 1). Duplicates were removed (n=436). Title, abstract and full-text screening of 306 articles were undertaken, resulting in the elimination of 216 articles. The remaining 90 full-text articles were assessed against the eligibility criteria, and the reference lists of these articles were manually searched. The reference list search revealed eight articles for full-text review; however, none met the inclusion criteria. In total, 64 articles met the inclusion criteria. Of these, 62 articles (97%) were identified through the database search and two (3%) were identified through the list provided by Ariadne Labs.

Study characteristics

The majority of articles (n=55) were original research articles, of which 13 used qualitative methods, 15 used some form of mixed-methods and 27 used quantitative methods. Nine were categorised as text and opinion articles. Of the 64 included articles, 54 represented unique studies. Seven study clusters were identified,

including the Dana-Farber Cancer Institute cluster (n=8),^{4 5 7 14-18} Brigham integrated Care Management Program cluster (n=2),^{19 20} Massachusetts General Hospital cluster 1 (n=2),^{21 22} Massachusetts General Hospital cluster 2 (n=2),^{23 24} University of Pennsylvania (n=2)^{25 26} and Meta-network Learning and Research Center Advance Care Planning cluster (n=2).^{27 28} The list of included articles, country, JBI checklist selection, clinical context and implementation/adaptation of the SICP and/or SICG is summarised in table 1.

Most articles were from North American inpatient clinical settings. Descriptions of SICP implementation and SICG version varied considerably. SICP implementation/adaptation ranged from none or unstructured training, to multiple hours of formal training. SICG implementation/adaptation was reported as including the original guide (various versions), to using a guide that had been modified for different patients, clinicians and clinical or cultural contexts.

Critical appraisals

Most articles reported clear aims and objectives. Strategies for sampling and data collection methods were largely well defined; however, strategies for dealing with confounding factors were often not stated. In studies that reported qualitative data, there was a lack of reflection about the influence of the researcher on the research (or vice versa), and few located researchers' cultural or theoretical backgrounds. Detailed JBI critical appraisal checklist responses are presented in online supplemental material B.




Figure 1 PRISMA Flow Diagram. SICG, Serious Illness Conversation Guide; SICP, Serious Illness Care Program.

Table 1 Summary of included articles, ordered by publication year

Author, year	Country	JBI checklist*	Clinical context	SICP implementation/adaptation	SICG implementation/adaptation
Bernacki <i>et al</i> , 2015 ⁷	USA	13	Oncology	Original SICP: 2.5-hour training to use SICG	Original SICG version: R4.2 12-10-13
Lakin <i>et al</i> , 2017 ¹⁹	USA	10	Primary care	Adapted SICP for the primary care setting—2.5 hours	SICG (version not stated)
Lamas <i>et al</i> , 2017 ³⁵	USA	1	Acute care	Not applicable as interviewer was from research team	Adapted SICG for long-term acute care patients
Mandel <i>et al</i> , 2017 ³⁷	USA	13	Nephrology	Proposed: 2.5 hours training in basic SICG competencies	Proposed: SICG (version not stated)
Miranda <i>et al</i> , 2018 ¹⁴	USA	1&9	Oncology	SICP implemented	SICG (version not stated)
O'Donnell <i>et al</i> , 2018 ⁵⁰	USA	10	Heart failure	Education about ACP and role of healthcare proxy	Discussion based on SICG (version not stated)
Baran <i>et al</i> , 2019 ⁵⁹	USA	13	Primary care	Not implemented	SICG version: 2015–2017
Geerse <i>et al</i> , 2019 ⁵	USA	9	Oncology	2.5 hours skills-based training to use the SICG	SICG version: R4.2 12-10-13
Lakin <i>et al</i> , 2019 ²⁰	USA	9	Primary care	SIC skills training—3-hour interactive session	SICG version: R4.2 12-10-13 (2012)
Massmann <i>et al</i> , 2019 ⁴³	USA	10	Primary care	2-hour training based on the SICP model	SICG (version not stated)
McGlinchey <i>et al</i> , 2019 ⁶³	UK	9	UK health setting	Adapted SICP for UK healthcare setting	Adapted SICG for UK healthcare setting
Paladino <i>et al</i> , 2019 ⁴	USA	11	Oncology	2.5-hour skills-based training session on the SICG	SICG version: R2.7 05-25-12
Tam <i>et al</i> , 2019 ⁵²	Canada	9&10	Internal medicine	2.5-hour small group session derived from SICP training	SICG version: 2017-04-18
Daubman <i>et al</i> , 2020 ²³	USA	1	Multiple contexts	Adapted 2.5–3 hours SICG training	Modified SICG - Partners SICG
Gace <i>et al</i> , 2020 ²¹	USA	5	General medical	2.5 hours of SIC training modified from Ariadne Labs	SICG (version not stated)
Gelfand <i>et al</i> , 2020 ⁴⁰	USA	13	Kidney care	NephroTalk, VitalTalk, Palliative Education	Adapted SICG from version: 2017-04-21
Greenwald <i>et al</i> , 2020 ²²	USA	5	Medical inpatient	SICP implemented	SICG (version not stated)
Jain <i>et al</i> , 2020 ²⁹	USA	13	Not stated	Recommends SICP training as communication resource	Refers to SICG as a communication tool
Ko <i>et al</i> , 2020 ³⁶	Canada	1	Oncology	15 min SICG introduction, no standardised training	SICG (version not stated)
Kumar <i>et al</i> , 2020 ⁸	USA	1&9	Outpatient oncology	3-hour SICP structured communication education	SICG (version not stated)
Lally <i>et al</i> , 2020 ⁵³	USA	10	Hospital patients	Communication skills training programme developed	Modified SICG for nurses (Administered by phone)
Ma <i>et al</i> , 2020 ⁵¹	Canada	10	Internal medicine	Adapted SICP, 2.5-hour workshop	SICG version: 2016
Manz <i>et al</i> , 2020 ²⁵	USA	11	Oncology	SICG training 3 months prior to the start of the trial	ACP template based on the SICG (version not stated)
Ouchi <i>et al</i> , 2020 ⁶⁸	USA	13	Emergency	Not stated	Code status conversation guide – adapted from SICG
Paladino <i>et al</i> , 2020 ¹⁶	USA	1&9	Oncology	SICP implemented	SICG version: 2015–2017
Paladino <i>et al</i> , 2020 ¹⁷	USA	11	Oncology	SICP implemented	SICG (version not stated)
Paladino <i>et al</i> , 2020 ⁵⁷	USA	9&10	Health systems (3)	SICP implemented—2.5–3 hour clinician training on SIC	SICG (version not stated)
Pasricha <i>et al</i> , 2020 ⁴⁴	USA	1&9	Intensive care	SICP training (3 hours)	SICG for surrogates. Based on SICG 2017-04-18
Pottash <i>et al</i> , 2020 ³⁹	USA	1&9	Ambulatory care	Short SICG introduction, video, and role play	Adapted SICG 'Advanced Illness Conversation Guide'
Sirianni <i>et al</i> , 2020 ³⁰	Canada	13	COVID-19	Yes, discussed resources for SIC	SICG version: 2018-04-18
van Breemen <i>et al</i> , 2020 ⁵⁸	Canada	3	Paediatrics	Training to use the SICG-Peds	Adapted SICG-Peds (12–2019)
Wasp <i>et al</i> , 2020 ⁵⁶	USA	9&10	Onco & Haematology	Adapted 3-hour SICG education with 4-hour VitalTalk session	SICG Communication Skills Assessment Tool

Continued

Table 1 Continued

Author, year	Country	JBI checklist*	Clinical context	SICP implementation/adaptation	SICG implementation/adaptation
Aaronson <i>et al</i> , 2021 ³⁸	USA	9	Emergency	Not stated	Adapted Partners SICG for social workers
Beddard-Huber <i>et al</i> , 2021 ³³	Canada	13	General	SIC interprofessional clinician workshop 2.5 hour	Adapted SICG for substitute decision-makers
Daly <i>et al</i> , 2021 ²⁷	USA	1	Family medicine	SICP implemented	SICG (version not stated)
DeCoursey <i>et al</i> , 2021 ⁴⁵	USA	9	Paediatrics	Adapted PediSICP	Adapted PediSICG
Geerse <i>et al</i> , 2021 ¹⁵	USA	1&9	Oncology	2.5-hour skills-based training to use the SICG	SICG version: R4.2 12-10-13
Greenwald <i>et al</i> , 2021 ⁴¹	USA	10	Hospital setting	Clinicians participated in two 1-hour training sessions	Adapted Partners SICG—COVID-19 (03-2020)
Hafid <i>et al</i> , 2021 ⁴⁶	Canada	9&10	Primary care	SICP adapted and implemented	SICG adapted (version not stated)
Karim <i>et al</i> , 2021 ³⁴	Canada	10	Outpatient oncology	Adapted 2-hour training session based on the SICP	SICG version: 2017-04-18
Lagrotteria <i>et al</i> , 2021 ⁵⁴	Canada	9	Tertiary hospitals	SICP implemented in 2.5-hour interactive training session	SICG version: draft R4.2 12-10-13
Lakin <i>et al</i> , 2021 ³¹	USA	10	General medicine	Adapted SICP—3-hour SAGE programme	SICG (version not stated)
Le <i>et al</i> , 2021 ⁴⁷	Canada	5	Acute medicine	SIC education provided in new employee orientation	SICG (version not stated)
Moye <i>et al</i> , 2021 ⁶⁹	USA	5&9	Older adults	SICP not implemented	Used six questions from the SICG
Paladino <i>et al</i> , 2021 ⁶⁰	USA	9	Inpatient, outpatient	Suggests adaptation of SICP to virtual training	Adapted COVID-19 Outpatient and Inpatient Guides
Paladino <i>et al</i> , 2021 ¹⁸	USA	9	Primary care	SICP training provided	SICG version: 2017-04-18
Reed-Guy <i>et al</i> , 2021 ⁴⁸	USA	1&9	Glioblastoma	SICP implemented	SICG used and adapted (version not stated)
Swiderski <i>et al</i> , 2021 ⁷¹	USA	9	Primary care	SICG training—Two 1-hour sessions	SICG (version not stated)
Thamcharoen <i>et al</i> , 2021 ⁴⁹	USA	1&9	Kidney disease	Interviewer trained in SICG	Adapted SICG for researcher
Borregaard Myrhøj <i>et al</i> , 2022 ³²	Denmark	9	Multiple myeloma	Team training in SIC focusing on existential issues	Modified SICG (Danish version)
Bowman <i>et al</i> , 2022 ⁶²	USA	1	Emergency/COVID-19	SICP training provided	SICG (version not stated)
Daly <i>et al</i> , 2022 ²⁸	USA	1	Family Medicine	Adapted SICP 1.5-hour in-person training	SICG (version not stated)
Davoudi <i>et al</i> , 2022 ⁶⁶	USA	1	Oncology	SICP implemented	SICG (version not stated)
Jacobsen <i>et al</i> , 2022 ²⁴	USA	1	Palliative care	SICP implemented	Partners SICG
Karim <i>et al</i> , 2022 ⁴²	USA	13	Oncology	SICP training in-person or virtual workshops	SICG (version not stated)
Hu <i>et al</i> , 2022 ⁷²	USA	1	General Surgery	Not stated	SICG (version not stated)
King <i>et al</i> , 2022 ⁵⁵	Canada	1	Internal Medicine	All components of SICP implemented	SICG (version not stated)
Li <i>et al</i> , 2022 ²⁶	USA	11	Oncology	All clinicians were trained in the use of the SICG	SICG (version not stated)
LoCastro <i>et al</i> , 2022 ⁶⁴	USA	9	Haematology	Adapted SICP for delivery via telehealth	Adapted SICG for delivery via telehealth
Sanders <i>et al</i> , 2022 ⁷⁰	USA	9&10	Multiple contexts	SICP 2.5-hour in-person training	SICG version: 04-2017 and revised SICG
Wasp <i>et al</i> , 2022 ⁷³	USA	10	Oncology	3-hour SICP training in use of SICG	SICG (version not stated)
Xu <i>et al</i> , 2022 ⁶⁷	USA	9	Primary Care	2.5-hour training in using SICG	SICG (version not stated)
Zehm <i>et al</i> , 2022 ⁶¹	USA	9&10	Education	Adapted SICP workshop with 2.5-hour training	Modified SICG—partners SICG
Garcia <i>et al</i> , 2023 ⁶⁵	USA	9	Inpatient Clinical	Adapted team-based SICP	SICG (version not stated)

*JBI Checklist number: 1—analytical cross sectional studies; 2— case control studies; 3—case reports; 4—case Series; 5—cohort studies; 6—diagnostic test accuracy studies; 7—economic evaluations; 8—prevalence studies; 9—qualitative research; 10—quasi-experimental studies; 11—randomised controlled trials; 12—systematic reviews; 13—text and opinion.

ACP, advance care planning; ICU, intensive care unit; JBI, Joanna Briggs Institute; NP, nurse practitioner; QoL, quality of life; SIC, serious illness conversation; SICG, Serious Illness Conversation Guide; SICP, Serious Illness Care Program.

Table 2 Overview of themes and subthemes

Themes	Subthemes
Serious illness conversations serve different functions that are reflected in how they are conveyed	Intentions and framing Expectations and directions
Serious illness conversations endeavour to discover what matters to patients	Current situation and possible trajectory Matters of importance
Serious illness conversations seek to align what patients want in their life and care	Preferences and priorities Preparation and recommendations

Thematic synthesis

Three themes and six subthemes emerged to describe the core elements of serious illness conversations (see table 2).

Serious illness conversations serve different functions that are reflected in how they are conveyed

The ways in which serious illness conversations were understood and conveyed impacted how the conversation was framed with respect to the clinician, the patient or the context. This theme is comprised of two subthemes: (a) intentions and framing and (b) expectations and directions.

Intentions and framing

The intentions and framing of the conversation described what clinicians wanted to accomplish using the guide, rather than the content of the guide itself. This included checking in, conveying medical updates or discussing the risks and benefits of treatment options,^{29 30} as well as allowing for the expression of goals, values^{23 31} and wishes and hopes for the future.³² Language varied when framing serious illness conversations for patients, such as: discussing future expectations,³³ discovering what is important,³⁴ conferring goals, expectations and experiences,³⁵ hoping for the best and preparing for the worst,³⁶ thinking and preparing,³⁷ looking at the bigger picture³⁸ and discussing health and future expectations.³⁹

Formally introducing the conversation involved explicitly stating what it would be about, establishing an agenda, or seeking permission.^{18 29 40} Clinicians could present the conversation as an opportunity to think ahead or plan in advance³³ in relation to the patient's care⁴⁰ or medical condition.^{31 41} It might be stated from the outset that the aim of the conversation was to inform future decisions and care,³³ or the decision-making aspect could be minimised.^{37 40 42} Serious illness conversations were articulated as being part of, conceptually overlapping with, or recorded as: Advance Care Planning^{5 7 15 24 25 27 34 36 43–49} (categorised as Advance Care Planning in the electronic medical record),^{4 21 23 26 31} End-of-Life conversations,^{4 7 8 14 19 31 43 47 48 50 51} Goals-of-Care conversations^{7 21 29 31 49 50 52–55} or Values and Goals conversations (inclusive of values-based/values-centred/goals-based/goals-centred conversations).^{7 8 17 18 20–23 31 33 35 42 48 56} Conversation framing was therefore informed by diverse understandings

of the concept of serious illness conversations and communicated in different ways depending on the perceived intention of the conversation.

Expectations and directions

Establishing expectations and directions included ascertaining what the conversation aimed to achieve, determining what subjects the conversation would address, and how much information the patient wanted or was ready to receive.^{18 29 30 33} Different versions of the SICG reflected variations in the preferred language used by clinicians, such as 'setting up the conversation', 'opening the conversation' or 'initiating the conversation'.^{33 37 40 57} It was important to ask patients about the amount and type of information they required^{30 40 49} so that they could indicate whether they wanted (or were ready) to have an in-depth conversation about specific concerns or questions.^{29 30 32 36} By establishing expectations from the outset, the discussion could be adapted to suit the needs of the patient before providing updates or clarifications.³⁰ This could help to focus only on issues that patients deemed relevant^{32 36} making the conversation less prescriptive and more collaborative.⁵⁸

Discussing the patient's lived experience was important to the conversation, but the ways in which this was broached varied depending on the clinician and whether the discussion was centred around the patient's understanding of their 'illness/medical condition',^{29 30 36 40 49 57 59} or their 'health'.^{38 41 60} Orientating the conversation around the 'illness' was thought to give the clinician insight into how the patient was coping, their awareness of what was ahead and the extent to which they had accepted their illness,^{29 37 59} particularly if their function or status had changed.⁵⁸ Centering the discussion around 'health' may be viewed as more holistic and could invite conversation about how patients from a variety of clinical contexts feel generally, not only in the context of their illness.^{37 38 41} It was therefore necessary to establish expectations surrounding patient understanding, acceptance, readiness and willingness early in the conversation as this could influence subsequent elements.^{22 37}

Serious illness conversations endeavour to discover what matters to patients

It was important to consider current and possible trajectories when seeking information from patients about what mattered to them in relation to their illness or health. This theme is comprised of two subthemes: (a) current situation and possible trajectory and (b) matters of importance.

Current situation and possible trajectory

Discussing the current situation was thought to enhance patients' understanding of their lived reality and possible trajectories. This was termed as delivering or conveying serious news,⁴⁷ giving medical updates,²⁹ delivering prognosis,^{29 61} sharing prognosis,^{44 58} clarifying prognostic awareness^{45 59 61} and assessment of prognostic understanding.^{8 16 48 51} This subject had to be broached with care as patients could feel anxious talking (or not talking) about death or dying, and some may not want to receive prognostic information.^{7 15} Asking for permission to divulge this information was therefore an important conversation element.^{29 48 52} A prognostic discussion was still thought to be possible even if the clinician was unsure of the exact prognosis.⁶² However, prognosis might be omitted due to comfort or confidence to discuss such topics, or if it was outside clinicians' professional scope of practice.^{5 15 35 38 53} Some elected not to focus on prognosis because discussing preferences in other domains was thought to be sufficient,^{35 38} but it was acknowledged that talking about prognosis could influence how patients answered subsequent questions.^{16 37 49 58 59} Indeed, if some form of prognosis or illness trajectory was not addressed, patients and clinicians may not be able to take full advantage of the possible benefits of the conversation.^{39 40 59}

Gauging the patient's level of trajectorial or situational awareness laid the foundation for how clinicians could clarify uncertainty for patients.^{29 33 41 48} Positive or negative wording could be used to portray information, with some recommending the use of hope/worry statements,^{23 29 63} wish/worry/wonder statements^{18 33 58} and hope for the best plan for the worst statements.^{33 37} The guide offered language templates for sharing time-based, function-based or uncertain prognoses.⁷ However, if clinicians were not comfortable providing a concrete time-based estimate, or if patients were ambivalent about receiving such information, more general information could be provided in the context of the patient's clinical condition.^{37 41 44 59 64} This might include expected decline (ie, function, cognition, condition),^{23 31 33 37 48} expected symptoms or events related to the illness/condition,^{18 46 48} worsening trajectories,^{33 47} quality of life,^{31 48} fragility/stability⁵⁸ and/or treatment options.^{5 31} Even if the prognosis was poor, it was important that hope and positivity was still conveyed⁶⁴ with a view towards supporting patients through their concerns.⁶⁵

Matters of importance

Discussion of important matters encompassed goals and fears, views on acceptable function (critical abilities), trade-offs and desires for family involvement.⁷ This provided opportunities for patients to express their thoughts and feelings, to discover what makes life meaningful and to reflect on important subjects to better plan care.^{5 19 23 26 29 30 32 63} The vernacular for exploring what was important varied in both the conversation guide and in the literature describing the conversation, but most encompassed some combination of the terms: values, goals, wishes, hopes, concerns, worries and fears.^{4 5 8 20–23 25 27 30–33 35 36 38 40 41 43 45 47–49 51 53 56 58 59 66 67} Structuring the conversation around important values and goals was viewed as differentiating serious illness conversations from other conversations in the care continuum⁴⁵ because this focus oriented the conversation towards how the person wanted to live, not necessarily how they wanted to die.³⁰ Asking about goals could lead clinicians to ask what patients would want if their goals were not within reach.²⁸ The conversation could also focus on soliciting views specifically related to illness, treatment or overall care.^{18 22 30 33 36 38 40 44 48–50 56 68 69} Value was noted in speaking about these topics more generally without necessarily linking it to an illness/health dichotomy.^{29 32} By exploring what was important through personal/clinical and concrete/existential lenses it was possible to gain insight into the patient's experience as a person in order to construct a sensitive and appropriate way forward.^{30 32 37 58 59}

Some serious illness conversations asked about sources of strength^{27 33 37 40 46 47 56 57 63} or prioritised values.⁶⁹ Conversation elements could be added to ask about sources of support,⁵⁶ including family support, coping resources, faith or spirituality.^{45 66 70} The subject of quality of life could likewise be introduced as a separate conversation element (eg, how would you describe your quality of life?), or it might be explored by delving deeper into the patient's goals, worries or priorities.^{32 48 49 55 56 68} If included, questions around critical abilities explored the way that the patient wanted to live (or what they could not live without) by surveying aspects of function, purpose and meaning in the face of potentially worsening health.^{16 27 33 36 40 46 48 49 53 58 63 68 69} Eliciting the patient's perspective regarding the functions and/or activities that were most important to them informed how to best support their needs and autonomy.^{37 60 66 68} Exploring possible trade-offs asked what patients would be willing to go through or concede in relation to, for example, gaining more time or mitigating possible losses.^{27 33 36 37 40 46 49 56 58 63} This prompted reflection and consideration of possible harms, benefits, burdens and risks related to care, as well as evaluation of what was both important and acceptable to the patient.⁵⁸ These questions could be focused around physical or cognitive abilities^{29 30} or

certain goals⁴⁸ to give clinicians insight into acceptable care and treatments.^{15 29 35 37 48}

Serious illness conversations seek to align what patients want in their life and care

Aligning what patients wanted in their life and care involved exploring what was most significant to patients and providing appropriate recommendations and interventions based on these preferences. This theme is comprised of two subthemes: (a) preferences and priorities and (b) preparation and recommendations.

Preferences and priorities

It was important for patients to be able to express their preferences^{4 8 17 20 42} as this could offer clinically significant insights in relation to their overall care.³⁵ However, eliciting preferences pertaining to specific medical treatments was not recommended early in the conversation as the emphasis should be on understanding the patient as a person first and foremost.²⁹ Discussion of preferences and priorities might be dependent on the patient's trajectory and whether decisions needed to be made sooner or later.²³ This involved providing patients with dedicated time and space to ask questions, reason, deliberate and express their preferences in relation to their future.^{8 32 67}

Asking about priorities was often addressed in relation to patients' aforementioned values, goals, wishes, hopes, worries, fears and preferences.^{16 23 32 40 47–49 52 57 69}

A goal was described as a specific want or desire related to a person's values, and a priority designated the importance of one goal or value over another.³⁷ The process of prioritisation was described as asking patients what matters, and then asking them what matters most.⁶⁹ Priorities could be explored in relation to health/illness goals and values, as well as in relation to familial, social or financial needs.⁷¹ Hence, it was important to ask questions to establish the meaning of a prioritised goal or value in the context of the patient's life, and in their own words, to guide decision making and recommendations.^{28 29 42 69}

Preparation and recommendations

Preparation of family, friends, surrogates, caregivers, healthcare proxies and substitute/medical decision makers was another important element.^{40 41 49 53 56} This included how much people in the patient's life knew about the health/illness situation,²² their level of involvement^{33 55 57 63} and whether support persons had (or required) support of their own.³¹ Patients could be asked to think about who they wanted to be their substitute decision maker^{36 41 47 53 60} and prepare that person for involvement in future decision making.^{30 53} Preparation for life events such as financial planning, travel or retirement might also occur.^{32 41} Including family or caregivers in the discussion supported the identification of barriers, prompting timely action and intervention to prevent possible care or discharge

delays.⁷² This was viewed as beneficial to the care partnership as it gave clinicians insight into patient and family preferences and provided family members with insight into care processes.⁷³ This encouraged clinicians to not simply discuss medical events, but to consider the human character of life and illness as part of a 'bigger picture'.^{32 48}

Recommendations for 'next steps'²⁸ could be context specific⁴⁴ or possibly dependent on the scope of practice of the clinician having the conversation.^{38 53} By eliciting aspects of life, health and illness significant to the patient, it was possible to tailor care and treatment plans that balanced the burdens/benefits of various treatment options and reflected the aspects identified as most important.^{16 29 31 33 40 45 58–60 63 67} In this way, recommendations were not simply prescribed, but were opportunities for person-centred shared decision making.^{29 30 67 68} Discussions provided space for clinicians and patients to express their thoughts about continuing, deferring or de-escalating certain care interventions.⁷² While it was not always necessary or possible to make decisions during the conversation, it was important for patients to be prepared to make decisions with a realistic understanding of what was happening, or could happen, with their illness.^{7 29 37 44 59 63} Other care planning matters could also be addressed,^{5 8 53} including life sustaining treatments (ie, intubation, cardiopulmonary resuscitation and tracheostomy).^{4 5 31 38 48 55 60 68} Establishing code status was part of some serious illness conversations,^{19 31 33 47 48 51 68 72} but was thought to only be possible once the patient's values and priorities were known.^{4 14 30} Others emphasised that establishing resuscitation orders should not be the focus of the conversation.^{30 54} Exploring end-of-life options could involve discussion of supportive/comfort care, hospice care, palliative care referrals and practical planning, such as assigning a healthcare proxy or establishing where the patient might like to die.^{8 14 18 19 31 32 43 47 48 50 55 65 66} Personalised preparation and recommendations provided opportunities for care to be proactive rather than reactive.⁶⁰

DISCUSSION

Main findings

This study reviewed literature pertaining to the SICP and SICG and explicated core conversation elements found therein. The three themes and six subthemes synthesise existent understandings, descriptions and interpretations of the core elements of serious illness conversations. The results revealed that the multifaceted nature and content of serious illness conversations could be framed, understood and communicated in numerous ways. While the serious illness conversation construct is relatively recent, this study has shown that its intention and subject matter is being iteratively defined and re-defined as it is adapted and applied in novel and varied contexts.

The original SICG outlined seven conversation components that were designed to support communication with patients who were often anxious due to lack of information about their prognosis or what to expect.⁷ However, this review showed that the conversation content has expanded through various adaptations to include other core elements, such as identifying a substitute or medical decision maker,^{20 30 36 41 45 53} providing clinical information,³⁰ expectations for health in the future,^{24 35 61} current quality of life,^{31 35 38 49} possibility of setbacks,³⁵ assess sources of strength or support,^{18 26 27 30 33 34 37 39 40 44–46 51 52 56–59 63 69} end of life care,^{4 14 19 31 50} code status^{14 19 30 31 33 38} and/or other needs^{14 26 41 50}; or perhaps exclude elements, such as prognosis.^{35 41 53} These changes are also seen in the fluidity of referring to serious illness conversations as advance care planning, end of life planning, goals of care conversations or values and goals conversations. Such alterations appear to reflect efforts to match differing intentions of the conversation, both from clinicians' 'sending' and patients' 'receiving' perspectives.

The patient and clinician-tested language of the SICG has been said to reduce the clinician's cognitive load, while modifications to the guide align information and recommendations based on real-time feedback to match the clinical context.^{31 74} While many changes have been made to the SICG, it is worth noting that the same core elements could be used in many clinical contexts. By discussing goals, values, fears, worries, hopes, desires and wishes, in context, it becomes possible to move beyond medicalisation of the illness experience, and explore the human experience of living with a serious illness.³² In this way, serious illness conversations do not seek to only discuss potential life expectancy, but life expectations as a whole.¹⁸ These results add to the literature supporting the concept of serious illness conversations as being guided by an ethos of person-centred and goal-concordant care.

The issue of prognostication is complicated, and the various modifications made to the guide reflect that inclusion of this element may be dependent on the patient, the method of identification, the clinician's scope of practice, and/or the clinical context.^{15 75} Discussion of prognosis was viewed by some as an indicator of a high-quality serious illness conversation.¹⁵ This is because talking about prognosis openly could help patients with psychological and existential coping mechanisms and the day-to-day reality of living with a serious illness.⁷⁶ Similarly, talking about what the patient experienced to be important could help them to articulate thoughts and opinions surrounding what would be acceptable to them in relation to their life and care. This highlights the importance of establishing expectations and intended outcomes for serious illness conversations in relation to the context in which they are conducted.

While the benefits (or possible drawbacks) of specific conversation domains require further exploration, this review contributes an important inventory and synthesis of existent core conversation elements. These findings pave the way for development of a conceptual framework for serious illness conversations that includes a holistic definition and content explication to further differentiate this activity in the care continuum.^{76 77} Future research could also explore the extent to which various conversation elements contribute to patient/family outcomes and clinician/organisation experiences.^{8 15 76 78}

Strengths and weaknesses

This review used rigorous methods to identify and synthesise literature pertaining to serious illness conversation core elements. Strict inclusion and exclusion criteria were adhered to, and transparent search, extraction, analysis and reporting methods were described. Thematic synthesis facilitated the analysis and inclusion of articles with varied methodologies in diverse clinical settings.

The SICP and SICG were developed by Ariadne Labs, a joint centre for health systems innovation at Brigham and Women's Hospital and the Harvard T.H. Chan School of Public Health. As this study only examined the SICP/SICG, it is likely that these themes reflect the content of the programme and guide in some way. Other serious illness conversation training programmes or guides were not included in this study and may contain other components. Most studies originated from North America, indicating a possible lack of cultural diversity. Due to several large-scale studies and secondary analyses of data, the number of included articles outnumbers the total number of studies. However, these articles were included and analysed individually because studies originating from the same cluster explored and described different aspects of the data.

This review was not limited to study type, participant or context, and included implementation studies as well as discussion articles, so these results combine patient, clinician and researcher descriptions of serious illness conversation elements across different methods and contexts. Further, it is impossible to know how closely clinicians followed the guide, or the extent to which documentation of conversations elements in the literature reflected the actual content of conversations. The authors acknowledge their knowledge of the serious illness conversation subject area and guide content may have impacted the interpretation. The lack of a second independent initial screener of the titles and abstracts is also recognised as a limitation. Two authors in the current study authored several articles included in the current review (JP^{4–8 14–20 51 57 60 70} (n=16, 24.6%) and EKF^{4–6 15–17 57 60} (n=8, 12%)). To minimise bias, JP and EKF were not involved in article selection, data extraction or quality appraisal.

Conclusions

This integrative systematic review explored how core elements of serious illness conversations were described in the literature and presented themes underpinning extant descriptions of these conversation elements. The results offer insights into the core elements of serious illness conversations in the context of the SICP/SICG and may be used to inform current and future clinical education and practice.

Twitter Rebecca Baxter @researchranger and Anna Sandgren @AnnaSandgren70

Contributors All authors made substantial contributions. SP performed the initial searches and screening in consultation with AS and RB. SP and RB conducted the full-text article screening in consultation with SA. SP evaluated the quality appraisals in consultation with RB. Data were extracted by RB, SP and SA. JP and EKF have extensive expertise in the SICP and provided critical contributions to the interpretation and results. All authors contributed to the analysis and construction of the final results. RB wrote the manuscript draft, and all authors critically reviewed, edited and revised the text. RB is the guarantor for the study. All authors approved the final version of the manuscript.

Funding The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the Kamprad Family Foundation for Entrepreneurship, Research, and Charity [grant number 20210163].

Competing interests EKF and JP are both faculty in Ariadne Labs' Serious Illness Care Program and are authors in several of the cited papers.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. The full dataset of included studies is available from the respective publishers.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Rebecca Baxter <http://orcid.org/0000-0001-6595-6298>

Susanna Pusa <http://orcid.org/0000-0002-4773-8796>

Sofia Andersson <http://orcid.org/0000-0002-1728-5722>

Erik K Fromme <http://orcid.org/0000-0002-6731-2338>

Joanna Paladino <http://orcid.org/0000-0001-9324-1943>

Anna Sandgren <http://orcid.org/0000-0002-3155-575X>

REFERENCES

- AriadneLabs. Serious illness care. 2022. Available: <https://www.ariadnelabs.org/serious-illness-care/> [Accessed 10 Sep 2022].
- Bernacki RE, Block SD, American College of Physicians High Value Care Task Force. Communication about serious illness care goals: a review and synthesis of best practices. *JAMA Intern Med* 2014;174:1994–2003.
- Bernacki R, Paladino J, Neville BA, *et al.* Effect of the serious illness care program in outpatient oncology: a cluster randomized clinical trial. *JAMA Intern Med* 2019;179:751–9.
- Paladino J, Bernacki R, Neville BA, *et al.* Evaluating an intervention to improve communication between oncology clinicians and patients with life-limiting cancer: a cluster randomized clinical trial of the serious illness care program. *JAMA Oncol* 2019;5:801–9.
- Geerse OP, Lamas DJ, Sanders JJ, *et al.* A qualitative study of serious illness conversations in patients with advanced cancer. *J Palliat Med* 2019;22:773–81.
- Sanders JJ, Miller K, Desai M, *et al.* Measuring goal-concordant care: results and reflections from secondary analysis of a trial to improve serious illness communication. *J Pain Symptom Manage* 2020;60:889–97.
- Bernacki R, Hutchings M, Vick J, *et al.* Development of the serious illness care program: a randomised controlled trial of a palliative care communication intervention. *BMJ Open* 2015;5:e009032.
- Kumar P, Wixon-Genack J, Kavanagh J, *et al.* Serious illness conversations with outpatient oncology clinicians: understanding the patient experience. *JCO Oncol Pract* 2020;16:e1507–15.
- CambridgeDictionary. Definition of 'core element' 2023. Available: <https://dictionary.cambridge.org/example/english/core-element> [Accessed 1 May 2023].
- Joanna Briggs Institute critical appraisal tools. n.d. Available: <https://jbi.global/critical-appraisal-tools>
- Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol* 2008;8:45.
- Harden A, Thomas J. Methodological issues in combining diverse study types in systematic reviews. *International Journal of Social Research Methodology* 2005;8:257–71.
- Booth A, Noyes J, Flemming K, *et al.* *Guidance on choosing qualitative evidence synthesis methods for use in health technology assessments of complex interventions*. Integrate-HTA, 2016.
- Miranda SP, Bernacki RE, Paladino JM, *et al.* A descriptive analysis of end-of-life conversations with long-term glioblastoma survivors. *Am J Hosp Palliat Care* 2018;35:804–11.
- Geerse OP, Lamas DJ, Bernacki RE, *et al.* Adherence and concordance between serious illness care planning conversations and oncology clinician documentation among patients with advanced cancer. *J Palliat Med* 2021;24:53–62.
- Paladino J, Koritsanszky L, Nisotel L, *et al.* Patient and clinician experience of a serious illness conversation guide in oncology: a descriptive analysis. *Cancer Med* 2020;9:4550–60.
- Paladino J, Koritsanszky L, Neal BJ, *et al.* Effect of the serious illness care program on health care utilization at the end of life for patients with cancer. *J Palliat Med* 2020;23:1365–9.
- Paladino J, Brannen E, Benotti E, *et al.* Implementing serious illness communication processes in primary care: a qualitative study. *Am J Hosp Palliat Care* 2021;38:459–66.
- Lakin JR, Koritsanszky LA, Cunningham R, *et al.* A systematic intervention to improve serious illness communication in primary care. *Health Affairs* 2017;36:1258–64.
- Lakin JR, Benotti E, Paladino J, *et al.* Interprofessional work in serious illness communication in primary care: a qualitative study. *J Palliat Med* 2019;22:751–63.

- 21 Gace D, Sommer RK, Daubman B-R, *et al.* Exploring patients' experience with clinicians who recognize their unmet palliative needs: an inpatient study. *J Palliat Med* 2020;23:1493–9.
- 22 Greenwald JL, Greer JA, Gace D, *et al.* Implementing automated triggers to identify hospitalized patients with possible unmet palliative needs: assessing the impact of this systems approach on clinicians. *J Palliat Med* 2020;23:1500–6.
- 23 Daubman B-R, Bernacki R, Stoltenberg M, *et al.* Best practices for teaching clinicians to use a serious illness conversation guide. *Palliat Med Rep* 2020;1:135–42.
- 24 Jacobsen J, Jackson V, Asfaw S, *et al.* One hospital's response to the Institute of medicine report, "dying in America". *J Pain Symptom Manage* 2022;63:e182–7.
- 25 Manz CR, Parikh RB, Small DS, *et al.* Effect of integrating machine learning mortality estimates with behavioral nudges to clinicians on serious illness conversations among patients with cancer: a stepped-wedge cluster randomized clinical trial. *JAMA Oncol* 2020;6:e204759.
- 26 Li EH, Ferrell W, Klaiman T, *et al.* Impact of behavioral nudges on the quality of serious illness conversations among patients with cancer: secondary analysis of a randomized controlled trial. *JCO Oncol Pract* 2022;18:e495–503.
- 27 Daly JM, Schmidt ME, Thoma KD, *et al.* Trained clinician's documentation of serious illness conversations and use of billing CPT 99497. *J Palliat Care* 2022;37:323–31.
- 28 Daly J, Schmidt M, Thoma K, *et al.* How well are serious illness conversations documented and what are patient and physician perceptions of these conversations *J Palliat Care* 2022;37:332–40.
- 29 Jain N, Bernacki RE. Goals of care conversations in serious illness: a practical guide. *Med Clin North Am* 2020;104:375–89.
- 30 Sirianni G, Torabi S. Addressing serious illness conversations during COVID-19. *Can Fam Physician* 2020;66:533–6.
- 31 Lakin JR, Arnold CG, Catzen HZ, *et al.* Early serious illness communication in hospitalized patients: a study of the implementation of the speaking about goals and expectations (SAGE) program. *Healthcare* 2021;9:100510.
- 32 Borregaard Myrholm C, Novrup Clemmensen S, Sax Røgind S, *et al.* Serious illness conversations in patients with multiple myeloma and their family caregivers: a qualitative interview study. *Eur J Cancer Care (Engl)* 2022;31:e13537.
- 33 Beddard-Huber E, Strachan P, Brown S, *et al.* Supporting Interprofessional engagement in serious illness conversations: an adapted resource. *J Hosp Palliat Nurs* 2021;23:38–45.
- 34 Karim S, Lupichuk S, Tan A, *et al.* Real world implementation of the serious illness care program in cancer care: results of a quality improvement initiative. *J Palliat Med* 2021;24:905–9.
- 35 Lamas DJ, Owens RL, Nace RN, *et al.* Conversations about goals and values are feasible and acceptable in long-term acute care hospitals: a pilot study. *J Palliat Med* 2017;20:710–5.
- 36 Ko JJ, Ballard MS, Shenkier T, *et al.* Serious illness conversation-evaluation exercise: a novel assessment tool for residents leading serious illness conversations. *Palliat Med Rep* 2020;1:280–90.
- 37 Mandel EI, Bernacki RE, Block SD. Serious illness conversations in ESRD. *Clin J Am Soc Nephrol* 2017;12:854–63.
- 38 Aaronson EL, Greenwald JL, Krenzel LR, *et al.* Adapting the serious illness conversation guide for use in the emergency department by social workers. *Pall Supp Care* 2021;19:681–5.
- 39 Pottash M, Joseph L, Rhodes G. Practicing serious illness conversations in graduate medical education. *Med Sci Educ* 2020;30:1187–93.
- 40 Gelfand SL, Mandel EI, Mendu ML, *et al.* Palliative care in the advancing American kidney health initiative: a call for inclusion in kidney care delivery models. *Am J Kidney Dis* 2020;76:877–82.
- 41 Greenwald JL, Abrams AN, Park ER, *et al.* Development of a peer support program for clinicians having serious illness conversations during COVID-19. *J Gen Intern Med* 2021;36:1094–7.
- 42 Karim S, Levine O, Simon J. The serious illness care program in oncology: evidence, real-world implementation and ongoing barriers. *Curr Oncol* 2022;29:1527–36.
- 43 Massmann JA, Revier SS, Ponto J. Implementing the serious illness care program in primary care. *J Hosp Palliat Nurs* 2019;21:291–9.
- 44 Pasricha V, Gorman D, Laothamatas K, *et al.* Use of the serious illness conversation guide to improve communication with surrogates of critically ill patients: a pilot study. *ATS Sch* 2020;1:119–33.
- 45 DeCoursey DD, Partin L, Revette A, *et al.* Development of a stakeholder driven serious illness communication program for advance care planning in children, adolescents, and young adults with serious illness. *J Pediatr* 2021;229:247–58.
- 46 Hafid A, Howard M, Guenter D, *et al.* Advance care planning conversations in primary care: a quality improvement project using the serious illness care program. *BMC Palliat Care* 2021;20:122.
- 47 Le K, Lee J, Desai S, *et al.* The surprise question and serious illness conversations: a pilot study. *Nurs Ethics* 2021;28:1010–25.
- 48 Reed-Guy L, Miranda SP, Alexander TD, *et al.* Serious illness communication practices in glioblastoma: an institutional perspective. *J Palliat Med* 2022;25:234–42.
- 49 Thamcharoen N, Nissaisorakarn P, Cohen RA, *et al.* Serious illness conversations in advanced kidney disease: a mixed-methods implementation study. *BMJ Support Palliat Care* 2021. 10.1136/bmjspcare-2020-002830 [Epub ahead of print 17 Mar 2021].
- 50 O'Donnell AE, Schaefer KG, Stevenson LW, *et al.* Social worker-aided palliative care intervention in high-risk patients with heart failure (SWAP-HF): a pilot randomized clinical trial. *JAMA Cardiol* 2018;3:516–9.
- 51 Ma C, Riehm LE, Bernacki R, *et al.* Quality of clinicians' conversations with patients and families before and after implementation of the serious illness care program in a hospital setting: a retrospective chart review study. *CMAJ Open* 2020;8:E448–54.
- 52 Tam V, You JJ, Bernacki R. Enhancing medical learners' knowledge of, comfort and confidence in holding serious illness conversations. *Am J Hosp Palliat Care* 2019;36:1096–104.
- 53 Lally K, Tuya Fulton A, Ducharme C, *et al.* Using nurse care managers trained in the serious illness conversation guide to increase goals-of-care conversations in an accountable care organization. *J Palliat Med* 2020;23:112–5.
- 54 Lagrotteria A, Swinton M, Simon J, *et al.* Clinicians' perspectives after implementation of the serious illness care program: a qualitative study. *JAMA Netw Open* 2021;4:e2121517.
- 55 King S, Douglas M, Javed S, *et al.* Content of serious illness care conversation documentation is associated with goals of care orders: a quantitative evaluation in hospital. *BMC Palliat Care* 2022;21:116.
- 56 Wasp GT, Cullinan AM, Chamberlin MD, *et al.* Implementation and impact of a serious illness communication training for hematology-oncology fellows. *J Cancer Educ* 2021;36:1325–32.
- 57 Paladino J, Kilpatrick L, O'Connor N, *et al.* Training clinicians in serious illness communication using a structured guide: evaluation of a training program in three health systems. *J Palliat Med* 2020;23:337–45.
- 58 van Breemen C, Johnston J, Carwana M, *et al.* Serious illness conversations in pediatrics: a case review. *Children (Basel)* 2020;7:102.

- 59 Baran CN, Sanders JJ. Communication skills: delivering bad news, conducting a goals of care family meeting, and advance care planning. *Prim Care* 2019;46:353–72.
- 60 Paladino J, Mitchell S, Mohta N, *et al.* Communication tools to support advance care planning and hospital care during the COVID-19 pandemic: a design process. *Jt Comm J Qual Patient Saf* 2021;47:127–36.
- 61 Zehm A, Scott E, Schaefer KG, *et al.* Improving serious illness communication: testing the serious illness care program with trainees. *J Pain Symptom Manage* 2022;63:e252–9.
- 62 Bowman JK, Aaronson EL, Petrillo LA, *et al.* Goals of care conversations documented by an embedded emergency department-palliative care team during COVID. *J Palliat Med* 2023;26:662–6.
- 63 McGlinchey T, Mason S, Coackley A, *et al.* Serious illness care programme UK: assessing the 'face validity', applicability and relevance of the serious illness conversation guide for use within the UK health care setting. *BMC Health Serv Res* 2019;19:384.
- 64 LoCastro M, Sanapala C, Mendler JH, *et al.* Adaptation of serious illness care program to be delivered via Telehealth for older patients with hematologic malignancy. *Blood Adv* 2023;7:1871–84.
- 65 Garcia R, Brown-Johnson C, Teuteberg W, *et al.* The team-based serious illness care program: a qualitative evaluation of implementation and teaming. *J Pain Symptom Manage* 2023;65:521–31.
- 66 Davoudi A, Tissot H, Doucette A, *et al.* Using natural language processing to classify serious illness communication with oncology patients. *Health Informatics* [Preprint] 2022.
- 67 Xu L, Sommer RK, Nyeko L, *et al.* Patient perspectives on serious illness conversations in primary care. *J Palliat Med* 2022;25:940–4.
- 68 Ouchi K, Lawton AJ, Bowman J, *et al.* Managing code status conversations for seriously ill older adults in respiratory failure. *Ann Emerg Med* 2020;76:751–6.
- 69 Moye J, Driver JA, Owsiany MT, *et al.* Assessing what matters most in older adults with multi-complexity. *Gerontologist* 2022;62:e224–34.
- 70 Sanders JJ, Durieux BN, Cannady K, *et al.* Acceptability of a serious illness conversation guide to black Americans: results from a focus group and oncology pilot study. *Palliat Support Care* 2022:1–10.
- 71 Swiderski D, Georgia A, Chuang E, *et al.* "I was not able to keep myself away from tending to her immediate needs": primary care physicians' perspectives of serious illness conversations at community health centers. *J Gen Intern Med* 2022;37:130–6.
- 72 Hu FY, O'Mara L, Tulebaev S, *et al.* Geriatric surgical service interventions in older emergency general surgery patients: preliminary results. *J Am Geriatr Soc* 2022;70:2404–14.
- 73 Wasp GT, Cullinan AM, Anton CP, *et al.* Interdisciplinary approach and patient/family partners to improve serious illness conversations in outpatient oncology. *JCO Oncology Practice* 2022;18:e1567–73.
- 74 Smith GM, Radigan NJ, Maloney FL, *et al.* Development, implementation, and outcomes of a serious illness care community of practice. *J Pain Symptom Manage* 2022;63:e160–7.
- 75 Baxter R, Fromme EK, Sandgren A. Patient identification for serious illness conversations: a scoping review. *Int J Environ Res Public Health* 2022;19:4162.
- 76 Jacobsen J, Bernacki R, Paladino J. Shifting to serious illness communication. *JAMA* 2022;327:321–2.
- 77 Sanders JJ, Paladino J, Reaves E, *et al.* Quality measurement of serious illness communication: recommendations for health systems based on findings from a symposium of national experts. *J Palliat Med* 2020;23:13–21.
- 78 Andersson S, Sandgren A. Organizational readiness to implement the serious illness care program in hospital settings in Sweden. *BMC Health Serv Res* 2022;22:539.