

Heart failure symptom burden in outpatient cardiology: observational cohort study

Maria Walsh ¹, Elizabeth Bowen,¹ Carl Vaughan,² Fiona Kiely¹

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/spcare-2023-004167>).

¹Department of Palliative Medicine, Marymount University Hospital and Hospice, Cork, Ireland

²Department of Cardiology, Mercy University Hospital, Cork, Ireland

Correspondence to

Dr Maria Walsh, Palliative Medicine, Marymount University Hospital and Hospice, T12 A710 Cork, Ireland; mariacatherine.walsh@gmail.com

This study was presented at the Annual Irish Association for Palliative Care Education and Research Seminar, February 2022.

Received 12 January 2023

Accepted 23 March 2023

Published Online First

19 April 2023



© Author(s) (or their employer(s)) 2023. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Walsh M, Bowen E, Vaughan C, et al. *BMJ Supportive & Palliative Care* 2023;**13**:e1280–e1284.

ABSTRACT

Objectives To assess the self-reported symptom burden in patients with a diagnosis of heart failure attending an outpatient cardiology clinic through the utilisation of validated patient-reported outcome measures.

Methods Eligible patients were invited to partake in this observational cohort study. Participant demographics and comorbidities were recorded, followed by participants recording their symptoms using the Integrated Palliative care Outcome Scale (IPOS) and Brief Pain Inventory (BPI) outcome measure tools.

Results A total of 22 patients were included in the study. The majority were male (n=15). The median age was 74.5 (range 55–94) years. Atrial fibrillation and hypertension were the most common comorbidities (n=10). Dyspnoea, weakness and poor mobility were the most prevalent symptoms, affecting 15 (68%) of the 22 patients. Dyspnoea was reported as being the most troublesome symptom. The BPI was completed by 68% (n=15) of the study participants. Median average pain score was 5/10; median worst pain score in the preceding 24 hours was 6/10 and median pain score at time of BPI completion was 3/10. The impact of pain on daily living during the preceding 24 hours ranged from impacting on all activities (n=7) to not impacting on activities (n=1).

Conclusions Patients with heart failure experience a range of symptoms that vary in severity. Introduction of a symptom assessment tool in the cardiology outpatient setting could help identify patients with a high symptom burden and prompt timely referral to specialist palliative care services.

INTRODUCTION

Heart failure (HF) is a progressive, life-limiting condition from structural and/or functional cardiac abnormalities along with objective evidence of pulmonary or systemic congestion.¹ Most recent data available from Ireland estimate 90 000

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Patients with advanced heart failure experience a range of symptoms impacting on quality of life.

WHAT THIS STUDY ADDS

⇒ Symptoms can be captured by using patient-reported outcome measures (PROM).
⇒ PROM tools are accessible in the outpatient setting.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Symptom assessment tools in the cardiology outpatient setting could ensure timely referral to specialist palliative care services.

Irish people [2% of the population] are living with HF.² An American study estimates the number will rise to >8 million in the USA by 2030.³ Globally, HF affects >64 million people; a global pandemic.⁴ Persons with HF can experience significant symptom burden with a high rate of hospital admission⁵ and lower quality of life.⁶

Palliative care is focused on improved quality of life in life-limiting illness. Early referral to palliative care services prevents and relieves suffering⁷ with clear health gains.^{8 9} Randomised controlled trials reveal improved symptom control, quality of life and mood.^{7 10} WHO estimates that up to one-third of adults diagnosed with cardiovascular disease would benefit from palliative care. The American Heart Association guidelines recommend palliative care be integrated into the care of all patients diagnosed with advanced cardiovascular disease.¹¹ A recent position paper from the Heart Failure Association of the European Society of Cardiology advised that all patients with HF be considered for palliative care throughout the disease

trajectory.¹² Despite this, there remains a shortfall in timely referral of end-stage HF to palliative care services, with most patients referred too late to maximise the benefits.¹³

Many symptoms have been reported by those living with advanced HF.^{6 14 15} Pain is an under-reported and undertreated symptom.¹⁶ The European Association for Palliative Care Task Force expert position statement of 2020 highlighted the need for symptoms to be assessed in a systematic fashion to identify those requiring specialist palliative care.¹⁷ The use of validated assessment tools is recommended. These can identify uncontrolled or escalating symptoms that should trigger a referral.

The aim of this study was to assess self-reported symptom burden in patients with a diagnosis of HF attending an outpatient cardiology clinic. Objectives included quantifying symptom burden and assessing pain scores through validated patient-reported outcome measures (PROM), specifically the Integrated Palliative care Outcome Scale (IPOS) and Brief Pain Inventory (BPI).

METHODS

Patients attending routine cardiology outpatient appointments at an acute general hospital were invited to participate in this observational cohort study. Inclusion criteria consisted of adult patients (>18 years of age) with a formal diagnosis of HF made by a consultant cardiologist. Eligible patients were identified by co-investigators (MW and EB) after review of medical charts—those without a diagnosis of HF were excluded. An information sheet was distributed to all patients suitable for study participation, and informed consent obtained (by MW or EB) before data collection. Patient demographics and comorbidities were recorded, with all data pseudoanonymised and stored on a password-protected laptop in compliance with General Data Protection Regulation standards.

The IPOS is a 20-part patient questionnaire about physical, psychological, emotional, spiritual and support needs (online supplemental appendix 2). Question 1 invites participants to highlight their main problems or concerns in an open format, focusing on their experiences over the preceding 3 days. The remainder of the questions use a Likert scale—ranging from ‘not at all’ to ‘overwhelmingly’—to measure the impact of specific symptoms on participants. Participant and family distress and practical concerns were also assessed. The feasibility and acceptability of the IPOS in a HF outpatient clinic has been successfully demonstrated.¹⁸ The BPI (Short Form) assesses severity and pain interference with activities of daily living (online supplemental appendix 1). Participants are asked to rate their pain severity at its worst, least and average during the previous 24 hours, and the time of questionnaire completion; 0 represents no pain and 10 the worst pain imaginable. Subsequent questions measure the impact of pain on daily function and quality of life on a similar 11-point scale, with 0 as pain that does not interfere and 10 complete interference. The BPI has been validated in both non-malignant and malignant pain.¹⁹

Patients completed the IPOS and BPI while waiting to be reviewed in the cardiology outpatient clinic. Those who could not complete the questionnaires independently were assisted by family members or a member of the research team. Data were collected over a 3-month period to facilitate a sample size of greater than 20, in alignment with a prior feasibility study within a similar population and context.¹⁸

RESULTS

A total of 22 patients were included. The majority were male (n=15). The median age was 74.5 (range 55–94) years. Atrial fibrillation and hypertension were the most common comorbidities (n=10). Sixty-seven per cent had three or more comorbidities (figure 1).




Figure 1 Patient comorbidities. HTN, hypertension; IHD, ischaemic heart disease; T2DM, type 2 diabetes mellitus.

Patients were at various disease stages, and New York Heart Association classification was inconsistently captured.

All participants completed the IPOS. Dyspnoea, weakness and poor mobility were the most prevalent symptoms, affecting 15 (68%) of the 22. Dyspnoea was the most troublesome symptom, with over 55% moderately and 13% severely overwhelmingly affected. Most of those affected by weakness/lack of energy described it as having a slight-moderate impact; similarly, 55% of patients affected by poor mobility described it as a slight-moderate effect. Participants who responded to the question ‘what have been your main problems or concerns over the past 3 days’ recorded dyspnoea (n=6), pain (n=2) and a range of other symptoms including oedema and altered bowel pattern. A third (36%) of participants did not answer this question.

Over two-thirds felt anxious or worried about their health (n=16) and a similar number reported family concerns (n=15). Most patients (77%) felt at peace most of the time or always, and only three felt depressed. Most (n=15) felt able to share how they were feeling with their family or friends; however, two did not feel able to share their feelings at all. Regarding information sharing, 77% of participants had as much information as they wanted (most of the time or always) and 82% described practical problems to their illness as addressed mostly or all of the time.

The BPI was completed by 68% (n=15), five of whom denied any pain on the day of study. Analgesics identified via a free-text question included paracetamol (n=3) and duloxetine (n=1). Median average pain score was 5/10; median worst pain score in the prior 24 hours was 6/10 and median pain score by BPI was 3/10. The impact of pain on daily living during the preceding 24 hours ranged from all activities (n=7) to none (n=1) (see figure 2).

DISCUSSION

There is value in using PROM to determine symptom burden in advanced HF given the inherent subjectivity of all symptoms. IPOS revealed that dyspnoea is one of

the most common symptoms and substantially impacts patients’ lives. Those experiencing shortness of breath often describe associated anxiety that can have a negative effect on quality of life,²⁰ and breathlessness is a leading cause of presentation to the emergency department.²¹ Poor mobility and weakness were highlighted as the other most frequent symptoms, both of which can be exacerbated by dyspnoea. Specialist palliative care teams have dyspnoea management expertise.^{22–24} There is significant scope for optimisation of breathlessness with appropriate and timely referral.

Almost half of the patients described pain. Further evaluation of causative factors is indicated—most patients with end-stage HF are elderly with multiple comorbidities.^{16 25} Pain management in life-limiting conditions can be challenging, particularly when pain encompasses all of a person’s physical, psychological, social, spiritual and practical struggles²⁶ and requires a multidimensional approach. Palliative care clinicians have a pivotal role in the education and support of non-specialist colleagues managing complex pain in this setting.

Strengths of our study include the use of PROM tools validated in both palliative care and HF populations. IPOS was deemed to be a feasible and acceptable method of identifying symptoms in HF.¹⁸ While the BPI was initially used to assess pain in cancer it has since been used in various populations with non-malignant diagnoses.¹⁹ The tools are accessible—patients were invited to participate in the waiting room of a busy clinic. Another potential method of administration would be their inclusion with appointment letters, so patients could complete the questions at home before medical review. Availability of PROM tool results at the review would allow focused interventions and help monitor the impact of symptoms on patients’ lives longitudinally.

This was a single-site study with a small number of participants, which limits generalisability. Patients were recruited from a general cardiology clinic rather than a dedicated HF clinic, with small numbers of eligible patients. Fifty per cent of participants completed the




Figure 2 Impact of pain on patient’s daily living.

questionnaires independently, giving the most accurate reports of symptom burden. Previous studies demonstrated that families tend to over-report patient symptoms when members are involved in completing symptom assessment scales, while in contrast health-care professionals may under-report symptom impact on patients.^{27–29}

CONCLUSIONS

Patients with HF experience a range of symptoms that vary in severity. Given the varied disease trajectory and uncertain prognosis in advanced HF brief interventions by the specialist palliative care team may be most beneficial if made in a timely manner, depending on need and patient wishes. Introduction of a symptom assessment tool in the cardiology outpatient clinic could help identify those with a high symptom burden and prompt timely referral to specialist palliative care.

Twitter Maria Walsh @drmariawalsh

Contributors All authors contributed to the study. MW, CV and FK conceived and developed the study design. MW and EB carried out the data collection. MW and FK analysed and interpreted the results and all authors were involved in draft manuscript preparation. All authors have read and approved the manuscript, and consented to manuscript submission and to subsequent publication. MW acted as guarantor for the study.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by the Clinical Research Ethics Committee of the Cork Teaching Hospitals (CREC), Cork, Ireland (Ref No ECM 4 (ccc) 12/02/19). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; internally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information.

Author note In 2022 the IAPC hosted its 21st Education and Research Seminar which was a virtual event given Covid19 restrictions. The IAPC have teamed with BMJ SPC for the last two years where the winner of the platform presentation is invited to submit their research and have it published by BMJ SPC. This is a prestigious award and we are grateful to Dr. Declan Walsh for continuing to support this initiative. The presentation winner Dr. Maria Walsh.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

This article is made freely available for personal use in accordance with BMJ's website terms and conditions for

the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

ORCID iD

Maria Walsh <http://orcid.org/0000-0002-4932-9742>

REFERENCES

- 1 Bozkurt B, Coats AJ, Tsutsui H, *et al.* Universal definition and classification of heart failure: a report of the heart failure society of america, heart failure association of the european society of cardiology, japanese heart failure society and writing committee of the universal definition of heart failure. *J Card Fail* 2021;27:387–413.
- 2 Heart Failure - HSE.ie. HSE.ie. 2021. Available: <https://www.hse.ie/eng/health/hl/living/heartfailure/>
- 3 Heidenreich PA, Albert NM, Allen LA, *et al.* Forecasting the impact of heart failure in the united states. *Circ: Heart Failure* 2013;6:606–19.
- 4 Savarese G, Becher PM, Lund LH, *et al.* Global burden of heart failure: a comprehensive and updated review of epidemiology. *Cardiovasc Res* 2023;118:3272–87. 10.1093/cvr/cvac013 Available: <https://doi.org/10.1093/cvr/cvac013>
- 5 Chamberlain AM, Dunlay SM, Gerber Y, *et al.* Burden and timing of hospitalizations in heart failure: a community study. *Mayo Clin Proc* 2017;92:184–92.
- 6 Alpert CM, Smith MA, Hummel SL, *et al.* Symptom burden in heart failure: assessment, impact on outcomes, and management. *Heart Fail Rev* 2017;22:25–39.
- 7 Temel JS, Greer JA, Muzikansky A, *et al.* Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 2010;363:733–42.
- 8 Kavalieratos D, Corbelli J, Zhang D, *et al.* Association between palliative care and patient and caregiver outcomes. *JAMA* 2016;316:2104.
- 9 Kaasa S, Loge JH, Apapro M, *et al.* Integration of oncology and palliative care: a lancet oncology commission. *Lancet Oncol* 2018;19:e588–653.
- 10 Zimmermann C, Swami N, Krzyzanowska M, *et al.* Early palliative care for patients with advanced cancer: a cluster-randomised controlled trial. *Lancet* 2014;383:1721–30.
- 11 Braun LT, Grady KL, Kutner JS, *et al.* Palliative care and cardiovascular disease and stroke: a policy statement from the american heart association/american stroke association. *Circulation* 2016;134:e198–225.
- 12 Hill L, Prager Geller T, Baruah R, *et al.* Integration of a palliative approach into heart failure care: a European Society of cardiology heart failure association position paper. *Eur J Heart Fail* 2020;22:2327–39.
- 13 Rogers J, Patel C, Mentz R, *et al.* Palliative care in heart failure: the PAL-HF randomized. *Controlled Clinical Trial Journal of the American College of Cardiology* 2017;70:331–41.
- 14 Zambroski CH, Moser DK, Bhat G, *et al.* Impact of symptom prevalence and symptom burden on quality of life in patients with heart failure. *Eur J Cardiovasc Nurs* 2005;4:198–206.
- 15 Nordgren L, Sörensen S. Symptoms experienced in the last six months of life in patients with end-stage heart failure. *Eur J Cardiovasc Nurs* 2003;2:213–7.
- 16 Light-McGroary K, Goodlin SJ. The challenges of understanding and managing pain in the heart failure patient. *Curr Opin Support Palliat Care* 2013;7:14–20.
- 17 Sobanski PZ, Alt-Epping B, Currow DC, *et al.* Palliative care for people living with heart failure: European association for palliative care Task force expert position statement. *Cardiovasc Res* 2020;116:12–27.

- 18 Kane PM, Daveson BA, Ryan K, *et al.* Feasibility and acceptability of a patient-reported outcome intervention in chronic heart failure. *BMJ Support Palliat Care* 2017;7:470–9.
- 19 Keller S, Bann CM, Dodd SL, *et al.* Validity of the brief pain inventory for use in documenting the outcomes of patients with noncancer pain. *Clin J Pain* 2004;20:309–18.
- 20 Carrel H. *Phenomenology of illness*. Oxford: Oxford University Press, 2016.
- 21 Hutchinson A, Pickering A, Williams P, *et al.* Breathlessness and presentation to the emergency department: a survey and clinical record review. *BMC Pulm Med* 2017;17:53.
- 22 Higginson IJ, Bausewein C, Reilly CC, *et al.* An integrated palliative and respiratory care service for patients with advanced disease and refractory breathlessness: a randomised controlled trial. *Lancet Respir Med* 2014;2:979–87.
- 23 Chin C, Booth S. Managing breathlessness: a palliative care approach. *Postgrad Med J* 2016;92:393–400.
- 24 Booth S, Moffat C, Burkin J, *et al.* Nonpharmacological interventions for breathlessness. *Curr Opin Support Palliat Care* 2011;5:77–86.
- 25 Triposkiadis F, Giamouzis G, Parissis J, *et al.* Reframing the association and significance of co-morbidities in heart failure. *Eur J Heart Fail* 2016;18:744–58. 10.1002/ejhf.600 Available: <http://doi.wiley.com/10.1002/ejhf.2016.18.issue-7>
- 26 Richmond C, Dame cicely Saunders. *BMJ* 2005;331:238.
- 27 Silveira MJ, Given CW, Given B, *et al.* Patient-Caregiver concordance in symptom assessment and improvement in outcomes for patients undergoing cancer chemotherapy. *Chronic Illn* 2010;6:46–56.
- 28 Nekolaichuk CL, Bruera E, Spachynski K, *et al.* A comparison of patient and proxy symptom assessments in advanced cancer patients. *Palliat Med* 1999;13:311–23.
- 29 Grossman SA, Sheidler VR, Swedeen K, *et al.* Correlation of patient and caregiver ratings of cancer pain. *J Pain Symptom Manage* 1991;6:53–7.