P 127

A RETROSPECTIVE REVIEW OF ANTIMICROBIAL PRESCRIBING AND INFECTION PREVALENCE IN A PALLIATIVE CARE UNIT

Jill McKane, Susan Addie, Mairi-clare McGowan. *St Vincent's Hospice, Howwood, UK*

10.1136/bmjspcare-2014-000654.168

Introduction Hospice patients are highly susceptible to infections of the urinary and respiratory tract.^{1 2 3 4} The use of antimicrobials for symptom control should be considered against the burden of treatment for the patient as well as antimicrobial resistance.^{1 3 5 6 7}

The Scottish Antimicrobial Prescribing Group produce guidance and targets for hospital based empirical prescribing. The aim is to reduce clostridium difficile infection.⁷ Empirical prescribing standards state 95% of antimicrobials prescribed should be in line with local policy with rationale for use documented.⁷ These hospital targets are applicable to the hospice setting.⁸

Method A consecutive retrospective case note audit was completed for a four month period at St Vincent's Hospice in Howwood. An audit proforma with data collection parameters was developed during this process.

Results Thirty-three case notes were audited: 76.7% of patients received at least one course of antimicrobials during their admission. The main indications for antibiotics were urinary tract(34%) and lower respiratory tract infections (60%). Indications for prescribing were specified in 88% and duration specified in 68% of cases. Of these 38% completed the course, 35% were continued longer and 26% were discontinued early. Antimicrobial selection, dosing and route were in line with the Greater Glasgow and Clyde policy in 78% of prescriptions; however only 51% also included documentation of duration. Of these prescriptions 91% were prescribed empirically without microbiology culture data.

Discussion The prevalence of infection rate, antibiotic use and improvement of symptom burden is variable within the literature.^{1 2 3 4 5} There is scope to improve practice surrounding antimicrobial initiation and documentation. Existing local guidelines are not specific to hospice care and should be adapted. We recommend and are working towards:

- Palliative Antimicrobial Guideline development and liaison with local antimicrobial stewards.
 - Development of St Vincent's Hospice antimicrobial toolkit with inclusion of medical handover tools.
- Re-audit and education.

REFERENCES

- 1 Raquel E, Reinbolt BS, Allison M, *et al*. Symptomatic treatment of infection in patients with advanced cancer receiving palliative care. *J Pain Symptom Manage* 2005;175:175–182.
- 2 White P, Kuhlenschmidt H, Vanvura B, Navari RM. Antimicrobial use in patients with advanced cancer receiving hospice care. *J Pain Symptom Manage* 2003;25:438–43.
- 3 Pereira J, *et al*. A retrospective review of the frequency of infections and patterns of antibiotic utilization on a palliative care unit. *J Pain Symptom Manage* 1998;16:374–81.
- 4 Vitetta L, Kenner D, Sali A. Bacterial infections in terminally ill hospice patients. *J Pain Symptom Manage* 2000;20:326–334.
- 5 Clayton J, Fardell B, Hutton-Potts J, *et al*. Parental antibiotics in palliative care unit: prospective analysis of current practice. *J Pall Med* 2003;17:44–48.
- 6 Van der Steen J, Radbruch L, Hertogh CMPM. White paper defining optimal palliative care in older people with dementia: A Delphi study and recommendations from the European Association for Palliative Care. *Pall Med* Published Online First: 4 Jul 2013. doi:10.1177/0269216313493685
- 7 Scottish Government - Scottish Medicines Consortium - Scottish Antimicrobial Prescribing Group CDI HEAT Target (CEL April 2009) Hospital-based Empirical Prescribing National Report, April 2011.
- 8 Scottish Government - Scottish Medicines Consortium - Scottish Antimicrobial Prescribing Group Good Practice Recommendations for antimicrobial use in frail elderly patients in NHS Scotland, February 2013.