




OPEN ACCESS

Acupuncture for palliative cancer pain management: systematic review

Juan Yang ,¹ Dietlind L Wahner-Roedler,¹ Xuan Zhou,² Lesley A Johnson,³ Alex Do,¹ Deirdre R Pachman,⁴ Tony Y Chon,¹ Manisha Salinas,⁵ Denise Millstine,³ Brent A Bauer¹

¹Division of General Internal Medicine, Mayo Clinic, Rochester, Minnesota, USA

²Formula-pattern Research Center, School of Traditional Chinese Medicine, Jinan University, Guangzhou, China

³Department of Women's Health/Internal Medicine, Mayo Clinic, Scottsdale, Arizona, USA

⁴Center for Palliative Medicine, Mayo Clinic, Rochester, Minnesota, USA

⁵Division of Family Medicine, Mayo Clinic, Jacksonville, Florida, USA

Correspondence to

Dr Brent A Bauer, Division of General Internal Medicine, Mayo Clinic, Rochester, Minnesota, USA 55905; Bauer.Brent@mayo.edu

DM and BAB are joint senior authors.

Received 13 August 2020
Revised 11 December 2020
Accepted 18 December 2020
Published Online First
13 January 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Yang J, Wahner-Roedler DL, Zhou X, et al. *BMJ Supportive & Palliative Care* 2021;**11**:264–270.

ABSTRACT

Background Pain is one of the most common and problematic symptoms encountered by patients with cancer. Due to the multifactorial aetiology, pain management of these patients frequently requires multidisciplinary interventions including conventional support and specialty palliative care. Acupuncture has been identified as a possible adjunctive therapy for symptom management in cancer pain, and there is currently no systematic review focused solely on the evidence of acupuncture on cancer pain in palliative care.

Objective To critically analyse currently available publications regarding the use of acupuncture for pain management among patients with cancer in palliative care settings.

Methods Multiple academic databases were searched from inception to 29 October 2020. Randomised controlled trials involving acupuncture in palliative care for treatment of cancer-related pain were synthesised. Data were extracted by two independent reviewers, and methodological quality of each included study was assessed using the Oxford Centre for Evidence-Based Medicine (OCEBM) 2011 Levels of Evidence.

Results Five studies (n=189) were included in this systematic review. Results indicated a favourable effect of acupuncture on pain relief in palliative care for patients with cancer. According to OCEBM 2011 Levels of Evidence, they were level 2 in one case (20%), level 3 in two cases (40%) and level 4 in the remaining (40%). Low-level evidence adversely affects the reliability of findings.

Conclusions Acupuncture may be an effective and safe treatment associated with pain reduction in the palliative care of patients with cancer. Further high-quality, adequately powered studies are needed in the future.

INTRODUCTION

Cancer is the second highest cause of death globally, with rates rising rapidly

due to an ageing population and widespread adoption of unhealthy lifestyles.^{1 2} Pain is one of the most frequent and problematic symptoms in cancer, with an incidence rate of 50.7% for all cancer stages, and 66.4% for advanced stages.³ It is also one of the most feared and burdensome issues of patients with cancer and interferes with significant aspects of their lives, including daily activity, social function, sleep quality and cognitive function.⁴ Pain can be caused by cancer and/or cancer-related treatments such as surgery, chemotherapy, radiation and hormone therapy.^{5 6} Uncontrolled pain can negatively impact the ability to complete activities of daily living, to maintain employment, to tolerate disease-directed therapy, and can impact emotional and psychological health. It is predicted that the number of cancer patients/survivors will reach over 20 million by 2026.⁷ Therefore, the population suffering from cancer-related pain will also increase. Barriers to optimal pain management in patients with cancer include medical providers' lack of training or comfort with pain management, time constraints in busy oncology practices, concerns of medication side effects or addiction, reimbursement challenges and problems of accessing prescriptions. These barriers remain a major persistent public health challenge, calling for optimised treatment regimens.⁸

Due to the multifactorial and complex nature of cancer pain, treatment should be based on a multidisciplinary collaboration that includes opioid and non-opioid medications, surgical and minimal invasive interventions, psychotherapies, manipulative therapies and integrative medicine (formerly referred to as complementary medicine treatments).⁹ Opioid analgesics are the mainstay for cancer pain, yet up to two-thirds of patients with cancer report inadequate pain management and/or

opioid-induced adverse events (AEs). Opioid analgesics may have unintended side effects including nausea, vomiting, constipation, sedation and cognitive impairment. Ten to twenty per cent of individuals cannot tolerate these AEs necessitating changes in approach to treatment.¹⁰ There is also an increased risk of addiction with long-term use of opioid medications, and many patients are reluctant to use these medications, in spite of their pain-relieving benefits.

Pain is the most common symptom among patients with cancer referred to palliative care.¹¹ Safe and effective management for patients with cancer requires interdisciplinary care, including conventional support and specialty palliative care.¹² Palliative care refers both to the active total care of patients with advanced, incurable disease, including preventing and relieving symptoms as well as improving the quality of life alongside curative treatment.¹³ The involvement of specialty palliative care services for patients with cancer can improve advanced care planning, quality of life (QOL), as well as patient and family satisfaction.^{14 15} There is an increasing number of symptom management techniques that are being offered to patients with cancer, with a lack of strong evidence about best modalities, including consideration for cost-effectiveness and sustainability.¹⁶

As pharmacological agents often are associated with side effects or potential concerns for drug interactions, other non-pharmacological interventions are especially important in the population with cancer. Ideally, these interventions would be effective, feasible and inexpensive. Acupuncture is one of the major modalities in integrative medicine. Having been practised for thousands of years, acupuncture is a traditional Chinese medical intervention involving the insertion of fine, metallic sterile needles into anatomical locations based on a system of meridians and channels theory.¹⁷ It has been considered an effective adjunct to conventional pharmacological treatments for many cancer-related symptoms, as well as medical conditions.¹⁸

Acupuncture has long been used for the treatment of pain, and substantial evidence supports acupuncture is effective in pain management.¹⁹⁻²¹ Although the exact analgesic mechanism is uncertain, it is speculated that the analgesic effect of acupuncture may be mediated by the release of opioid peptides and serotonin.²² In the past decade, acupuncture has been actively used for relieving cancer pain, as well as decreasing analgesic medication doses and side effects.^{23 24} Acupuncture offers a promising yet understudied approach to control the pain experienced by many patients with cancer.²⁵ Acupuncture may be a safer approach to cancer-related pain, and could have synergy with conventional interventions by reducing pain, improving fatigue and improving QOL of patients receiving palliative care.²⁶ There is additional evidence suggesting that acupuncture has a role in the supportive care of patients with cancer by relieving

chemotherapy-induced or radiotherapy-induced nausea and vomiting, xerostomia and leucopenia.²⁷ This has already been recommended in Clinical Practice Guideline through expert groups in oncology.^{28 29}

A number of systematic reviews and meta-analyses have examined acupuncture for pain relief in patients with cancer. While previous findings have suggested that acupuncture has positive results in reducing cancer-related pain, studies inadequately addressed the issue of pain management in palliative care settings.³⁰⁻³⁵ The impact of acupuncture on cancer pain in palliative care was sparsely addressed in reviews of acupuncture on pain improvement, though reviews have included other symptoms such as fatigue, nausea, depression, chemotherapy-induced or radiotherapy-induced side effects for palliative care in patients with cancer.^{18 36-38}

To our knowledge, this current systematic review is the first to address the use of acupuncture to treat cancer pain in palliative care. Therefore, the purpose of this review is to critically assess available publications regarding the use of acupuncture as palliative care for pain management in patients with cancer.

METHODS

Inclusion and exclusion criteria

Studies were included if they met all of the following requirements: (1) participants ≥ 18 years old; (2) participants diagnosed with cancer; (3) received acupuncture therapy; (4) palliative care setting; (5) pain intensity assessed via validated measurement. We excluded studies that were not in English, research involving animals, cellular level or studies not relating to pain management.

Data sources and search strategies

A comprehensive electronic search of several databases was conducted from inception to 30 October 2020. The databases included Ovid MEDLINE(R) and Epub Ahead of Print, In-Process and Other Non-Indexed Citations and Daily, Ovid Embase, Ovid Cochrane Database of Systematic Reviews, Ovid Cochrane Central Register of Controlled Trials and Scopus.

The search strategy was designed and performed by an experienced librarian in Rochester Campus, Mayo Clinic with input from the investigators. Controlled vocabulary supplemented with keywords was used to search for studies describing acupuncture for palliative care in pain management of patients with cancer.

Study selection and data extraction

All the articles were initially selected based on their titles and abstracts. Eligibility criteria were applied to the full-text articles during the final selection. Predefined data, including the first author, publication year, country location, condition, target population, sample size, interventions, outcome, efficacy, duration, follow-up and AEs were extracted. Selection and extraction were independently completed

by two investigators. Discrepancies in data selected and extracted by the investigators were resolved either through discussion with a third author or by resorting to data obtained from the original author. The whole process of study selection is summarised in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram.³⁹

Study appraisal

Quality of evidence was rated according to the Oxford Centre for Evidence-Based Medicine.⁴⁰ Levels of Evidence evaluated included: level 1, randomised trials or systematic reviews of randomised trials; level 2, randomised trial or (exceptionally) observational study with dramatic effect; level 3, non-randomised controlled cohort/follow-up study; level 4, case series, case-control study or historically controlled studies; and level 5, mechanism-based reasoning. Level 1 represented the likely strongest evidence, and level 5 represented the likely weakest evidence.

RESULTS

A total of 941 publications from the electronic databases of Ovid MEDLINE(R), Epub Ahead of Print, Embase, In-Process and Other Non-Indexed Citations and Ovid MEDLINE(R) Daily, EBM Reviews—Cochrane Database of Systematic Reviews, EBM

Reviews—Cochrane Central Register of Controlled Trials using a cut-off date of 29 October 2020, and 49 from Google Scholar were reviewed in this study. Of the original 990 reviewed articles, 17 repetitious studies were excluded. Further screening excluded non-acupuncture, non-palliative, non-cancer, non-pain related, non-English and abstracted publications by examining their titles and abstracts. At the end of this screening process, 26 possible relevant studies remained. Based on selection criteria, five articles relating to the use of acupuncture for pain management in patients with cancer in palliative care settings were included in this study. Details of study selection are presented in the PRISMA flow chart (figure 1).

Five studies with 189 participants were included in this review.^{41–45} Reviewed articles were published between 1999 and 2019, with three studies performed in the USA,^{41 43 45} one in the UK⁴⁴ and one in Canada.⁴² Two studies were retrospective chart review,^{41 43} and the other three were a pilot randomised controlled trial (RCT),⁴² a pilot prospective trial⁴⁵ and a non-randomised cohort study.⁴⁴ All included participants were adults. Sample sizes ranged from 20 to 68 cases. The observed interventions varied with traditional acupuncture,^{41 43} electroacupuncture,⁴² percutaneous electrical nerve stimulation (PENS), Korean




Figure 1 Flowchart of the selection of clinical studies.

Table 1 Summary of demographic information of the included studies

Author/year	Country	Type	Age (years)	N	Indication	Intervention	Assessment tool	Treatment sessions	Provider	Efficacy	Adverse events	Evidence level*
Miller <i>et al</i> 2019 ⁴³	USA	Retrospective chart review	31–89	68	Cancer-related pain	Traditional acupuncture/Seirin filiform needles	Edmonton Symptom Assessment Scale (ESAS)	15–40 min/ treatment , median number of treatments was 2 (range 1–13)	Palliative medicine physician certified in medical acupuncture	Acupuncture improved cancer-related pain and other symptoms	Not reported	4
Lim <i>et al</i> 2011 ⁴²	Canada	Pilot RCT	31–81	20	Advanced cancer pain	Electroacupuncture	(ESAS)	20 min/ treatment weekly for 4 weeks	Principal investigator, a radiation oncologist and a certified medical acupuncturist	Acupuncture had an immediate effect on all symptoms including pain and it was well tolerated	Leg stiffness; falling asleep sensation	2
Dean-Clower <i>et al</i> 2010 ⁴⁵	USA	Pilot prospective trial	>21	32	Advanced cancer pain	Acupuncture	Brief Pain Inventory	12 acupuncture sessions over 8 weeks	Licensed acupuncturists	Patients' pain severity and pain interference scores were significantly improved	None	3
Johnstone <i>et al</i> 2002 ⁴¹	USA	Retrospective chart review	19–80	41	Cancer or cancer-related pain	Traditional Chinese acupuncture, auricular acupuncture, percutaneous electrical nerve stimulation, and Korean hand and Japanese scalp acupuncture+standard allopathic care	Visual Analogue Scale	Unknown number of acupuncture sessions over 4 months	Licensed physician; acupuncturist	55% pain patients showed at least 30% improvement, and 33% showed at least 80% improvement	None	4
Dillon and Lucas 1999 ⁴⁴	UK	Non-randomised cohort study	22–78	28	Malignant bone pain	Auricular stud acupuncture/Seirin semipermanent press needle	Subjective assessment of pain relief (0–4 scale)	2-week intervals for 4 weeks	–	Auricular acupuncture significantly reduced pain levels	None	3

* Levels of Evidence based on the Quality Rating Scheme for Studies and Other Evidence modified from the Oxford Centre for Evidence-Based Medicine for rating of individual studies; available online at <https://www.cebm.net/2016/05/occbm-levels-of-evidence/>.

RCT, randomised controlled trial.

hand and Japanese scalp acupuncture,⁴¹ and auricular stud.⁴⁴ Stud referred to a standard 'Seirin' semipermanent press needle (2.9 mm×1.55 mm) inserted into the sensitive area of ears under aseptic conditions.⁴⁴ The frequency and duration of the treatment session varied among the included trials. Except for the non-randomised cohort study in which medical providers were not clearly specified,⁴⁴ the other acupuncture treatments of the retrospective chart review were provided by physicians certified in medical acupuncture⁴³ or licensed acupuncturists. All the included studies focused primarily on malignant bone pain,⁴⁴ cancer and cancer-related pain.

The assessment tools used to evaluate analgesic effectiveness were varied. One used subjective assessment of pain relief (ie, none, slight, moderate, good and complete) on a scale (0–4),⁴⁴ and the other used the Edmonton Symptom Assessment Scale (0–10).⁴³ All of the trials reported effective outcomes related to acupuncture therapies.

Study results showed that acupuncture provided a favourable effect on pain management for patients with cancer in palliative care. In the five studies, only one study did not mention AEs associated with acupuncture.⁴⁴ Three studies reported no AEs, and one included leg stiffness and a 'falling asleep' sensation, which cleared without treatment after 1 hour. Therefore, in these studies, acupuncture therapy appeared to be a safe treatment for pain management. The methodological quality of the included trials was generally not strong. Based on the Oxford grading of evidence, one study was graded as level 2 (20%),⁴² two studies were classified as level 3 (40%)^{44 45} and the remaining two were level 4 (40%).^{41 43} Study characteristics are detailed in [table 1](#).

DISCUSSION

Our systematic review aimed to evaluate the current available evidence to determine whether acupuncture therapy is effective and safe for palliative care among patients with cancer pain. Although a considerable number of reviews concerning acupuncture for cancer pain have been published, the efficacy and safety of acupuncture for cancer pain in palliative care settings has been relatively unknown. The synthesis of the reviewed studies generated five publications including two retrospective chart reviews, one pilot RCT, a pilot prospective trial and one non-randomised cohort study. The findings from these previous studies indicated that acupuncture might be effective and safe in providing pain relief in palliative care in patients with cancer. Robust supporting evidence in the setting of palliative care is lacking, making it difficult to draw firm conclusions. More academically rigorous studies are required to examine this subject in the future.

The optimal observed interventions for studies of acupuncture are controversial. Taking into account the potential active interference of physiological or

psychological effects, strict inclusion criteria were applied to reduce the heterogeneity in this review, and non-invasive acupuncture techniques in treatment groups, such as acupressure and laser acupuncture, were excluded in the screening process. Acupuncture techniques contained in this review varied, the treatment tool ranged from filiform needle, electroacupuncture, PENS, scalp acupuncture to auricular acupuncture; reported duration and frequency of sessions also varied making it difficult to define an optimal therapy session duration or frequency.

Strengths and limitations

This study is the first review acupuncture as a component of palliative care in pain management for patients with cancer. The evidence suggests that acupuncture is effective and safe in managing cancer pain in palliative settings, but the results should be interpreted cautiously due to the small number and low quality of the included studies. More high-quality studies with large sample sizes are needed in the future. However, we recognise that there are numerous hurdles to such trials, such as difficulties with recruitment, attrition, randomisation, consistency of treatment approach, duration and frequency of treatments, and conducting trials in palliative care.^{46 47}

Beyond the research challenges, there remains a challenge from a practice perspective regarding palliative care in general. Though the WHO extended palliative care as a component of integrated treatment throughout the whole process of life-threatening illness, the clinical practice continues to face major challenges. Palliative care is very important to patients with life-threatening illnesses, but it is often misconstrued as end-of-life care, consequently resulting in referrals happening too late, and with only 10% of patients needing palliative care actually receiving it.⁴⁸ Our review addresses the unmet need for acupuncture in palliative care treatment and can serve to guide further clinical studies and practice from a methodological standpoint.

Conclusions

Overall, the limited data suggest acupuncture may be safe and provide benefit in the palliative management of cancer pain. Given the methodological limitation of included trials, more high-quality clinical studies with large sample sizes are needed in the future.

Acknowledgements The authors would like to thank the experienced librarian, Leslie C Hassett, Mayo Clinic Rochester Campus, for her significant contributions to the literature research and valuable constructive advice. We would also thank The HEAD Foundation for their support of this work.

Contributors BAB and DM conceptualised the study. JY, DW and DRP designed the study. JY, XZ and MS searched, screened, extracted and analysed the data. JY and LAJ wrote the original draft. TYC, DM, BAB, AD, MS and DRP reviewed and edited the paper. All authors approved the final paper.

Funding This study was funded by The HEAD Foundation, Singapore.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Juan Yang <http://orcid.org/0000-0002-0850-679X>

REFERENCES

- Bray F, Ferlay J, Soerjomataram I, *et al.* Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394–424.
- Torre LA, Bray F, Siegel RL, *et al.* Global cancer statistics, 2012. *CA Cancer J Clin* 2015;65:87–108.
- van den Beuken-van Everdingen MHJ, Hochstenbach LMJ, Joosten EAJ, *et al.* Update on prevalence of pain in patients with cancer: systematic review and meta-analysis. *J Pain Symptom Manage* 2016;51:1070–90.
- Rodriguez C, Ji M, Wang H-L, *et al.* Cancer pain and quality of life. *J Hosp Palliat Nurs* 2019;21:116–23.
- Fallon M, Giusti R, Aielli F, *et al.* Management of cancer pain in adult patients: ESMO clinical practice guidelines. *Ann Oncol* 2018;29:iv166–91.
- Chiu HY, Hsieh YJ, Tsai PS. Systematic review and meta-analysis of acupuncture to reduce cancer-related pain. *Eur J Cancer Care* 2017;26:e12457.
- Miller KD, Siegel RL, Lin CC, *et al.* Cancer treatment and survivorship statistics, 2016. *CA Cancer J Clin* 2016;66:271–89.
- Sanft T, Denlinger CS, Armenian S. NCCN guidelines insights: survivorship, version 2.2019: Featured updates to the NCCN guidelines. *J Natl Compr Canc Netw* 2019;17:784–94.
- Silverman J, Gulati A, Pain C. *Academic pain medicine*. Springer, 2019: 177–210.
- Wiffen PJ, Wee B, Derry S. Opioids for cancer pain—an overview of Cochrane reviews. *Cochrane Database Syst Rev* 2017;7.
- Are M, McIntyre A, Reddy S. Global disparities in cancer pain management and palliative care. *J Surg Oncol* 2017;115:637–41.
- Bruera E, Paice JA. Cancer pain management: safe and effective use of opioids. *Am Soc Clin Oncol Educ Book* 2015;35:e593–9.
- Hartl S. *Palliative care. ERS Handbook of respiratory medicine*, 2019: 249.
- Singer AE, Goebel JR, Kim YS, *et al.* Populations and interventions for palliative and end-of-life care: a systematic review. *J Palliat Med* 2016;19:995–1008.
- Kavalieratos D, Corbelli J, Zhang D, *et al.* Association between palliative care and patient and caregiver outcomes: a systematic review and meta-analysis. *JAMA* 2016;316:2104–14.
- Soares LGL. The pain Watch: it's about time to relieve cancer pain. *Ann Palliat Med* 2019;8:523–4.
- Helms JM. *Acupuncture energetics: a clinical approach for physicians*. Thieme, 1995.
- Wu X, Chung VCH, Hui EP, *et al.* Effectiveness of acupuncture and related therapies for palliative care of cancer: overview of systematic reviews. *Sci Rep* 2015;5:16776.
- Vickers AJ, Cronin AM, Maschino AC, *et al.* Acupuncture for chronic pain: individual patient data meta-analysis. *Arch Intern Med* 2012;172:1444–53.
- Cho Y-H, Kim C-K, Heo K-H, *et al.* Acupuncture for acute postoperative pain after back surgery: a systematic review and meta-analysis of randomized controlled trials. *Pain Pract* 2015;15:279–91.
- Murakami M, Fox L, Dijkers MP. Ear acupuncture for immediate pain Relief-A systematic review and meta-analysis of randomized controlled trials. *Pain Med* 2017;18:551–64.
- Kaptchuk TJ. Acupuncture: theory, efficacy, and practice. *Ann Intern Med* 2002;136:374–83.
- Lu W, Dean-Clower E, Doherty-Gilman A, *et al.* The value of acupuncture in cancer care. *Hematol Oncol Clin North Am* 2008;22:viii:631–48.
- NIN Consensus Development Panel on Acupuncture. Acupuncture. *JAMA* 1998;280:1518–24.
- Hershman DL, Unger JM, Greenlee H, *et al.* Effect of acupuncture vs sham acupuncture or Waitlist control on joint pain related to aromatase inhibitors among women with early-stage breast cancer: a randomized clinical trial. *JAMA* 2018;320:167–76.
- Wu X, Chung VCH, Hui EP, *et al.* Effectiveness of acupuncture and related therapies for palliative care of cancer: overview of systematic reviews. *Deutsche Zeitschrift für Akupunktur* 2017;60:28–9.
- Crew KD, Capodice JL, Greenlee H, *et al.* Randomized, blinded, sham-controlled trial of acupuncture for the management of aromatase inhibitor-associated joint symptoms in women with early-stage breast cancer. *J Clin Oncol* 2010;28:1154–60.
- Birch S, Lee MS, Alraek T, *et al.* Evidence, safety and recommendations for when to use acupuncture for treating cancer related symptoms: a narrative review. *Integr Med Res* 2019;8:160–6.
- Paice JA, Portenoy R, Lacchetti C, *et al.* Management of chronic pain in survivors of adult cancers: American Society of clinical oncology clinical practice guideline. *J Clin Oncol* 2016;34:3325–45.
- Chiu HY, Hsieh YJ, Tsai PS. Systematic review and meta-analysis of acupuncture to reduce cancer-related pain. *Eur J Cancer Care* 2017;26:e12457.
- He Y, Guo X, May BH, *et al.* Clinical evidence for association of acupuncture and Acupressure with improved cancer pain: a systematic review and meta-analysis. *JAMA Oncol* 2020;6:271–278.
- Hu C, Zhang H, Wu W, *et al.* Acupuncture for pain management in cancer: a systematic review and meta-analysis. *Evid Based Complement Alternat Med* 2016;2016:1–13.
- Lee H, Schmidt K, Ernst E. Acupuncture for the relief of cancer-related pain - a systematic review. *Eur J Pain* 2005;9:437–44.
- Paley CA, Johnson MI, Tashani OA, *et al.* Acupuncture for cancer pain in adults. *Cochrane Database Syst Rev* 2015;2015:CD007753.
- Paley CA, Tashani OA, Bagnall A-M, *et al.* A Cochrane systematic review of acupuncture for cancer pain in adults. *BMJ Support Palliat Care* 2011;1:51–5.
- Lau CHY, Wu X, Chung VCH, *et al.* Acupuncture and related therapies for symptom management in palliative cancer care: systematic review and meta-analysis. *Medicine* 2016;95:e2901.
- Lian W-L, Pan M-qi, Zhou D-han, *et al.* Effectiveness of acupuncture for palliative care in cancer patients: a systematic review. *Chin J Integr Med* 2014;20:136–47.
- Towler P, Molassiotis A, Brearley SG. What is the evidence for the use of acupuncture as an intervention for symptom

- management in cancer supportive and palliative care: an integrative overview of reviews. *Support Care Cancer* 2013;21:2913–23.
- 39 Liberati A, Altman DG, Tetzlaff J, *et al.* The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med* 2009;6:e1000100.
- 40 Howick J, Chalmers I, Glasziou P. *OCEBM Levels of Evidence Working Group 'The Oxford 2011 Levels of Evidence': Oxford Centre for Evidence-Based Medicine.* Oxford, 2011.
- 41 Johnstone PAS, Polston GR, Niemtow RC, *et al.* Integration of acupuncture into the oncology clinic. *Palliat Med* 2002;16:235–9.
- 42 Lim JTW, Wong ET, Aung SKH. Is there a role for acupuncture in the symptom management of patients receiving palliative care for cancer? A pilot study of 20 patients comparing acupuncture with nurse-led supportive care. *Acupunct Med* 2011;29:173–9.
- 43 Miller KR, Patel JN, Symanowski JT, *et al.* Acupuncture for cancer pain and symptom management in a palliative medicine clinic. *Am J Hosp Palliat Care* 2019;36:326–32.
- 44 Dillon M, Lucas C. Auricular stud acupuncture in palliative care patients. *Palliat Med* 1999;13:253–4.
- 45 Dean-Clover E, Doherty-Gilman AM, Keshaviah A, *et al.* Acupuncture as palliative therapy for physical symptoms and quality of life for advanced cancer patients. *Integr Cancer Ther* 2010;9:158–67.
- 46 Hudson P, Aranda S, McMurray N. Randomized controlled trials in palliative care: overcoming the obstacles. *Int J Palliat Nurs* 2001;7:427–34.
- 47 Holm M, Carlander I, Fürst C-J, *et al.* Delivering and participating in a psycho-educational intervention for family caregivers during palliative home care: a qualitative study from the perspectives of health professionals and family caregivers. *BMC Palliat Care* 2015;14:16.
- 48 Hjerstad MJ, Aass N, Aielli F, *et al.* Characteristics of the case mix, organisation and delivery in cancer palliative care: a challenge for good-quality research. *BMJ Support Palliat Care* 2018;8:456–67.