



OPEN ACCESS

# Pain management in advanced cancer: physical activity as an outcome – accelerometer feasibility study

Sarah Lord ,<sup>1</sup> Phillip Good ,<sup>2,3</sup> Gregore Iven Mielke<sup>4</sup>

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/spcare-2022-003940>).

<sup>1</sup>Palliative Care Department, The Prince Charles Hospital, Chermside, Queensland, Australia

<sup>2</sup>Supportive and Palliative Care, Mater Health Services, Brisbane, Queensland, Australia

<sup>3</sup>Department of Palliative Care, St Vincent's Private Hospital, Brisbane, Queensland, Australia

<sup>4</sup>School Of Public Health, The University of Queensland, Saint Lucia, Queensland, Australia

## Correspondence to

Dr Sarah Lord, Palliative Care Department, The Prince Charles Hospital, Chermside, Australia; [Sarah.Lord@health.qld.gov.au](mailto:Sarah.Lord@health.qld.gov.au)

Received 29 August 2022

Accepted 30 August 2022

Published Online First

13 September 2022



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Lord S, Good P, Mielke GI. *BMJ Supportive & Palliative Care* 2023;**13**:e843–e846.

## ABSTRACT

**Objective** Cancer pain is a common distressing symptom. Numerical Pain Scales (NPS) assess pain but lack information about function and quality of life. This feasibility study assesses the use of triaxial accelerometers to measure function as an outcome measure in pain studies in advanced cancer.

**Methods** Advanced cancer participants were recruited from two palliative care services, with an average pain score of  $\geq 3$  on NPS. ActiGraph wGT3X-BT Accelerometers were worn for 1 week on the wrist. Patients recorded daily pain scores, Edmonton Symptom Assessment Scale (ESAS) scores, and their daily opioid use.

**Results** 24 participants were recruited. A total of 142 days of accelerometer data was collected (5.9 days/participant). The average daily step count was 5723.7. The average acceleration was 14.4 milligravity units/day. An average of 93 min/day total activity across all intensities was recorded. No correlation was seen between acceleration or average daily minutes in activity and total daily oral morphine equivalent, ESAS, 'average pain' score or 'worst pain' scores using spearman's correlation coefficients. Overall, participants were satisfied with the study.

**Conclusions** Accelerometers are a feasible method to measure activity as an outcome measure in advanced cancer. Further study is required to assess the impact of pain management strategies on function.

## INTRODUCTION

Triaxial accelerometers offer a way to measure activity and pain management in patients with advanced cancer. This new technology uses motion sensors to detect the body's acceleration in up to three planes (triaxial) and record physical function,<sup>1</sup> activity and sleep. It is lightweight and can be attached to the body or

### WHAT WAS ALREADY KNOWN?

- ⇒ Pain commonly impacts on function in advanced cancer.
- ⇒ Ability to function is one of the main goals of pain management.

### WHAT ARE THE NEW FINDINGS?

- ⇒ Patients with advanced cancer function significantly less than healthy adults.
- ⇒ Triaxial accelerometers effectively measure function in this group.

### WHAT IS THEIR SIGNIFICANCE?

- Clinical
  - Function can be measured easily.
- Research
  - Further studies into how pain impact's function.

worn on the wrist and is an objective and accurate method of measuring activity for patients with advanced cancer.<sup>2</sup>

Studies suggest that two-thirds of patients with advanced cancer experience pain,<sup>3</sup> that 50% of these patients describe the pain as moderate or severe<sup>3</sup> and their main goal of pain management, is the ability to live normally and complete everyday tasks.<sup>4</sup> Many studies use Numerical Pain Scales to assess pain and its response to treatment<sup>5</sup> but these scales are unidimensional, only measuring severity, and not other aspects of pain. Self-reporting of pain and physical activity are subjective, rely on recall, are known to be inaccurate and prone to bias<sup>6</sup> and also significantly underestimates sedentary time.<sup>7</sup> Few studies exist that objectively measure physical activity in this patient group to provide accurate data about physical function in relation to pain management.

This study investigates the use of accelerometers to measure physical activity

in patients experiencing cancer-related pain and how activity correlates with pain and other symptoms.

## METHODS

This study is a prospective, multicentre study of patients with advanced cancer and an average pain score of  $\geq 3$  (Numerical Rating Scale or NRS). Patients were recruited by the palliative care teams at two centres in Brisbane, via outpatient clinic, community services or the inpatient palliative care ward. There were no exclusion criteria.

The primary outcome was to assess the feasibility of using accelerometers to monitor pain in patients with advanced cancer. This was achieved if  $\geq 60\%$  of participants wore the accelerometer for  $\geq 80\%$  of the time during the 6-day trial period. Secondary outcomes included pain scores, total symptom scores and opioid dosing. The study was powered to define feasibility (20 participants).

Participants wore a commercial accelerometer (Acti-Graph wGT3X-BT) on their wrists for 6–7 days. Baseline demographic data and total daily oral morphine equivalent (OME) doses was collected at day 0. Participants recorded daily pain relief usage, pain scores (brief pain inventory) and symptom scores (Edmonton Symptom Assessment Scale (ESAS)).


## Statistical analysis

The accelerometer recorded raw acceleration in three axes and provided raw data in gravitational equivalent units (g) ( $1\text{ g}=9.81\text{ m/s}^2$ ). Raw data were processed in R using the most up to date GGIR package, a widely used open-source code.<sup>8</sup> The vector magnitude of the three axes was used to calculate activity-related acceleration using Euclidian Norm minus 1 g [ $\text{ENMO}=\sqrt{(x^2+y^2+z^2)}-1$ ]. Data were initially aggregated in 5 s time series and included if wear time was at least 600 min/day. Data were used to quantify overall physical activity expressed as acceleration in milligravity units (mg), as well as time spent in activities at different intensities. Active minutes were defined as activities with average acceleration  $>30\text{ mg}$ .

Daily acceleration and active minutes were correlated with daily scores for OME (mg), total ESAS, worst and average pain scores (0–10 on NRS) to produce scatterplots. Correlation was analysed using Spearman's correlation to produce spearman correlation coefficient with 95% CIs.

## RESULTS

The recruitment period was extended (December 2018 to May 2021) due to the Coronavirus Pandemic. A total of 24 patients (online supplemental file 1) were recruited, with 142 days of accelerometer data recorded for those participants, equating to an average of 5.9 days per participant. An average 20.8 hours (SD: 5.2 hours) was recorded for each day of data; collecting a total of 2953.6 hours of accelerometer data.



**Figure 1** Daily minutes of physical activity compared with physical activity intensity and daily scores for pain, opioid use and symptoms. (A) average minutes in different physical activity intensities. ( $n=142$  days) ( $<50\text{ mg}$  units is considered inactive); (B) correlation between daily average pain score and daily minutes in activity (Spearman's  $r=0.10$ ;  $p=0.260$ ); (C) correlation of oral morphine equivalent (OME) (total mg/day) with minutes in activity per day (Spearman's  $r=-0.05$ ;  $p=0.630$ ); (D) correlation between total daily Edmonton Symptom Assessment Scale (ESAS) scores and daily minutes in activity (Spearman's  $r=-0.14$ ;  $p=0.109$ ).

The average daily step count of participants was 5723.69, with a range from 1117 to 21205 steps per day when incomplete days were removed. The average acceleration of participants was 14.4 milli gravity units (SD: 7.3) across all days. Most time during the day was spent inactive (including sleep time), with an average total activity time of 93 mi/day (figure 1). Most of the time spent active was in activities with an average acceleration between 50 and 99 mg (light intensity activity). Daily median duration of light intensity activity was 73 min. Using a cut-off from the healthy adult population, median moderate intensity activity was 20 min, with nearly three-quarters of this time in activity with an average acceleration of 100–149 mg. There was no activity of vigorous intensity recorded.

No correlation was seen between total daily OME dosing and either acceleration or minutes in activity (figure 1) (Spearman's  $r=-0.12$ ;  $p=0.179$  and Spearman's  $r=-0.05$ ;  $p=0.630$ , respectively). A trend towards lower activity (both average acceleration and minutes in activity (figure 1)) at higher OME doses above 200 mg day was seen, but this was not statistically significant. No correlation was demonstrated between total symptom score (measured using ESAS) and either minutes in activity (figure 1) or acceleration.

The median average pain score at baseline was 4, with a range of 2–9 (0–10). Baseline median 'least pain' was 2 (range 0–9) and 'worst pain' was 6 (range 4–9). No correlation was seen between average pain score and acceleration (Spearman's  $r=0.10$ ;  $p=0.261$ ) or the number of daily minutes in activity (Spearman's  $r=0.10$ ;  $p=0.260$ ). The full range of activity was seen

at all pain scores, with the most minutes in activity seen with an average pain score of 6. There was also no correlation seen between daily 'worst pain' scores and average acceleration (Spearman's  $r = -0.09$ ;  $p = 0.298$ ) or daily minutes in activity (Spearman's  $r = -0.11$ ;  $p = 0.237$ ).

Overall participants were satisfied with the study, with 74% stating they were satisfied or mostly satisfied. Only one participant was somewhat dissatisfied with their involvement (patient global impression of change scale).

## DISCUSSION

The primary outcome of assessing the feasibility of using accelerometers to measure activity in advanced cancer was achieved. The amount of accelerometer data collected per patient suggests it is acceptable for participants with advanced cancer to wear these devices. While other clinical studies<sup>9</sup> highlighted compliance as a major issue, it was not an issue in our study and patients reported a high level of satisfaction in wearing the device.

As expected, the daily average of active time is lower than the average healthy adult population (209 min/day),<sup>10</sup> and the average step count (5723.69) is consistent with prior studies.<sup>11</sup> Most of the activity was low intensity activity, and about half that of the healthy population (141 min/day).<sup>10</sup> Interestingly some participants had higher than average activity despite their disease state, living very active lives despite their pain and symptoms. Importantly, while the average activity is low, it was possible to capture this accurately using the accelerometer.

Opioid use (below 200 mg OME a day) appears to have little impact on activity levels. There is a trend for higher doses of opioids to be correlated with less activity; possibly due to a higher rate of sedative side effects, worse total pain or higher doses in response to pain control. The sample in this study is small and other disease or treatment factors have not been accounted for, so this trend is uncertain and worth investigating further.

Pain scores did not appear to impact level of function. Pain (highly subjective)<sup>12</sup> and symptom scores vary in the individual and between individuals with similar pathology.<sup>13</sup> The ability to measure a change in function might enable us to better understand the dynamic of pain and function in an individual patient.

The total symptom burden also does not appear to impact physical activity levels. Some patients function well despite high overall symptom burden. Our results align with another study that assessed quality of life and self-reported activity in palliative cancer patients, which also identified the association of higher activity with lower fatigue subscores.<sup>14</sup> The relationship between patient activity and specific symptoms or disease burden is complex, and further studies are

needed to assess the impact of specific symptoms on function.

Patients with advanced cancer desire increased function with pain, and improved quality of life (known association with increased activity).<sup>14</sup> Examining the change in activity and pain levels before and after intervention with medications, and assessing toxicity of these medications could help guide which medications provide the most functional benefit to patients in pain, and appropriate dosing. Future studies using accelerometers could provide valuable information as to how best to achieve this for a patient group where function is particularly important.

## CONCLUSIONS

Measuring pain using standard questionnaires and patient recall is subjective, inaccurate and prone to bias.<sup>6</sup> Accelerometers are a promising tool to measure activity in advanced cancer, to further understand how we can help maximise function and improve the quality of life in our patients. We demonstrated it is feasible to use accelerometers to measure activity in advanced cancer, however, correlation between pain, opiate use, total symptom distress scores and function cannot be drawn from this study.

**Twitter** Gregore Iven Mielke @GregoreMielke

**Contributors** PG designed the study. PG and SL were responsible for recruitment and data collection. SL completed initial data analysis, with accelerometer and statistical analysis by GIM. SL wrote the initial draft of the paper with contributions by both PG and GIM.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants and was approved by the Mater Human Research Ethic Committee (Brisbane Australia) HREC/17/MHS/131. Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; internally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-

commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Sarah Lord <http://orcid.org/0000-0002-7764-5064>  
Phillip Good <http://orcid.org/0000-0002-8198-0375>

#### REFERENCES

- 1 Welk GJ, Schaben JA, Morrow JR. Reliability of accelerometry-based activity monitors: a generalizability study. *Med Sci Sports Exerc* 2004;36:1637–45.
- 2 Maddocks M, Byrne A, Johnson CD, *et al*. Physical activity level as an outcome measure for use in cancer cachexia trials: a feasibility study. *Support Care Cancer* 2010;18:1539–44.
- 3 van den Beuken-van Everdingen MHJ, Hochstenbach LMJ, Joosten EAJ, *et al*. Update on prevalence of pain in patients with cancer: systematic review and meta-analysis. *J Pain Symptom Manage* 2016;51:1070–90.
- 4 Gibbins J, Bhatia R, Forbes K, *et al*. What do patients with advanced incurable cancer want from the management of their pain? A qualitative study. *Palliat Med* 2014;28:71–8.
- 5 Farrar JT, Young JP, LaMoreaux L, *et al*. Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale. *Pain* 2001;94:149–58.
- 6 Prince SA, Adamo KB, Hamel M, *et al*. A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act* 2008;5:56–24.
- 7 Prince SA, Cardilli L, Reed JL, *et al*. A comparison of self-reported and device measured sedentary behaviour in adults: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act* 2020;17:31.
- 8 Migueles JH, Rowlands AV, Huber F, *et al*. GGIR: a research community-driven open source R package for generating physical activity and sleep outcomes from multi-day RAW accelerometer data. *J Meas Phys Behav* 2019;2:188–96.
- 9 Maddocks M, Granger CL. Measurement of physical activity in clinical practice and research: advances in cancer and chronic respiratory disease. *Curr Opin Support Palliat Care* 2018;12:219–26.
- 10 Mielke GI, Burton NW, Brown WJ. Raw triaxial wrist accelerometer-measured physical activity in mid-age Australian adults, March 2022, PREPRINT (version 1) available at research square.
- 11 Maddocks M, Wilcock A. Exploring physical activity level in patients with thoracic cancer: implications for use as an outcome measure. *Support Care Cancer* 2012;20:1113–6.
- 12 Selby D, Cascella A, Gardiner K, *et al*. A single set of numerical cutpoints to define moderate and severe symptoms for the Edmonton symptom assessment system. *J Pain Symptom Manage* 2010;39:241–9.
- 13 Lowe SS, Danielson B, Beaumont C, *et al*. Associations between objectively measured physical activity and quality of life in cancer patients with brain metastases. *J Pain Symptom Manage* 2014;48:322–32.
- 14 Lowe SS, Watanabe SM, Baracos VE, *et al*. Associations between physical activity and quality of life in cancer patients receiving palliative care: a pilot survey. *J Pain Symptom Manage* 2009;38:785–96.