

(5%) had severe refractory pain requiring palliative sedation with levomepromazine and/or phenobarbitone.

Discussion and Conclusion Our findings showed that complex pain is highly prevalent in young adult cancer patients at the end of life requiring opioids, adjuncts and interventional pain strategies for management. This may be explained by age specific tumour types and predilection for bone and pelvic pain syndromes. Our data offer important information to the limited existing information in this area. In comparison standard palliative care cohort, our data showed that a large proportion of young adult patients require high doses of opioids for analgesia defined as OMEDD of more 300 mg2. Despite this pain remained poorly controlled in some patients requiring benzodiazepine or palliative sedation at the end of life, suggestive of total pain phenomenon, similarly reflected in several case reports¹. When required, these patients require high doses of sedatives. Future research is needed to identify tailored treatment regimens for young adults with cancer-related pain, focusing on psychological symptom profile encompassing total pain.

REFERENCES

1. Abdelaal, *et al*. Palliative care for adolescents and young adults with advanced illness: a scoping review. *Palliative Medicine* 2023;**37**(1):88–107.
2. Bercovitch, *et al*. High dose morphine use in the hospice setting. *Cancer* 2000;**86**(5):735–903.

P-29

THE EVOLUTION OF 'PALLIATIVE CARE: ITS MORE THAN THEY THINK' CAMPAIGN

^{1,2,3}Sarah Lord, ¹Elise Power, ⁴Stuart Ekberg. ¹The Prince Charles Hospital, Chermiside, Australia; ²Ipswich Hospital, Ipswich, Australia; ³University of Queensland, Brisbane, Australia; ⁴Queensland University of Technology, Brisbane, Australia

10.1136/spcare-2024-ANZSPM.77

Research conducted in The Prince Charles Hospital, Brisbane using conversational analysis of direct observation during initial Palliative Care outpatient consultations highlighted that patients have a very limited understanding of what palliative care is. Often palliative care is thought to involve care that occurs only immediately before death. This occurs despite referrals being made by medical professionals. Therefore, we surmised that either these clinicians also do not understand the scope of palliative care, or they are not effectively communicating this to the patients they are referring.

We set out to try to better understand clinicians' attitudes to palliative care and how we could improve their understanding of the full scope of palliative care enabling them to be able to communicate this better with patients.

The project was focused on the thoracic team at The Prince Charles Hospital, being one of the major referring teams. To assess understanding an anonymised survey was undertaken using the Knowledge and attitudes towards hospital and palliative care (KAHP) scale and several free text questions aimed at barriers and facilitators to palliative care referrals. Both medical practitioners and senior nursing staff were invited to complete the survey electronically.

19 responses from roughly 100 invited clinicians were received. 13 of these were from medical practitioners. Overall, clinicians felt that patients would benefit if palliative care was initiated earlier in the course of the illness, that palliative care improved symptom control and met the needs of the family better than conventional care. However, clinicians felt that

discussing palliative care could cause patients and families to lose hope and that telling patients that they are dying is difficult. Despite this more than half of the respondents felt knowledgeable enough to discuss palliative care and well trained to take care of the symptoms in life limiting illnesses. Highlighted barriers to palliative care referral included a lack of time, not discussing palliative care with the patient and patients not being ready for palliative care. Potential facilitators to easier referral included clearer referral processes, increased knowledge about local resources, access to patient brochures and an increased profile of palliative care.

This led to the development of educational materials for clinicians with Palliative Care Australia (PCA) and the 'More than they think campaign'. Resources developed included a brochure, posters, slide show, video and a local services fact-sheet. These materials have been rolled out in The Prince Charles Hospital as part of an education program to the Thoracic department. They have also been made available via the PCA website for use by other services. We are currently in the process of evaluating the effectiveness of this education campaign.

P-30

IL-16 BLOOD LEVELS AND COMBINED POLYMORPHISM OF CCL11 AND IL-16 ARE THE BIOMARKERS TO SELECT OXYCODONE FOR CANCER PAIN MANAGEMENT

^{1,2}Hiromichi Matsuoka, ¹Yoshihiko Fujita, ¹Junji Tsurutani, ¹Takeshi Yoshida, ¹Atsuko Koyama, ¹Kazuto Nishio, ¹Kazuhiko Nakagawa. ¹Kindai University Faculty of Medicine, Osakasayama, Japan; ²National Cancer Center Hospital, Chuoku, Tsukiji, Japan

10.1136/spcare-2024-ANZSPM.78

Background For precision medicine for cancer pain, we identified a SNP in CCL11 (rs17809012) as one of the biomarkers significantly associated with the analgesic effect of morphine by screening 74 pain-related single nucleotide polymorphisms (SNPs).¹ In this study, to explore biomarkers for predicting opioid efficacy, we aimed to evaluate whether plasma concentrations of chemokines/cytokines and their SNPs in combination can accurately predict the most appropriate opioid for pain relief in cancer patients.

Methods In this study, plasma concentrations of several chemokines/cytokines were determined in pretreatment plasma samples obtained from a total of 138 patients enrolled in our previous clinical trial² who were randomly assigned to the morphine (N=70) and oxycodone (N=68) groups. The relationship between pre-treatment blood concentrations of various chemokines/cytokines and NRS (opioid analgesic effect) in the oxycodone group was investigated using simple regression analysis. Regarding IL-16, which showed promising results, we performed simple regression analysis using opioid type as independent variable and Δ NRS as dependent variable and multiple regression analysis using opioid type and IL-16 concentration (high or low) and opioid type IL-16 concentration (interaction term) as independent variables and NRS as dependent variable among all patients. Finally, we evaluated the relationship between the combination of both CCL11 and IL-16 SNPs and opioid efficacy using multiple regression analysis.

Results In the oxycodone group, there was a significant difference in NRS between groups ($p=0.013$) of patients with high ($n=34$) and low ($n=34$) blood levels of IL-16, and oxycodone was more effective in patients with lower IL-16 levels ($p=0.038$), whereas morphine was more effective in patients