


Asymptomatic COVID-19 infection: diagnosis, transmission, population characteristics

Yaxian You,^{1,2} Xinyuan Yang,^{1,2} Dongni Hung,^{1,2} Qianxi Yang,^{1,2} Ting Wu ,^{1,3} Meichun Deng^{1,2,4}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjspcare-2020-002813>).

¹Department of Biochemistry and Molecular Biology & Hunan Province Key Laboratory of Basic and Applied Hematology, School of Life Sciences, Central South University, Changsha, Hunan, China

²Xiangya School of Medicine, Central South University, Changsha, Hunan, China

³Department of Cardiovascular Medicine, The Third Xiangya Hospital, Central South University, Changsha, Hunan, China

⁴Hunan Key Laboratory of Animal Models for Human Diseases, Hunan Key Laboratory of Medical Genetics, School of Life Sciences, Central South University, Changsha, Hunan, China

Correspondence to

Dr Meichun Deng, Central South University, Changsha 410013, Hunan, China; dengmch@csu.edu.cn

YY and XY contributed equally.

Received 21 November 2020

Revised 26 February 2021

Accepted 27 March 2021

Published Online First

30 July 2021



© Author(s) (or their employer(s)) 2024. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: You Y, Yang X, Hung D, et al. *BMJ Supportive & Palliative Care* 2024;**14**:e220–e227.

ABSTRACT

A novel coronavirus first discovered in late December 2019 has spread to many countries around the world. An increasing number of asymptomatic patients have been reported and their ability to spread the virus has been proven. This brings major challenges to the control of the transmission. The discovery and control of asymptomatic patients with COVID-19 are the key issues in future epidemic prevention and recovery. In this narrative review, we summarise the existing knowledge about asymptomatic patients and put forward detection methods that are suitable for finding such patients. Besides, we compared the characteristics and transmissibility of asymptomatic patients in different populations in order to find the best screening, diagnosis and control measures for different populations. Comprehensive preventive advice is also provided to prevent the spread of infection from asymptomatic patients.

INTRODUCTION

COVID-19 is an acute respiratory tract infection that arose in late 2019. The causative pathogen of COVID-19 has been confirmed to be a novel coronavirus. The virus was named SARS-CoV-2 by the WHO.¹ As of 14 November 2020, there have been 52 852 674 confirmed cases of COVID-19, including 1 295 328 deaths.² Many potential drugs have been used to treat COVID-19.³ As the pandemic has developed, an increasing number of asymptomatic patients has been found, causing worldwide anxiety. Two conditions must be met for asymptomatic COVID-19 cases: first, the absence of self-perceived or clinically recognisable symptoms; and second, a positive test for SARS-CoV-2. This definition actually includes patients during the latent period and covert infected patients who are asymptomatic from onset to recovery. For

Key messages

What was already known?

- ⇒ Asymptomatic patients have the potential for transmission.
- ⇒ Asymptomatic patients are ubiquitous in people of all ages.

What are the new findings?

- ⇒ The characteristics of asymptomatic patients vary in different groups.
- ⇒ Prevention of asymptomatic transmission requires context-specific methods.

What is their significance?

- ⇒ Clinical: Assist in the diagnosis of asymptomatic patients in different populations.
- ⇒ Research: Suggest the proportion and characteristics of asymptomatic patients.

some latent cases, although patients do not show any symptoms at the time of testing, they may eventually develop symptoms as the virus replicates in the body. Presymptomatic patients will be used in this review to represent latent cases, thus differentiating between the two when needed. When not specifically stated, this article focuses on cases that do not exhibit symptoms at the time of initial diagnosis of COVID-19. Current studies have shown that the combined proportion of patients who are asymptomatic at initial testing reaches 15.6%, of which approximately 48.9% are presymptomatic infections.⁴ The proportion of asymptomatic patients varies considerably among different groups (children, pregnant women, elderly, etc). The potential transmission of asymptomatic viruses imposes a significant burden on countries with secondary attack rates (SARs) ranging from 0% to 80%.⁵ This review aims to summarise the diagnosis, transmission and prevention of asymptomatic patients with COVID-19,




Figure 1 Examples of detection, transmission and screening of asymptomatic patients. Existing detection methods for asymptomatic patients include imaging, nucleic acid and serological diagnostics. Joint detection methods are highly efficient, accurate and more suitable for the discovery of asymptomatic patients. These methods are applied to screen out among discharged patients and persons released from isolation. It is also essential to conduct general monitoring and follow-up among the general population. These all will contribute to preventing further transmission of asymptomatic patients in homes, hospitals, and transmission through patient contaminants, thereby reducing the risk of infection in healthy people.

analyse the differences in the proportion, transmission capacity and characteristics of asymptomatic patients in different populations and finally propose methods for effective surveillance and control of the disease. We have summarized the main findings of this article using an (figure 1), which captures the content of the article for readers at a single glance.

Detection of asymptomatic patients

Clinical testing

There is no significant change in blood cell count and inflammation indicators in asymptomatic compared with patients with symptomatic COVID-19.⁶ Multiple organ dysfunction often occurs in symptomatic patients with little change in organ function in asymptomatic patients.⁷ An analysis of the imaging features of an asymptomatic infection showed that nearly half of the asymptomatic infections (47.62%) had lung abnormalities, and among the covert infections, 64.39% had lung abnormalities.⁸ Meng *et al* found that ground-glass opacity on CT scan was the main manifestation of asymptomatic patients (94.8%), with peripheral (75.9%) or unilateral (58.6%) location. The lower lung lobe (68.9% right vs 62.1% left) is more involved than the upper lobe (51.7% right vs 53.4% left).⁹ CT detects symptoms of COVID-19 earlier than nucleic acid testing does. About 75% of the patients

who were diagnosed negative by reverse transcription-polymerase chain reaction (RT-PCR) had abnormalities in their chest CT and had positive chest CT for COVID-19, but the negative result of

RT-PCR turned positive after 5.1 ± 1.5 days. In addition, studies have shown that chest CT (98%) has higher sensitivity than RT-PCR (71%) for detecting patients with COVID-19.¹⁰ A recent study found that fluorodeoxyglucose positron emission tomography/CT can identify possible COVID-19 disease in the absence of or before symptom onset.¹¹ Care should be taken when considering CT as a screening tool. CT scan has high sensitivity (86%–98%), but its diagnosis of COVID-19 often overlaps with that of other viral pneumonia, resulting in low specificity (about 25%).¹² Moreover, the sensitivity of CT testing is affected by a variety of factors, such as disease severity, the proportion of patients with comorbidities, the proportion of asymptomatic patients and the age of asymptomatic patients.¹³ Therefore, we do not recommend the use of CT alone in the initial diagnosis of COVID-19. In low-prevalence populations, CT chest scans may be more effective in detecting asymptomatic cases.¹⁴ It has also been found to be more reliable for early diagnosis in the elderly, but less reliable in young children.^{15 16} We suggest adopting the method of joint testing. It is

possible to combine both CT and RT-PCR to achieve adequate sensitivity and specificity. Research has shown that CT of the chest has been considered as a very important strategy for supplementary diagnosis.¹⁷

Laboratory testing

Nucleic acid detection

RT-PCR of respiratory specimens is currently the gold standard for diagnosis of COVID-19, with a sensitivity of approximately 96%. Both the virus load of the sampling site and operation of the test influence the accuracy of the results. In the early stages of viral infection, there may be some false-negative results due to insufficient amount of virus on the swab. Therefore, it may be better to perform multiple swab tests from different parts of the body when necessary.¹⁸ There are a variety of non-neuronal cell types in the olfactory epithelium of the nasal cavity, expressing two host receptors, angiotensin-converting enzyme 2 and transmembrane protease serines 2. These two receptors are considered to play an essential role in the binding and accumulation of SARS-CoV-2, which may make the nasal epithelium an enhanced binding site for the virus.¹⁹ For asymptomatic patients, a nasopharyngeal swab may be a better sampling method than the commonly used sputum and throat swabs. More recently, further studies have shown that nucleic acid testing using saliva samples appears to be more sensitive than nasopharyngeal swabs.²⁰ The sensitivity and accuracy of existing PCR methods are highly dependent on the performance of the kits and negatively correlated with the proportion of elderly patients.¹³ The prevalence of false-negative results suggests that nucleic acid testing is not entirely appropriate for the detection of asymptomatic patients. Any missed infections pose a threat to public safety.

Additionally, the cost of testing is high, it takes a long time, and health professionals are needed to perform the sampling. These factors all limit the large-scale application of RT-PCR for the screening of asymptomatic patients. In response to these problems, LabCorp Laboratories, Inc has developed an RT-PCR test kit for home specimen collection, enabling individuals to collect their own nasal swab samples at home, partially solving the problem of location limitation for nucleic acid testing.¹⁸ In addition, loop-received isothermal amplification (LAMP) is an economical and efficient alternative to standard PCR developed for the rapid detection of SARS-CoV-2. At present, a RT-LAMP kit developed by Sherlock Biosciences, Inc has been approved by Food and Drug Administration and European Union Association. According to Zhu *et al*, the accuracy and specificity of the nanoparticle-based RT-LAMP assay can reach 100%.²¹

Serological testing

Testing for specific antibodies in patients' blood is another option for COVID-19 diagnosis. Li *et al*'s

research showed that IgM antibodies and IgG antibodies appeared successively within 1 to 2 weeks after infection. The amount of IgG antibody reaches its highest value a few weeks after the onset of the disease. It will persist for months or even years, but the duration is still unclear.²² Many rapid tests for IgM or IgG have been produced so far. In these tests, using the principle of colloidal gold immunochromatography, results can be obtained within 15 min using 20 µL whole blood sample (or 10 µL of serum/plasma samples).²³ This method shortens the detection time. The tests are simple to perform and the results are easy to interpret, which breaks through the limitations of personnel and location of the existing detection methods. Antibody detection makes large-scale screening possible, which is important for the detection of asymptomatic patients. Some people who tested negative for nucleic acid tested positive for antibodies, suggesting that antibody testing can be used as a supplementary method for RT-PCR to improve diagnostic accuracy and reduce missed diagnosis.²⁴ In particular, for patients whose CT scan shows lung lesions but with negative pharyngeal swabs, specific IgM screening is useful for early detection and isolation. IgM-IgG combined antibody detection is considered to be a more reliable detection method, and it has greater specificity and sensitivity compared with single IgM or IgG tests. However, there are also some problems with antibody detection. There are significant individual differences in the IgG response to SARS-CoV-2 in different populations. Virus-specific IgG levels were significantly lower in asymptomatic patients compared with the symptomatic group, and a proportion of patients may not even have a SARS-CoV-2-IgG response.²⁵ Additionally, the tests do not confirm the presence of the virus, but only provide evidence of recent infection. Even after recovery, IgG can persist for a long time.²⁶ Furthermore, antibody test results are susceptible to interference from other viral infections, resulting in some false-positive results.

Joint testing

To improve the accuracy of detection and prevent potential virus transmission by asymptomatic patients, joint detection may be the most effective way to fill the gap. Symptoms of COVID-19 pneumonia frequently overlap with those of other viral infections, leading to low specificity of chest CT scans.²⁷ We do not recommend chest CT as the main screening tool. However, the sensitivity of chest CT is higher than that of RT-PCR; thus, it is reasonable to consider combining these two methods to achieve better sensitivity and accuracy.²⁸ Moreover, it is reported that combined use of nucleic acid detection and specific serum detection can help the discovery of asymptomatic patients.²⁹ The combination of serum antibody ELISA and quantitative PCR improves the detection rate of early diagnosis.³⁰

Asymptomatic patients in different populations

Regarding the proportion of asymptomatic patients in the general population, a systematic review study found that the proportion of diagnosed patients who were asymptomatic at initial testing ranged from approximately 20%–75%. After at least 7 days of follow-up of these patients, most reported symptoms within the first week. Only 4.1% remained asymptomatic throughout the disease.⁵ However, another study declared that the overall estimate of those who remained asymptomatic throughout the course of infection is 20%.³¹ Participant selection bias can affect the estimate. The exact proportion of asymptomatic patients needs to be further studied. Studies of viral shedding have found that asymptomatic viral hyperload periods can last up to 9 days. For pre-symptomatic patients, this number was 6 days before the onset of symptoms.³² The viral load of asymptomatic appears similar to that of symptomatic patients.³³

Regarding transmission in asymptomatic patients, the SARs in asymptomatic patients in the home setting was 0.7%, which was lower than the 18% in symptomatic cases.³⁴ In another study, SARs and observed reproduction in asymptomatic index cases were found to be about one-seventh of those in symptomatic index cases, while those in presymptomatic cases were two-thirds of those in symptomatic cases.³⁵ Adults showed a higher susceptibility than children.³⁶ Transmission was mainly associated with group activities, such as sharing meals, and not with the disease status of the cases.³⁵

Asymptomatic patients in children

A meta-analysis showed that about 26% of paediatric patients were asymptomatic.³⁷ Compared with adults, children may be more inclined to have mild or asymptomatic infections,³⁸ but they may have higher viral loads and longer latency and duration of viral shedding than adults.³⁹ Viral loads in the nasopharynx/oropharynx of asymptomatic children are lower compared with symptomatic children. However, viral load is often influenced by the duration of infection and cannot be used as evidence of the strength of transmission.⁴⁰ Some children do not show any symptoms but may develop radiological changes.⁴¹ Recently, a small number of children without respiratory symptoms have been found to develop a rare complication of the multisystem inflammatory syndrome, with overlapping features with Kawasaki disease. Also, some children are often accompanied by symptoms such as rhabdomyolysis and acute renal failure.⁴² These atypical symptoms should be addressed to optimise early identification and hospital management of the child.

Asymptomatic patients among pregnant women

The proportion of obstetric patients presenting to the hospital with initial testing asymptomatic was 95.1%, and the proportion remaining asymptomatic

throughout the course of the infection was approximately 58.8%.⁵ SARS-CoV-2 infection in pregnant women usually seems to be asymptomatic.⁴³ Vertical transmission of SARS-CoV-2 to the fetus has been demonstrated in cases, but the risk of vertical transmission is low, at approximately 2%.⁴⁴ However, infants whose mothers are infected with COVID-19 are more likely to experience adverse effects such as fetal distress, premature birth, respiratory distress and even death.⁴⁵ Ventilation support and earlier infection time of pregnant women seem to be the main determinants of adverse fetal outcomes.⁴⁶ Normal IgM and elevated IgG antibodies in newborns born to asymptomatic mothers have been reported.⁴⁷ Notably, Molina *et al* found that pregnant women may experience prolonged viral shedding of SARS-CoV RNA.⁴⁸ During a pandemic, healthcare workers should be cautious about delivering high-risk pregnant women and universal testing for COVID-19 should be considered in certain populations.⁴⁹

Asymptomatic patients among the elderly

The proportion of COVID-19 confirmed residents in nursing homes who were asymptomatic at initial testing was approximately 53.6%, and the proportion who remained asymptomatic throughout the follow-up period was approximately 27.9%.⁵ Compared with younger patients, elderly asymptomatic patients are more likely to develop symptoms during the course of the disease, which is consistent with the finding that disease severity is higher in older patients. Additionally, older patients often have a much longer latency period than younger patients.⁵⁰ Furthermore, Hitoshi *et al* found that the duration of illness in the asymptomatic group was positively correlated with patient age. Comorbidities in elderly patients, particularly hypertension, may be predictive of symptom development in initially asymptomatic carriers.⁵¹ It was also found in a nursing home report that elderly asymptomatic patients with COVID-19 often exhibit certain symptoms of the geriatric syndrome, such as falls, loss of appetite and altered consciousness.⁵² We may need to be vigilant for older people with such symptoms.

Transmission and prevention

According to reports, there is no significant difference in the viral load of respiratory samples from asymptomatic and symptomatic patients with COVID-19 as defined by RT-PCR.⁵³ The duration of virus shedding in the asymptomatic group was 19 days, obviously longer than the symptomatic group.⁵⁴ Asymptomatic patients are thought to make an essential contribution to virus transmission. There is growing evidence that SARS-CoV-2 can be transmitted by droplets, aerosol and conjunctiva. Faecal–oral transmission is also a possibility, whereas vertical mother-to-child transmission has not been confirmed.⁵⁵ The rate of secondary episodes of asymptomatic infection seems to be lower

than that of symptomatic infection, indicating that asymptomatic patients may be less infectious than symptomatic ones.³¹ Chen *et al* recently claimed that asymptomatic cases are more likely to produce asymptomatic cases, but this study contains limitations and warrants further research.⁵⁵⁶ Due to the concealment of the infection, asymptomatic patients still pose a serious threat to public health. The exact proportion of asymptomatic patients needs to be further evaluated, which will affect the effectiveness of subsequent epidemic prevention. At present, to control the further spread of SARS-CoV-2, understanding the mode of transmission in different places is important for timely detection and isolation of asymptomatic patients.

Intrafamily transmission

SARS-Cov-2 mainly spreads among family members.⁵⁵⁵ A study in the UK reported that nearly 70% of cases were transmitted within the family after extensive community control measures were taken.⁵⁵⁷ It is worth noting that both children and adults are susceptible to SARS-CoV-2 infection. Bi *et al* showed that children and the general population have a similar risk of infection (7.4% in children aged <10 years vs population average of 6.6%).⁵⁵⁸ In families, susceptible individuals have more time and opportunity for exposure to the virus; therefore, the probability of transmission in the family is higher than that in the community.⁵⁵⁹ Considering the difficulty in identifying asymptomatic patients, once a case occurs in the family, the health of the whole family is in jeopardy. In Du *et al*'s article, they reported 14 children with mild or conventional types of COVID-19; all of which were transmitted within the family.³⁸ Other studies have demonstrated that intrafamily transmission may be the main route of SARS-CoV-2 infection in children.⁵⁶⁰ It is estimated that about 75% of paediatric patients have a history of family exposure.³⁷ Moreover, Xiao *et al* have detected the presence of SARS-CoV-2 in contaminated household environments, and the virus remains viable and infectious on surfaces for up to 7 days.⁵⁶¹ The risk of transmission increases in contaminated household environments. Maintaining a clean environment, ventilation for more than 2 hours and avoiding non-biological surface transmission are key to avoiding home transmission.

Hospital transmission

Hospitals are high-risk areas. Effective measures must be taken to protect medical staff and prevent nosocomial infections. Examinations in dermatology, ophthalmology, head and neck surgery are considered to be high-risk procedures for transmission of SARS-CoV-2 from asymptomatic patients.⁵⁶² Screening of patients and hospital staff for atypical symptoms is recommended. Laboratory tests should also be conducted when necessary. Frequent testing and rapid reporting of results are particularly important to

reduce hospital-acquired infections. For patients with any suspected symptoms or histories of contact with suspected cases, medical personnel should wear full personal protective equipment to evaluate whether patients should be referred for treatment. Unnecessary surgery should be postponed.⁵⁶³ For essential surgery with a high risk of infection, operating room personnel should study the surgical pandemic guidelines and take special precautions to prevent intraoperative aerosols.⁵⁶⁴

Blood safety is also a factor worth considering in blood transfusion. SARS-CoV and Middle East respiratory syndrome virus RNA has been found in blood samples.^{565 566} Even in respiratory diseases, viral shedding in plasma and serum is common.⁵⁶⁷ For this problem, blood centres should conduct rigorous testing and detailed epidemiological investigations of blood donors to rule out the risk of asymptomatic infections. Blood samples obtained from suspected patients with COVID-19 should be recalled. Blood samples can be subjected to pathogen inactivation with photochemical treatment (PCT), if necessary; however, the use of PCT is limited because of its damage to blood components and high cost.⁵⁶⁸

Due to the unique physical condition of pregnant women, testing strategies based on currently known symptoms may miss pregnant women infected with COVID-19, posing a risk to hospitals. Timely detection of pregnant women with COVID-19 and timely preventive measures are important for the protection of pregnant women, parents, newborns, other patients and hospital staff.⁵⁶⁹ There is currently no evidence that SARS-CoV-2 can use breast milk as a medium.⁵⁷⁰ However, the shedding of the virus in asymptomatic women may also affect the treatment of newborns.⁵⁷¹ Women who are asymptomatic or have mild symptoms after delivery are recommended to breast feed directly under strict infection control measures.⁵⁶⁹

Transmission of pollutants from patients

Current statistics show that the positive rate of the SARS-CoV-2 nucleic acid test for nasopharyngeal swabs is 38.13%.⁵⁷² Due to the existence of false-negative results, a negative test does not mean that there is no viral on the swab.⁵⁷³ Given this, biological specimen handling is essential for the prevention of laboratory-related infections. Besides, even after the results of respiratory samples in patients have turned negative, faecal samples remain positive.⁵⁷⁴ A meta-analysis indicated that SARS-CoV-2 RNA was detected in the stool samples of 48.1% of patients. Even after the nasopharyngeal test is negative, the test result may still be positive.⁵⁷⁵ Virus shedding from the digestive system may be more severe and last longer than from the respiratory tract.⁵⁷⁴ Especially in asymptomatic children, this phenomenon can last up to 54 days after admission or 30 days after discharge.⁵⁷⁶ Although experts believe that a positive SARS-CoV-2 nucleic

acid test from a rectal swab does not indicate the presence of live viruses, faecal–oral transmission may still occur.^{S70} Long-term excretion of viruses through the intestine has been proved in children, reminding us to focus on the management of faeces of patients with COVID-19, especially children.^{S77} Therefore, special disinfection of used swabs and patient excreta is necessary. Additionally, since it is arduous to determine the presence of asymptomatic patients before laboratory testing, masks used by the public should also be disinfected daily. Global guidelines for the handling of biological samples and contaminants should also be drawn up as soon as possible so that effective disposal can be carried out safely and securely while protecting handlers.

The mutation of SARS-CoV-2

SARS-CoV-2 has a certain mutation frequency as RNA viruses since the epidemic of COVID-19. Data published by the WHO indicate that there are four major mutations currently. The D614G mutation in the SARS-CoV-2 has become the main type and transmitted globally by June 2020.^{S78} The ‘Cluster 5’ variant does not appear to be widely disseminated.^{S79} The VOC 202012/01 variant is more transmissible, but the severity of disease (as assessed by length of hospitalisation and 28-day mortality) and reinfection have not changed.^{S80} Preliminary studies suggest that the 501Y.V2 variant is associated with a higher viral load and may increase infectivity, but there is no evidence to prove that it causes more severe disease yet. So further investigation is needed.^{S81}

A number of countries have seen the recurrence of cured cases or rapid spread of the disease recently. It may indicate the spread of the SARS-CoV-2 variant to some extent with the involvement of asymptomatic carriers.^{S82}

Recently, a variant of SARS-CoV-2, SARS-CoV-2 GZ69, was isolated from an asymptomatic healthcare worker. It was found that the mutation of residues 203–204 in SARS-CoV-2 GZ69 may reduce the anti-proliferation ability of N protein, which may explain the virus persistence in asymptomatic patients.^{S83} Fabrizio Maggi reported an asymptomatic traveller from Brazil who tested positive for the SARS-CoV-2 P.1 variant in a screening nasopharyngeal swab sample. The results suggest that this may lead to a wider circulation of SARS-CoV-2 variant P.1 in areas other than the Brazilian state of Amazonas.^{S84} We recommend that strict surveillance measures are necessary, because asymptomatic patients may lead to further transmission of SARS-CoV-2 variant. Moreover, designing specific primer sets for viral variants is useful to assist in large-scale screening.

Screening of discharged patients and persons released from isolation
According to the *Regulations for the Administration of Asymptomatic Infected Patients of New Coronavirus*

issued in China, patients who are asymptomatic and their close contacts should be quarantined and observed for 14 days. Only after 14 days and two consecutive negative nucleic acid tests (sampling interval ≥ 24 hours), can medical observation be discontinued. For convenience and non-invasiveness, throat swab samples are generally collected for nucleic acid detection.^{S85} However, Xu *et al* found that after nasopharyngeal swabs turn negative, rectal swabs may still show a positive result.^{S86} The timing of viral nucleic acid shedding differs among tissue types. A meta-analysis proved that PCR using sputum samples to diagnose COVID-19 is more sensitive than stool samples.^{S87} Rectal swabs are not routinely used in the diagnosis of COVID-19 in many countries. The current clinical application cases are insufficient. Further studies are still needed to confirm its feasibility. Besides, rehabilitation patients may also experience intermittent viral shedding. Some people who have reached the discharge standard may still be carriers.^{S88} We recommend 14 days of medical observation after reaching discharge standards to prevent the possibility of the result being positive again due to intermittent virus shedding.

Mass screening

At present, to prevent the spread of SARS-CoV-2 among travellers, symptom-based screening is often set up at airports, railway stations and ports. Due to their rapidity and ease of use, non-contact infrared thermometers are widely used in this type of screening.^{S88} However, as more asymptomatic infections are discovered and some patients deliberately hide symptoms, the effectiveness of fever screening has major challenges. The proportion of asymptomatic patients is a key factor for further study, which can seriously affect the effectiveness of screening.^{S89} Rapid identification, registration, reporting, supervising and epidemiological tracking of asymptomatic patients are significant. Special attention should be paid to screening the population who need to return to work and production to prevent ‘import cases’ between regions. The use of thermal scanners and other symptom detection equipment may improve the efficiency and reliability of screening.^{S90} Setting up multiple sites to filter people multiple times is also worth considering.^{S91} Moreover, recent studies have shown that nucleic acid testing using saliva samples is highly sensitive, especially for recognising patients in the early stages of the disease.^{S92} The ease of saliva collection makes self-collection of saliva samples a potentially effective method for large-scale screening of asymptomatic populations. Developing nucleic acid or antibody detection devices that are more suitable for screening at these sites remains the best option for improving the accuracy and sensitivity of the tests.^{S93} In addition to traveller screening, some countries, such as South Korea, Iceland and Germany have also conducted universal nucleic acid

testing in the population to detect patients and assess the proportion of asymptomatic patients.^{S94 S95} The risk of transmission of asymptomatic infections is multifaceted. The concealment of transmission makes it difficult to prevent. It seems that the most reliable method of prevention currently is to identify patients with symptoms and track their movements in real time. Many new internet-based technologies have been used to enable the detection of close contacts. Self-reporting based on smartphones and continuous tracking with the code can achieve inexpensive and extensive mass tracking.^{S96} The establishment of the rapid suspicious case notification platform is also conducive to the self-discovery of close contacts, reminding them to actively seek detection. These measures will contribute to the discovery of asymptomatic patients.

CONCLUSION

The emergence of asymptomatic patients with COVID-19 poses new challenges for infection prevention and control. More work is needed to clarify the exact proportion of asymptomatic patients among the overall total. Available data found significant differences in the proportion of asymptomatic patients diagnosed in different populations. It has been found that asymptomatic patients have the potential for transmission. For the discovery of asymptomatic carriers, the main existing detection methods include imaging, nucleic acid and serological diagnostics. Joint detection methods can improve the accuracy of early diagnosis. To slow down or stop the spread of SARS-CoV-2, our top priority is to take more effective prevention and control measures at home, in hospitals and elsewhere, and find more appropriate methods for population screening. For different groups of people such as children, pregnant women and the elderly, different prevention and treatment measures should be taken according to their specific symptoms and different transmission capabilities. For the public, measures such as wearing face masks, maintaining physical distance, washing hands frequently and not gathering are still the most desirable protective measures.

Contributors YY, XY, DH, QY, TW and MD contributed to the material collection and manuscript writing. All authors read and approved the final manuscript.

Funding This work was supported by the National Natural Science Foundation of China under contract (Nos. 31672290, 31100764), the Natural Science Foundation of Hunan Province, China (No. 2016JJ3180), Open Sharing Fund for the Large-scale Instruments and Equipments of Central South University (No. CSUZC2020043), National Undergraduate Innovation Training Program of Central South University (No. S2020105330717).

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Ting Wu <http://orcid.org/0000-0003-1708-2869>

REFERENCES

- Kang S, Peng W, Zhu Y, *et al.* Recent progress in understanding 2019 novel coronavirus (SARS-CoV-2) associated with human respiratory disease: detection, mechanisms and treatment. *Int J Antimicrob Agents* 2020;55:105950.
- WHO Coronavirus Disease (COVID-19) Dashboard. Available: <https://who.sprinklr.com/> [Accessed 25 Feb 2021].
- Wang M, Wu T, Zuo Z, *et al.* Evaluation of current medical approaches for COVID-19: a systematic review and meta-analysis. *BMJ Support Palliat Care* 2021;11:45–52.
- He J, Guo Y, Mao R, *et al.* Proportion of asymptomatic coronavirus disease 2019: a systematic review and meta-analysis. *J Med Virol* 2021;93:820–30.
- Yanes-Lane M, Winters N, Fregonese F, *et al.* Proportion of asymptomatic infection among COVID-19 positive persons and their transmission potential: a systematic review and meta-analysis. *PLoS One* 2020;15:e0241536.
- Pan Y, Yu X, Du X, *et al.* Epidemiological and clinical characteristics of 26 asymptomatic severe acute respiratory syndrome coronavirus 2 carriers. *J Infect Dis* 2020;221:1940–7.
- Wu T, Zuo Z, Kang S, *et al.* Multi-Organ dysfunction in patients with COVID-19: a systematic review and meta-analysis. *Aging Dis* 2020;11:874–94.
- Chen C, Zhu C, Yan D, *et al.* The epidemiological and radiographical characteristics of asymptomatic infections with the novel coronavirus (COVID-19): a systematic review and meta-analysis. *Int J Infect Dis* 2021;104:458–64.
- Meng H, Xiong R, He R, *et al.* Ct imaging and clinical course of asymptomatic cases with COVID-19 pneumonia at admission in Wuhan, China. *J Infect* 2020;81:e33–e39:e33–9.
- Taleghani N, Taghipour F. Diagnosis of COVID-19 for controlling the pandemic: a review of the state-of-the-art. *Biosens Bioelectron* 2021;174:112830.
- Setti L, Kirienko M, Dalto SC, *et al.* Fdg-Pet/Ct findings highly suspicious for COVID-19 in an Italian case series of asymptomatic patients. *Eur J Nucl Med Mol Imaging* 2020;47:1649–56.
- Udugama B, Kadhiresan P, Kozlowski HN, *et al.* Diagnosing COVID-19: the disease and tools for detection. *ACS Nano* 2020;14:3822–35.
- Kim H, Hong H, Yoon SH. Diagnostic performance of CT and reverse transcriptase polymerase chain reaction for coronavirus disease 2019: a meta-analysis. *Radiology* 2020;296:E145–55.
- Shah AS, Walkoff LA, Kuzo RS, *et al.* The utility of chest computed tomography (CT) and RT-PCR screening of asymptomatic patients for SARS-CoV-2 prior to semiurgent or urgent Hospital procedures. *Infect Control Hosp Epidemiol* 2020;41:1375–7.
- Ng BH, Nuratiqah NA, Andrea YLB, *et al.* Lung computed tomography patterns of a cluster of asymptomatic young males with COVID-19 admitted to a teaching hospital in Kuala Lumpur. *Med J Malaysia* 2020;75:368–71.
- Yoon S, Li H, Lee KH, *et al.* Clinical characteristics of asymptomatic and symptomatic pediatric coronavirus disease 2019 (COVID-19): a systematic review. *Medicina* 2020;56:474.
- Fan L, Liu S. CT and COVID-19: Chinese experience and recommendations concerning detection, staging and follow-up. *Eur Radiol* 2020;30:5214–6.
- Yüce M, Filiztekin E, Özkaya KG. COVID-19 diagnosis -A review of current methods. *Biosens Bioelectron* 2021;172:112752.
- Wang X, Tan L, Wang X, *et al.* Comparison of nasopharyngeal and oropharyngeal swabs for SARS-CoV-2 detection in 353

- patients received tests with both specimens simultaneously. *Int J Infect Dis* 2020;94:107–9.
- 20 Nagura-Ikeda M, Imai K, Tabata S, *et al.* Clinical evaluation of self-collected saliva by quantitative reverse transcription-PCR (RT-qPCR), direct RT-qPCR, reverse transcription-loop-mediated isothermal amplification, and a rapid antigen test to diagnose COVID-19. *J Clin Microbiol* 2020;58. doi:10.1128/JCM.01438-20. [Epub ahead of print: 24 Aug 2020].
 - 21 Zhu X, Wang X, Han L, *et al.* Multiplex reverse transcription loop-mediated isothermal amplification combined with nanoparticle-based lateral flow biosensor for the diagnosis of COVID-19. *Biosens Bioelectron* 2020;166:112437.
 - 22 Li Y, He Q, Yu R, *et al.* Highlighted prospects of an IgM/IgG antibodies test in identifying individuals with asymptomatic severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. *Arch Pathol Lab Med* 2021;145:39–45.
 - 23 Li Z, Yi Y, Luo X, *et al.* Development and clinical application of a rapid IgM-IgG combined antibody test for SARS-CoV-2 infection diagnosis. *J Med Virol* 2020;92:1518–24.
 - 24 Dong X, Cao Y-Y, Lu X-X, *et al.* Eleven faces of coronavirus disease 2019. *Allergy* 2020;75:1699–709.
 - 25 Wellinghausen N, Voss M, Ivanova R, *et al.* Evaluation of the SARS-CoV-2-IgG response in outpatients by five commercial immunoassays. *GMS Infect Dis* 2020;8:p. Doc22.
 - 26 Wang B, Wang L, Kong X, *et al.* Long-Term coexistence of SARS-CoV-2 with antibody response in COVID-19 patients. *J Med Virol* 2020;92:1684–9.
 - 27 Ai T, Yang Z, Hou H, *et al.* Correlation of chest CT and RT-PCR testing for coronavirus disease 2019 (COVID-19) in China: a report of 1014 cases. *Radiology* 2020;296:E32–40.
 - 28 Wang Y, Hou H, Wang W, *et al.* Combination of CT and RT-PCR in the screening or diagnosis of COVID-19. *J Glob Health* 2020;10:010347.
 - 29 Wu X, Fu B, Chen L, *et al.* Serological tests facilitate identification of asymptomatic SARS-CoV-2 infection in Wuhan, China. *J Med Virol* 2020;92:1795–6.
 - 30 Zhang W, Du R-H, Li B, *et al.* Molecular and serological investigation of 2019-nCoV infected patients: implication of multiple shedding routes. *Emerg Microbes Infect* 2020;9:386–9.
 - 31 Buitrago-Garcia D, Egli-Gany D, Counotte MJ, *et al.* Occurrence and transmission potential of asymptomatic and presymptomatic SARS-CoV-2 infections: a living systematic review and meta-analysis. *PLoS Med* 2020;17:e1003346.
 - 32 Le TQM, Takemura T, Moi ML, *et al.* Severe Acute Respiratory Syndrome Coronavirus 2 Shedding by Travelers, Vietnam, 2020. *Emerg Infect Dis* 2020;26:1624–6.
 - 33 Zou L, Ruan F, Huang M, *et al.* SARS-CoV-2 viral load in upper respiratory specimens of infected patients. *N Engl J Med* 2020;382:1177–9.
 - 34 Madewell ZJ, Yang Y, Longini IM, *et al.* Household transmission of SARS-CoV-2: a systematic review and meta-analysis. *JAMA Netw Open* 2020;3:e2031756.
 - 35 Qiu X, Nergiz AI, Maraolo AE, *et al.* Defining the role of asymptomatic and pre-symptomatic SARS-CoV-2 transmission - a living systematic review. *Clin Microbiol Infect* 2021. doi:10.1016/j.cmi.2021.01.011. [Epub ahead of print: 20 Jan 2021].
 - 36 Koh WC, Naing L, Chaw L, *et al.* What do we know about SARS-CoV-2 transmission? A systematic review and meta-analysis of the secondary attack rate and associated risk factors. *PLoS One* 2020;15:e0240205.
 - 37 Chang T-H, Wu J-L, Chang L-Y. Clinical characteristics and diagnostic challenges of pediatric COVID-19: a systematic review and meta-analysis. *J Formos Med Assoc* 2020;119:982–9.
 - 38 Du W, Yu J, Wang H, *et al.* Clinical characteristics of COVID-19 in children compared with adults in Shandong Province, China. *Infection* 2020;48:445–52.
 - 39 Li B, Zhang S, Zhang R, *et al.* Epidemiological and clinical characteristics of COVID-19 in children: a systematic review and meta-analysis. *Front Pediatr* 2020;8:591132.
 - 40 Dong Y, Mo X, Hu Y, *et al.* Epidemiology of COVID-19 among children in China. *Pediatrics* 2020;145:e20200702.
 - 41 Zhou M-Y, Xie X-L, Peng Y-G, *et al.* From SARS to COVID-19: what we have learned about children infected with COVID-19. *Int J Infect Dis* 2020;96:710–4.
 - 42 Peterson N, Sagdeo K, Tyungu D, *et al.* Discovering associations: Kawasaki disease and COVID-19. *Case Rep Pediatr* 2020;2020:1–3.
 - 43 Pettiroso E, Giles M, Cole S, *et al.* COVID-19 and pregnancy: a review of clinical characteristics, obstetric outcomes and vertical transmission. *Aust N Z J Obstet Gynaecol* 2020;60:640–59.
 - 44 Noor A, Krilov LR. An overview of the ongoing clinical issues of COVID-19. *Pediatr Ann* 2021;50:e84–9.
 - 45 Chen H, Guo J, Wang C, *et al.* Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet* 2020;395:809–15.
 - 46 Di Mascio D, Sen C, Saccone G, *et al.* Risk factors associated with adverse fetal outcomes in pregnancies affected by coronavirus disease 2019 (COVID-19): a secondary analysis of the WAPM study on COVID-19. *J Perinat Med* 2020;48:950–8.
 - 47 Verma S, Bradshaw C, Auyeung NSF, *et al.* Outcomes of Maternal-Newborn dyads after maternal SARS-CoV-2. *Pediatrics* 2020;146. doi:10.1542/peds.2020-005637. [Epub ahead of print: 31 Jul 2020].
 - 48 Molina LP, Chow S-K, Nickel A, *et al.* Prolonged detection of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) RNA in an obstetric patient with antibody seroconversion. *Obstet Gynecol* 2020;136:838–41.
 - 49 Tanacan A, Erol SA, Turgay B, *et al.* The rate of SARS-CoV-2 positivity in asymptomatic pregnant women admitted to hospital for delivery: experience of a pandemic center in turkey. *Eur J Obstet Gynecol Reprod Biol* 2020;253:31–4.
 - 50 Dai J, Yang L, Zhao J. Probable longer incubation period for elderly COVID-19 cases: analysis of 180 contact tracing data in Hubei Province, China. *Risk Manag Healthc Policy* 2020;13:1111–7.

Additional references 51–96

- S51 Yu C, Zhou M, Liu Y, et al. Characteristics of asymptomatic COVID-19 infection and progression: a multicenter, retrospective study. *Virulence* 2020;11:1006–14.
- S52 Sacco G, Foucault G, Briere O, et al. COVID-19 in seniors: findings and lessons from mass screening in a nursing home. *Maturitas* 2020;141:46–52.
- S53 Ra SH, Lim JS, Kim G-U, et al. Upper respiratory viral load in asymptomatic individuals and mildly symptomatic patients with SARS-CoV-2 infection. *Thorax* 2021;76:61–3.
- S54 Hung IF-N, Cheng VC-C, Li X, et al. SARS-CoV-2 shedding and seroconversion among passengers quarantined after disembarking a cruise SHIP: a case series. *Lancet Infect Dis* 2020;20:1051–60.
- S55 He D, Zhao S, Lin Q, et al. The relative transmissibility of asymptomatic COVID-19 infections among close contacts. *Int J Infect Dis* 2020;94:145–7.
- S56 Chen Y, Wang AH, Yi B, et al. [Epidemiological characteristics of infection in COVID-19 close contacts in Ningbo city]. *Zhonghua Liu Xing Bing Xue Za Zhi* 2020;41:667–71.
- S57 Haroon S, Chandan JS, Middleton J, et al. Covid-19: breaking the chain of household transmission. *BMJ* 2020;370:m3181.
- S58 Li W et al. The characteristics of household transmission of COVID-19. *Clin Infect Dis* 2020:ciaa450.
- S59 Su L, Ma X, Yu H, et al. The different clinical characteristics of corona virus disease cases between children and their families in China - the character of children with COVID-19. *Emerg Microbes Infect* 2020;9:707–13.
- S60 Tan Y-P, Tan B-Y, Pan J, et al. Epidemiologic and clinical characteristics of 10 children with coronavirus disease 2019 in Changsha, China. *J Clin Virol* 2020;127:104353.
- S61 Xiao F, Chen B, Xiao T, et al. Children with SARS-CoV-2 infection during an epidemic in China (outside of Hubei Province). *Ann Transl Med* 2020;8:849.
- S62 Goldman RA, Swendseid B, Chan JYK, et al. Tracheostomy management during the COVID-19 pandemic. *Otolaryngol Head Neck Surg* 2020;163:67–9.
- S63 Langer PD, Bernardini FP. Oculofacial Plastic Surgery and the COVID-19 Pandemic: Current Reactions and Implications for the Future. *Ophthalmology* 2020;127:e70–1.

S64 Diaz A, Sarac BA, Schoenbrunner AR, et al. Elective surgery in the time of COVID-19. *Am J Surg* 2020;219:900–2.

S65 Ng EKO, Hui DS, Chan KCA, et al. Quantitative analysis and prognostic implication of SARS coronavirus RNA in the plasma and serum of patients with severe acute respiratory syndrome. *Clin Chem* 2003;49:1976–80.

S66 Ng EKO, Ng P-C, Hon KLE, et al. Serial analysis of the plasma concentration of SARS coronavirus RNA in pediatric patients with severe acute respiratory syndrome. *Clin Chem* 2003;49:2085–8.

S67 Corman VM, Albarak AM, Omrani AS, et al. Viral shedding and antibody response in 37 patients with middle East respiratory syndrome coronavirus infection. *Clin Infect Dis* 2016;62:477–83.

S68 Chang L, Yan Y, Wang L. Coronavirus disease 2019: coronaviruses and blood safety. *Transfus Med Rev* 2020;34:75–80.

S69 Søreide K, Hallet J, Matthews JB, et al. Immediate and long-term impact of the COVID-19 pandemic on delivery of surgical services. *Br J Surg* 2020;107:1250–61.

S70 Davanzo R, Moro G, Sandri F, et al. Breastfeeding and coronavirus disease-2019: Ad interim indications of the Italian Society of Neonatology endorsed by the Union of European Neonatal & Perinatal Societies. *Matern Child Nutr* 2020;16:e13010.

S71 Breslin N, Baptiste C, Gyamfi-Bannerman C, et al. Coronavirus disease 2019 infection among asymptomatic and symptomatic pregnant women: two weeks of confirmed presentations to an affiliated pair of new York City hospitals. *Am J Obstet Gynecol MFM* 2020;2:100118.

S72 Wu J, Liu J, Li S, et al. Detection and analysis of nucleic acid in various biological samples of COVID-19 patients. *Travel Med Infect Dis* 2020;37:101673.

S73 Li Y, Yao L, Li J, et al. Stability issues of RT-PCR testing of SARS-CoV-2 for hospitalized patients clinically diagnosed with COVID-19. *J Med Virol* 2020;92:903–8.

S74 Tahamtan A, Ardebili A. Real-Time RT-PCR in COVID-19 detection: issues affecting the results. *Expert Rev Mol Diagn* 2020;20:453–4.

S75 Cheung KS, Hung IFN, Chan PPY, et al. Gastrointestinal manifestations of SARS-CoV-2 infection and virus load in fecal samples from a Hong Kong cohort: systematic review and meta-analysis. *Gastroenterology* 2020;159:81–95.

- S76 Xie J, Long X, Ren C, et al. Follow-Up study of long-time positive RT-PCR in stool specimens from asymptomatic children infected with SARS-CoV-2. *Pediatr Infect Dis J* 2020;39:e315–7.
- S77 Yeo C, Kaushal S, Yeo D. Enteric involvement of coronaviruses: is faecal-oral transmission of SARS-CoV-2 possible? *Lancet Gastroenterol Hepatol* 2020;5:335–7.
- S78 WHO. SARS-CoV-2 variants. Available: <http://www.who.int/csr/don/31-december-2020-sars-cov2-variants/en/> [Accessed 25 Feb 2021].
- S79 Hammer AS, Quaade ML, Rasmussen TB, et al. SARS-CoV-2 transmission between mink (*Neovison vison*) and humans, Denmark. *Emerg Infect Dis* 2021;27:547–51.
- S80 Leung K, Shum MH, Leung GM, et al. Early transmissibility assessment of the N501Y mutant strains of SARS-CoV-2 in the United Kingdom, October to November 2020. *Euro Surveill* 2021;26.
- S81 Tang JW, Toovey OTR, Harvey KN, et al. Introduction of the South African SARS-CoV-2 variant 501Y.V2 into the UK. *J Infect* 2021. doi:10.1016/j.jinf.2021.01.007. [Epub ahead of print: 17 Jan 2021].
- S82 Rahimi F, Talebi Bezmin Abadi A. Implications of the emergence of a new variant of SARS-CoV-2, VUI-202012/01. *Arch Med Res* 2021. doi:10.1016/j.arcmed.2021.01.001. [Epub ahead of print: 22 Jan 2021].
- S83 Caccuri F, Zani A, Messali S, et al. A persistently replicating SARS-CoV-2 variant derived from an asymptomatic individual. *J Transl Med* 2020;18:362.
- S84 Maggi F, Novazzi F, Genoni A, et al. Imported SARS-CoV-2 variant P.1 in traveler returning from Brazil to Italy. *Emerg Infect Dis* 2021;27:1249–51.
- S85 Lan L, Xu D, Ye G, et al. Positive RT-PCR test results in patients recovered from COVID-19. *JAMA* 2020;323:1502–3.
- S86 Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020;323:1061–9.
- S87 Böger B, Fachi MM, Vilhena RO, et al. Systematic review with meta-analysis of the accuracy of diagnostic tests for COVID-19. *Am J Infect Control* 2021;49:21–9.
- S88 Dietz L, Horve PF, Coil DA, et al. 2019 Novel Coronavirus (COVID-19) Pandemic: Built Environment Considerations To Reduce Transmission. *mSystems* 2020;5.

- S89 Xu Y, Li X, Zhu B, et al. Characteristics of pediatric SARSCoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat Med* 2020;26:502–5.
- S90 Gostic K, Gomez AC, Mummah RO, et al. Estimated effectiveness of symptom and risk screening to prevent the spread of COVID-19. *Elife* 2020;9. doi:10.7554/eLife.55570. [Epub ahead of print: 24 Feb 2020].
- S91 Aggarwal N, Garg M, Dwarakanathan V, et al. Diagnostic accuracy of non-contact infrared thermometers and thermal scanners: a systematic review and meta-analysis. *J Travel Med* 2020;27. doi:10.1093/jtm/taaa193. [Epub ahead of print: 23 Dec 2020].
- S92 Kwon KT, Ko JH, Shin H, et al. Drive-Through screening center for COVID-19: a safe and efficient screening system against massive community outbreak. *J Korean Med Sci* 2020;35:e123.
- S93 Yokota I, Shane PY, Okada K, et al. Mass screening of asymptomatic persons for SARS-CoV-2 using saliva. *Clin Infect Dis* 2020. doi:10.1093/cid/ciaa1388. [Epub ahead of print: 25 Sep 2020].
- S94 Esbin MN, Whitney ON, Chong S, et al. Overcoming the bottleneck to widespread testing: a rapid review of nucleic acid testing approaches for COVID-19 detection. *RNA* 2020;26:771–83.
- S95 Cohen J, Kupferschmidt K. Countries test tactics in 'war' against COVID-19. *Science* 2020;367:1287–8.
- S96 Gudbjartsson DF, Helgason A, Jonsson H, et al. Spread of SARS-CoV-2 in the Icelandic population. *N Engl J Med* 2020;382:2302–15.