P 104

FIVE YEARS ON: THE INCREASING USE OF ULTRASOUND IN A HOSPICEBernadette Lee. *Princess Alice Hospice, Esher, UK*

10.1136/bmjspcare-2014-000654.145

Background Since 2008, several new doctors have attended the Focused Abdominal UltraSound in Palliative Care course. The course taught the examination protocols for ascites, liver, urinary tract and deep vein thrombosis (DVT). A log book was kept with the scanner.

Aims The aims were to audit:
 the frequency of use
 whether the course examination protocols were followed
 the impact of having ultrasound
 whether cleaning of the transducer was recorded
 Information from the log book and patient records were used to complete the audit.

Results In the last 3 years, the number of scans increased from 22 to 40. There was a trend for patients who might need paracentesis to be assessed before admission was requested.

All 125 scans followed examination protocols.

The clinical indications were confirmation of ascites (97), urinary bladder (14), hydronephrosis (3) and groin DVT (1).

Year	New doctors trained	Number of doctors doing scans	Number of scans	Number of patients	Location			
					ward	OPD	home	nursing home
2008	6	3	24	16	24	0	0	0
2009	0	3	15	13	15	0	0	0
2010	1	4	22	16	20	2	0	0
2011	4	7	24	20	17	2	5	0
2012	1	6	40	29	30	7	2	1
total	12	—	125	94	106	11	7	1

Each scan had been helpful in the clinical decision making: paracentesis were done on 22 occasions, prescription of diuretics (1), referral for PleurX drain (1) and insertion of urine catheter (12). 92% (115/125) log entries recorded cleaning of transducer.

Conclusions There has been increasing and appropriate use of ultrasound last 5 years. The use of inpatient beds were optimised. Cleaning of the transducer needs to be done each time and recorded. Annual audit of ultrasound use is planned.