OPEN ACCESS

Prospective comparative effectiveness cohort study comparing two models of advance care planning provision for Australian community aged care clients

Karen Margaret Detering,^{1,2} Rachel Zoe Carter,¹
 Marcus William Sellars,^{1,3} Virginia Lewis,⁴ Elizabeth Anne Sutton¹

¹Advance Care Planning Australia, Austin Health, Heidelberg, Victoria, Australia

²Faculty of Medicine, Dentistry and Health Sciences, Melbourne University, Melbourne, Victoria, Australia

³Kolling Institute, Northern Clinical School, The University of Sydney, Camperdown, New South Wales, Australia

⁴Australian Institute for Primary Care and Ageing, La Trobe University, Melbourne, Victoria, Australia

Correspondence to

Dr Karen Margaret Detering, Advance Care Planning Australia, Austin Health, Heidelberg, Victoria, Australia; karen.detering@austin.org.au

Received 1 May 2017

Revised 20 July 2017

Accepted 30 August 2017

Published Online First

16 September 2017

ABSTRACT

Objectives Conduct a prospective comparative effectiveness cohort study comparing two models of advance care planning (ACP) provision in community aged care: ACP conducted by the client's case manager (CM) ('Facilitator') and ACP conducted by an external ACP service ('Referral') over a 6-month period.

Methods This Australian study involved CMs and their clients. Eligible CM were English speaking, ≥ 18 years, had expected availability for the trial and worked ≥ 3 days per week. CMs were recruited via their organisations, sequentially allocated to a group and received education based on the group allocation. They were expected to initiate ACP with all clients and to facilitate ACP or refer for ACP. Outcomes were quantity of new ACP conversations and quantity and quality of new advance care directives (ACDs).

Results 30 CMs (16 Facilitator, 14 Referral) completed the study; all 784 client's files (427 Facilitator, 357 Referral) were audited. ACP was initiated with 508 (65%) clients (293 Facilitator, 215 Referral; $p < 0.05$); 89 (18%) of these (53 Facilitator, 36 Referral) and 41 (46%) (13 Facilitator, 28 Referral; $p < 0.005$) completed ACDs. Most ACDs (71%) were of poor quality/notvalid. A further 167 clients (facilitator 124; referral 43; $p < 0.005$) reported ACP was in progress at study completion.

Conclusions While there were some differences, overall, models achieved similar outcomes. ACP was initiated with 65% of clients. However, fewer clients completed ACP, there was low numbers of ACDs and document quality was generally poor. The findings raise

questions for future implementation and research into community ACP provision.

INTRODUCTION

Advance care planning (ACP) is a coordinated communication process between a person, their family/carer(s) and healthcare providers and aims to clarify the person's values, treatment preferences and goals of medical treatment should the person lose capacity to make or communicate such decisions in the future.¹ In Australia, formal ACP programmes usually operate within health, institutional or aged care settings and involve trained staff.¹ These programmes are often located at public health services. While discussions are the main focus, an important and often desirable outcome of ACP is the completion of a written advance care directive (ACD) that documents the person's preferences and/or the appointment of a substitute decision maker.¹ ACP has been shown to improve care, including end-of-life care,^{2,3} to improve the likelihood that a person's preferences will be known and respected²⁻⁴ and to improve the psychological outcomes in surviving relatives.^{2,4}

The Australian Government Home Care Package (HCP) programme provides funding for personal/health/nursing support to frail or unwell Australians to assist them to remain at home rather than enter residential care.⁵ HCPs assisted 60 000 people in 2013; this number is expected to increase to 100 000 by

CrossMark

To cite: Detering KM, Carter RZ, Sellars MW, et al. *BMJ Supportive & Palliative Care* 2017;7:486–494.

2016/2017, with predictions that 80% of aged care services will be delivered in this form by 2050.⁵ Each client is assigned a case manager who, in partnership with the client, coordinates the services that the client receives.⁵ There are four levels of HCP support based on needs ranging from level 1 (basic) to level 4 (high).⁵

ACP may be both beneficial and achievable for HCP clients. Older people believe ACP is important, and they are enthusiastic about participation.⁶ Although most HCP clients are elderly, frail and have chronic illnesses, they are more likely to have decision-making capacity than residential care residents^{7,8} and thus more able to participate in ACP. Additionally, care providers visit clients in their homes, possibly a more amenable environment for ACP conversations than either aged care facilities or hospitals.⁹ It is hypothesised that ACP discussions while the client is at home and likely to be stable is more appropriate and timely than during hospital admissions, when the person may be too unwell to undertake ACP. Evidence shows that preferences for life-sustaining treatments are dependent on the context in which they are made, with a 'hospitalisation dip' during and immediately following admission, and return of preferences to their prehospitalisation status after some time, thus supporting the notion of conducting ACP when clinically stable.¹⁰

Despite the perceived value of ACP for HCP clients,^{11–14} limited international and Australian research has shown that generally the uptake of ACP within community care is low^{12,13,15–17} and that case managers vary in their ACP practice, knowledge and attitudes.^{12,15,17} These studies show lack of a systematic approach to ACP within services that most services do not provide ACP training for staff, that minimal resources are allocated to ACP and that the quality and prevalence of ACP varies between service providers.^{12,13,15–17}

Given the increase in community-based aged care programmes such as HCP, and the recognised value of undertaking ACP earlier and prior to loss of decision-making capacity, there is a strong impetus to provide ACP in this setting. However, how best to provide ACP in this setting is unknown. We conducted a prospective comparative effectiveness cohort study comparing two models of ACP provision to HCP clients:

1. ACP conducted by the client's case manager ('Facilitator' model), based on previous ACP implementation in other settings^{2,18}
2. ACP provided by referral to external ACP services ('Referral' model). This model is based on Australian research showing current practice within HCP services includes ACP initiation, followed by referral to complete ACP.¹⁷

The aim of this study was to compare the effectiveness of these two distinct models of ACP provision, in terms of quantity of ACP discussions and quantity and

quality of ACD completion, over a 6-month intervention period.

METHODS

Study design

This study was conducted in Victoria, Australia. Eligible participants were case managers. HCP services were recruited via an information session and invited their case managers to participate. Eligible case managers were ≥ 18 years, English speaking and expected to be available ≥ 3 days per week for the 6-month intervention period. Participants, provided consent, were stratified by site and sequentially allocated to Facilitator or Referral groups. No data were collected regarding the number of case managers not invited to participate.

Recruitment occurred in August 2014, the intervention period was 1 September 2014 to 28 February 2015, and data collection occurred during March 2015.

Models of ACP provision

Two models of ACP provision were utilised: the Facilitator model, based on previous ACP implementation in other health settings,^{2,18} and the Referral model, based on the findings from Australian research showing current practice includes initiation of ACP and then referral for ACP completion.¹⁷ Details of these models are outlined in [table 1](#).

The expectations of the Facilitator case managers were that they would initiate and conduct ACP with their clients including ACD completion. These case managers received training based on Austin Health's model of ACP facilitator training.^{19,20} ([table 1](#)) The expectations of the Referral case managers were that they would initiate ACP with clients, and then refer them to a specialist service for ACP discussions and ACD completion.

HCP services received remuneration for the case manager's time as follows:

Facilitator group: \$A5000 per full-time case manager for time required for initiating and conducting ACP and providing study data.

Referral group: \$A2500 per full-time case manager for time required for initiating ACP, referral to a specialist service, and providing study data.

Case managers in both groups were asked to initiate ACP with all clients during the intervention period. ACP initiation was defined as raising the topic and explaining what it is and possible benefits. Case managers provide written information about ACP and assessed interest in further ACP discussions. Where the client was interested, the case manager would facilitate ACP or refer the person to a specialist service depending on the group allocation. Where clients had previously undertaken ACP, a review of ACP was offered and either conducted or referred.

Case managers recorded details, including duration, of all ACP activity during the intervention period.

Table 1 Facilitator and Referral models of advance care planning (ACP) provision

	Facilitator model of ACP	Referral model of ACP
Description of model	Case managers are trained to facilitate ACP with their clients. They will conduct the ACP discussions and complete advance care directives (ACDs) (where appropriate).	ACP is conducted by an ACP facilitator at a specialist service located at public health services and use trained non-medical facilitators (usually nurses) to conduct ACP. Each service determines how their ACP facilitators are trained. ACP is provided free of charge.
Potential advantages	Case managers have ongoing relationships with clients. Case managers are part of the usual care team. ACP is conducted in the client's home.	Clients have access to expert facilitators. Specific time is allocated for ACP.
Potential disadvantages	Case managers need to find time within current workloads and roles. Case managers may not be expert enough as ACP facilitators	Client needs to travel to appointments. Clients do not have existing relationships with facilitators. The facilitator is not part of the client's usual care team.
Training provided	Three-module learning package (ACP theory, ethics/law, doing ACP) expected to take 2 hours to complete. This is completed prior to the an 8-hour (face-to-face) experiential workshop consisting of facilitated discussion and role play. The workshop's primary focus is 'how to have the conversation'. During the workshop, participants were expected to complete a full ACP discussion with another participant and document the outcome in an ACD.	Two-hour face-to-face training session, providing an overview of ACP, interactive training in initiation of ACP and instructions on how to refer clients and follow-up requirements for clients who attended an ACP service.

ACD, advance care directive.

Telephone mentoring by project staff was provided to case managers in both groups.

Audit of client files

At study completion, all client files of the participating case managers were audited. This included clients who had commenced or ceased receiving services during the period. Using a specifically designed template, researchers and case managers, or another HCP organisational representative, examined the files together to extract the required information. Data collected included demographics, detail of pre-existing (prior to intervention) ACP/ACDs and detail of new ACP discussions and ACDs. Researchers did not directly contact clients or have access to client identifying information.

A completed conversation was defined as one or more discussions (with/without completion of ACD) with no wish from the client for further discussion.

Document audit

Deidentified copies of all newly completed ACDs were assessed for quality and validity. These ACDs included statutory documents for appointing substitute decision makers (medical enduring power of attorney) and documents outlining preferences, either statutory (Refusal of Treatment Certificate (RTC)) or common law ACDs (CL-ACD).

Two researchers independently assessed ACDs for compliance with best practice and validity, using a specifically developed tool, based on previous research.¹⁸ Disagreement was resolved by discussion until consensus was achieved. Parameters included whether: the client's name was clearly documented; only one cardiopulmonary resuscitation (CPR) preference was chosen (where relevant); only one life-prolonging treatment option was selected (where relevant) and this was compatible with the CPR option; that

the document was signed by the client (or substitute decision maker if client was non-competent); and that witnessing requirements were met.

Outcome measures

These were the quantity of ACP discussions and the outcomes of these discussions including quantity and quality of ACDs completed and reported pre-existing ACP activity.

Statistical methods

Continuous data were assessed for normal distribution using a Kolmogorov-Smirnov statistic with a Lilliefors significance level for testing normality. Normally distributed data are reported as mean±SD and where appropriate hypothesis testing was performed using Student's t-test. Continuous data not normally distributed are reported as median±IQR and where appropriate hypothesis testing was performed using a Mann-Whitney U test. Categorical data are reported as median±IQR or frequency plus percentage. Statistical testing was performed using Pearson χ^2 test. A p value of ≤0.05 was considered statistically significant.

RESULTS

Case manager and client demographics

Thirty-five case managers were recruited and trained; 19 were allocated to Facilitator group and 16 to Referral group. Five case managers subsequently withdrew (four changed jobs, one went on extended leave). Thus, data for 30 case managers (16 Facilitator, 14 Referral), and their clients, are included.

There were no significant demographic differences between Facilitator and Referral case managers (table 2), and they are similar to that of Australian case managers^{17 21} in that their median age was 47 years; most were female, were born in Australia, New

Table 2 CM demographics and client load

	Case managers			p=
	Overall n=30, n (%)	Facilitator n=16, n (%)	Referral n=14, n (%)	
Gender: female	27 (90)	14 (88)	13 (93)	0.63
Age (years), median (IQR)	47 (38–55)	48 (39–55)	45 (36–56)	0.69
Country of birth				0.62
Australia or New Zealand	16 (53)	8 (50)	8 (57)	
UK	6 (20)	4 (25)	2 (14)	
Other	8 (27)	4 (25)	4 (29)	
Work classification				0.26
Full-time	17 (57)	11 (69)	6 (43)	
Part-time (0.6–0.9 EFT)	12 (40)	5 (31)	7 (50)	
Part-time (0.1–0.5 EFT)*	1 (3)	0 (0)	1 (7)	
Training classification – multiple responses possible				
Nursing	9 (30)	5 (31)	4 (29)	
Social work	9 (30)	5 (31)	4 (29)	
Occupational therapy	3 (10)	1 (6)	2 (14)	
Community services	7 (23)	4 (25)	3 (21)	
Disability services	2 (7)	1 (6)	1 (7)	
Aged care/care coordination	9 (10)	5 (31)	4 (29)	
Case management	14 (47)	7 (44)	7 (50)	
Other†	3 (10)		3 (21)	
Years worked as case manager				0.48
Less than 1 year	3 (10)	2 (13)	1 (7)	
1–5 years	14 (47)	9 (56)	5 (36)	
5.1–10 years	9 (30)	4 (25)	5 (36)	
>10 years	4 (13)	1 (6)	3 (21)	
Clients in case load, median (IQR)	25 (20–28)	26 (21–29)	23 (19–28)	0.66
Full-time CM	26 (20–37)	26 (20–36)	29 (23–42)	0.54
Part-time CM	23 (20–26)	26 (24–27)	20 (18–23)	0.52

*One CM only worked 2 days per week (0.4 EFT).

†Other: health promotion, counselling, master of law.

CM, case manager;

Zealand or UK and they had a median client case load of 25. One included Referral group case manager's hours changed after recruitment and training, and therefore did not meet inclusion criteria in that they did not work 3 or more days per week.

Client data for all 784 clients of the 30 case managers are included: 427 (54%) from the Facilitator, and 357 (46%) from the Referral groups. Clients had a median age of 83 years, most spoke English (89%) and most (73%) were on low level (one or two) packages (table 3). Client demographic data were similar in the two groups, apart from a slightly higher proportion of females in the referral group. Client demographics were representative of Australian HCP clients,²² where two-thirds are female, 90% are ≥70 years and three-quarters receive level one or two packages.

ACP activity

Figure 1 presents a flow chart depicting the baseline and 6-month intervention ACP activity for the entire group of clients.

ACP activity during the intervention period

Overall, case managers initiated ACP with 508 (65%) clients during the intervention period, with a higher proportion in Facilitator compared with the Referral groups (69% vs 60%, $p<0.05$). Of the remaining 276 clients, reasons for not initiating ACP were documented in 193 (70%) of files (figure 1).

Of those where ACP was initiated, 89 clients (11% of total 784 clients; 18% of the 508 clients where ACP was initiated) completed ACP conversations, with no significant difference between groups (figure 1, table 4). A further 167 clients (Facilitator 124, Referral 43; $p<0.005$) were interested in progressing ACP further, with 40 stating they would do this themselves without further assistance, and 127 clients noted to be progressing with but not completed ACP by the end of the study. The remaining 252 clients (Facilitator 116, Referral 136) either declined ACP discussion (173) or stated they had previously undertaken ACP (79) and did not wish further discussion (figure 1).

Table 3 Client demographics

	Overall n=784, n (%)	Facilitator n=427, n (%)	Referral n=357, n (%)	p=
Package level				0.26
1–2	572 (73)	321 (75)	251 (70)	
3–4	212 (27)	106 (25)	106 (30)	
Gender: female	512 (65)	264 (62)	248 (70)	0.03
Marital status				0.15
Married/long-term partner	291 (37)	175 (41)	116 (33)	
Widowed	380 (49)	192 (45)	188 (53)	
Divorced/separated/single	112 (14)	60 (14)	52 (15)	
Can communicate in English	694 (89)	384 (90)	310 (87)	
Age				0.22
Median (IQR)	83 (77–88)	84 (77–89)	83 (77–88)	
Medical conditions* – Multiple responses possible				0.18
Circulatory system	466 (59)	245 (57)	221 (62)	
Musculoskeletal and connective tissue	411 (52)	218 (51)	193 (54)	
Endocrine, nutritional and metabolic disorders	212 (27)	107 (25)	105 (29)	
Dementia	181 (23)	105 (25)	76 (21)	
Mental illness	160 (20)	88 (21)	72 (20)	

*Up to three medical conditions (coded according to the Aged Care Assessment Service classifications) per client, only conditions occurring for more 20% of clients are listed

Completed ACP conversations

Where initiated (n=508), ACP was completed in 89 (18%) clients, with 55 (66%) taking 60 min or less (table 4). Significantly more time was required when case managers facilitated conversations than when the clients were referred ($p<0.05$). However, the time taken at the ACP service was not recorded. Thus, total time required (HCP case manager and referral staff) for the Referral group clients is unknown. Clients in both groups required a median of 2 discussions.

Completed advance care directives

Of the 89 people completing ACP, 41 (46%) completed ACDs (table 4), with a higher proportion in the Referral (75%), compared with the Facilitator group (25%) ($p<0.005$).

Overall quality of documentation was poor. Only 12 (5 Facilitator, 7 Referral) of the 41 clients (29%) had all completed documents being valid (table 5). The substitute decision-maker documents were more likely to be correctly completed than either the RTC or the common law ACDs (60% vs 33% vs 29% respectively). The most common errors were absent, incomplete or inappropriate document signing or witnessing and inconsistencies within documents regarding treatments (eg, person indicating they did not want life-prolonging treatment but did want CPR performed, which is life prolonging).

Prior levels of ACP

Of the 784 clients, 322 clients (41%) had evidence in their records of prior ACP (table 6), with no difference between groups. The majority of these involved substitute decision-maker appointments. Only a minority

of these documents were present in the files (24 decision-maker appointments, 4 ACDs).

DISCUSSION

This is the first known study evaluating the effectiveness of models of ACP implementation in community aged care. During the 6-month intervention period, case managers initiated ACP with 65% of their clients. Although some differences between the two models were observed, that approximately two-thirds of clients in both groups achieved this result is a positive indicator for the implementation of both models, especially considering the minimal scope of the training interventions. However, a much smaller proportion of clients went on to complete ACP discussions and ACDs. Thus, it is important to consider why those where ACP was initiated did not complete it.

ACP may be considered a health behaviour where individuals have varied motivations, barriers, enablers and self-efficacy.^{23 24} ACP consists of multiple components including identifying values, and treatment goals, choosing substitute decision makers, discussions with family and health providers and completion of ACDs. Theories of behaviour change²⁵ recognise the importance of a stepped approach (precontemplation, contemplation, preparation, action and maintenance) towards target behaviours. In this study, the principal outcomes were the completed ACP discussions and ACDs. Interestingly, of clients where ACP was initiated but incomplete, one-third reported they were in progress. Similar observations over the same time period have been made elsewhere.²⁶ It is reasonable to suppose that over time ACP completion rates might

Figure legend:

Fac = client of case manager in "Facilitator" arm of study

Ref = client of case manager in "Referral" arm of study

ACD = Advance care directive (Includes statutory and common law ACDs)

MEPOA= Medical enduring power of attorney (Substitute decision maker)

Figure 1 Study flow chart. ACP, advance care planning.

be higher. Also, similar to previous studies,^{2 20} almost half of those who completed ACP discussions also completed ACDs.

The HCP population is heterogeneous. It is therefore reasonable to expect clients to be at differing stages of readiness for each of the ACP components. Despite being mostly an elderly, frail population, as other studies have postulated,^{23 27 28} many of this group and their families may not have viewed ACP as relevant. They may not have considered themselves sick enough, may have lacked information regarding their illness and prognosis, may not have wished to discuss their likely future worsening health state or death, may have been occupied with other priorities or assigned a low priority to ACP without clear triggers or health crises. Previous research has shown that illness severity is not necessarily associated with ACD completion.²⁴ Other elements of ACP, such as

values clarification and communication with family/carer(s), may have occurred but were not measured here. The findings in this study therefore may not be surprising. Supporting case managers to use tools to assist with prognosis, such as the Supportive and Palliative Care Indicator Tool²⁹ or the 'surprise question'³⁰ may help prioritise clients for ACP. Furthermore, incorporating ACP into routine assessments, such as on commencement of an aged-care package, following hospital admission and during specific assessments (health assessment for people aged 75 and older, chronic disease assessment), may help to overcome the perceived lack of a trigger or health crisis.

ACP activity prior to the intervention in this study found that a third of the group reported having already appointed substitute decision maker(s), and 5% had completed ACDs. These reported levels of existing

Research

Table 4 Outcomes of advance care planning (ACP), including documentation

		Overall n (%)	Facilitator n (%)	Referral n (%)	P=*
Total clients in group		784	427	357	
Clients with whom ACP was initiated		508 (65)	293 (69)	215 (60)	0.01
Clients completing ACP (% clients where ACP was initiated)		89 (18%)	53 (18%)	36 (17%)	0.31
Time taken	≤30 min	31	15	16	0.02
	31–60 min	24	9	15	
	61–90 min	7	5	2	
	90–120 min	6	6	0	
	>120 min	15	13	2	
	Missing data	6	5	1	
All documents n (% who completed ACP)		41 (46)	13 (25)	28 (78)	<0.005
Medical enduring power of attorney	Total	15	5	10	0.30
	Alone	3	2	1	
	Plus CL-ACD	9	2	6	
	Plus CL-ACD and RTC	4	1	3	
Common law advance care directives (CL-ACD)	Total	38	11	27	0.10
	Alone	23	6	17	
	Plus MEPOA	9	3	6	
	Plus MEPOA and RTC	4	1	1	
	Plus RTC	2	1	1	

*For facilitator compared with referral.

MEPOA, medical enduring power of attorney (substitute decision maker); RTC, Refusal of Treatment Certificate.

appointments of substitute decision makers are higher than other studies have found.^{27 31} The majority of these pre-existing documents were absent from the client's file, and therefore it is unclear as to their existence and

validity. Anecdotally, our experience has found people wrongly believe that they have already completed them, confusing ACDs with wills and financial powers of attorney.

Table 5 Validity of completed advance care directives

	Overall n (%)	Facilitator n (%)	Referral n (%)
Medical enduring power of attorney	15	5	10
Valid as per legal requirements n (%)	9 (60)	3 (60)	6 (60)
Validity errors	6 (40)	2 (40)	4 (40)
Not signed by client	1	—	1
Witnessing of document incomplete, incorrect*	5	2	2
Refusal of Treatment Certificate	6	2	4
Valid as per legal requirements	2 (33)	0	2 (50)
Validity errors	4 (67)	2 (100)	2 (50)
Treatment refused option not indicated	2	1	1
'Medical treatment generally' chosen but specific treatment specified	1	1	—
Witnessing of document incomplete†	1	—	1
Advance care directive (common law)	38	11	27
Complies with best practice	11(29)	4 (36)	7 (26)
Best practice non-compliance element (one or more)			
Incompatible CPR/LPT option‡	14	5	9
Not signed by client	14	2	12
Not signed by witness	20	4	16
Not signed by doctor	17	2	15
Number with both treatment inconsistencies and signing/witness issues	11	3	8

*Must be signed in the presence of two witnesses. Neither witness can be the agent/alternative agent. One witness must be authorised by law to witness the signing of statutory declarations.

†Must be signed in the presence of two witnesses, one must be a registered medical practitioner.

‡Person has selected that they do not want LPT, but want CPR, or have stated they want CPR, but have also stated they do not want to be resuscitated. CPR, cardiopulmonary resuscitation; LPT, life-prolonging treatment.

Table 6 Pre-existing advance care planning (ACP)

Type of pre-existing ACP	Type of ACD (where applicable)	Overall n (%)	Facilitator n (%)	Referral n (%)	p=*
Appointment of substitute decision maker	MEPOA only	229 (71.1)	129 (70.9)	100 (71.4)	0.91
ACD outlining preferences only	Common law ACD	11 (3.4)	5 (2.7)	6 (4.3)	0.45
	RTC	2 (0.6)	0 (0.0)	2 (1.4)	0.11
MEPOA and ACD	Common law ACD	22 (6.8)	10 (5.5)	12 (8.6)	0.28
	RTC	2 (0.6)	0 (0.0)	2 (1.4)	0.11
	Common law ACD and RTC	5 (1.6)	2 (1.1)	3 (2.1)	0.45
ACP discussion, no documents		51 (15.8)	36 (19.8)	15 (10.7)	0.22
Total clients with pre-existing ACP		322	182	140	0.27

*For facilitator compared with referral.

ACD, advance care directive; MEPOA, medical enduring power of attorney (substitute decision maker); RTC, Refusal of Treatment Certificate.

When ACDs were completed, they were generally of poor quality, with less than half being legally valid. The reasons for this are unclear and raise issues of how quality control of ACP and ACD should be managed and also raise concern that the ACD documents themselves are overly complicated or difficult to complete. Both groups experienced issues regarding signing and witnessing of documents, something that may be improved by embedding ACP into the broader health-care system and including the clients' usual doctors in the process. Neither model in this study addressed this issue specifically.

While ACP completion rates were similar in both groups, more Facilitator clients had ACP initiated (Facilitator 69%, Referral 60%; $p<0.05$) and more (Facilitator 124, Referral 43; $p<0.005$) reported that ACP was in progress at the end of the study. This finding raises the possibility that with longer follow-up, there might be differences in favour of the Facilitator group. This may be a reflection of the 'Facilitator' group receiving more extensive ACP training and, therefore, feeling more comfortable/skilled in ACP. Despite small numbers overall, there was a significantly higher percentage of document completion in the Referral group. Thus, while both models of ACP have potential advantages, this study has shown similar outcomes overall.

The client population and case manager characteristics were representative of Australian HCP services, and therefore results are likely to be generalisable to the Australian HCP population. However, this intervention was only of 6 months' duration, and the outcomes measured were completion of ACP discussions and ACDs. While these measurements are consistent with existing ACP literature, recently there is ongoing interest as to how to best capture the efficacy of ACP interventions and indeed which ACP components are most critical to its success. This may be particularly important given the transtheoretical model of behavioural change theory and the relevance to ACP. Other steps in the ACP process were not measured in this study. Furthermore, this study did not look at whether ACP affected future decision making and

end-of-life care—the ultimate aims of ACP. Although the HCP organisations received payment and case managers committed to the requirements of the study, it is unknown whether any additional time or funding was made available to case managers to conduct ACP discussions. Finally, this study did not actively source pre-existing ACDs, thus it is not possible to compare the quantity and quality of pre-existing documentation with those achieved in this study.

CONCLUSION

This study found that during a 6-month period, both Facilitator and Referral models of ACP provision achieved similar results, in terms of ACP initiation, completed conversations and ACD documentation. While ACP was initiated with many clients, there was low numbers of completed discussions and documents, and documentation quality was generally poor. Both models have merit, and how providers offer ACP services may depend on factors, such as access to a suitable referral service, access to training and support for case managers and preferences of the clients. The findings raise questions for future research into ACP practice in the community aged care, including when and how to offer and conduct ACP, what training is required to provide the necessary skills and what are the most useful outcome measures for determining whether an ACP intervention has been successful.

Acknowledgements Previous staff: William Silvester who was involved in the design of the study and Catherine O'Leary who was involved in training and support.

Contributors KMD, RZC and EAS: conception, design, interpretation and drafting of manuscript; MWS and VL: conception, design, interpretation and review of manuscript; and MWS and RZC: participant training and support and data collection. KMD is the guarantor.

Funding This work was supported by programme funding received from the Australian Government Department of Health.

Competing interests None declared.

Ethics approval Austin Health's institutional research ethics committee (Ref LNR/14/Austin/309).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 The Australian Commission of Quality and Safety in Health Care. The National Consensus Statement: Essential elements for safe and high-quality end-of-life care. 2015 <http://www.safetyandquality.gov.au/publications/national-consensus-statement-essential-elements-for-safe-high-quality-end-of-life-care/> (accessed 25 March 2017)
- 2 Detering KM, Hancock AD, Reade MC, *et al.* The impact of advance care planning on end of life care in elderly patients: randomised controlled trial. *BMJ* 2010;340:c1345.
- 3 Brinkman-Stoppelenburg A, Rietjens JA, van der Heide A. The effects of advance care planning on end-of-life care: a systematic review. *Palliat Med* 2014;28:1000–25.
- 4 Wright AA, Zhang B, Ray A, *et al.* Associations between end-of-life discussions, patient mental health, medical care near death, and caregiver bereavement adjustment. *JAMA* 2008;300:1665–73.
- 5 Australian Government Department of Social Services. The home care packages program guidelines. https://www.dss.gov.au/sites/default/files/documents/09_2015/hcp_operational_manual_finalseptember2015_3.pdf (accessed 25 March 2017).
- 6 Auret K, Sinclair C, Thomas J, *et al.* When paperwork does not represent the patient's wishes for surrogate decision-making. *Intern Med J* 2015;45:1200.
- 7 Auret K, Sinclair C, Averill B, *et al.* Advance care planning and end-of-life care in a network of rural Western Australian hospitals. *Aust J Rural Health* 2015;23:195–200.
- 8 Australian Institute of Health and Welfare. Residential aged care in Australia 2009–10: a statistical overview. <http://www.aihw.gov.au/WorkArea/DownloadAsset.aspx?id=10737421028> (accessed 25 Mar 2017)
- 9 Ratner E, Norlander L, McSteen K. Death at home following a targeted advance-care planning process at home: the kitchen table discussion. *J Am Geriatr Soc* 2001;49:778–81.
- 10 Ditto PH, Jacobson JA, Smucker WD, *et al.* Context changes choices: a prospective study of the effects of hospitalization on life-sustaining treatment preferences. *Med Decis Making* 2006;26:313–22.
- 11 Baughman KR, Aultman J, Hazelett S, *et al.* Managing in the trenches of consumer care: the challenges of understanding and initiating the advance care planning process. *J Gerontol Soc Work* 2012;55:721–37.
- 12 Black K. Correlates of case managers' advance care planning practices. *Clin Gerontol* 2010;33:124–35.
- 13 Golden AG, Tewary S, Qadri S, *et al.* The positive attitudes and perceptions of care managers about advance directives. *Am J Hosp Palliat Care* 2011;28:98–101.
- 14 Molloy DW, Russo R, Pedlar D, *et al.* Implementation of advance directives among community-dwelling veterans. *Gerontologist* 2000;40:213–7.
- 15 Baughman KR, Ludwick RE, Merolla DM, *et al.* Professional judgments about advance care planning with community-dwelling consumers. *J Pain Symptom Manage* 2012;43:10–19.
- 16 Black K, Fauske J. Measuring case managers' advance care planning practice: translating focus group findings to survey development. *Care Manag J* 2008;9:166–76.
- 17 Sellars M, Detering KM, Silvester W. Current advance care planning practice in the Australian community: an online survey of home care package case managers and service managers. *BMC Palliat Care* 2015;14:15.
- 18 Silvester W, Fullam RS, Parslow RA, *et al.* Quality of advance care planning policy and practice in residential aged care facilities in Australia. *BMJ Support Palliat Care* 2013;3:349–57.
- 19 Advance care planning Australia. <http://www.advancereplanning.org.au> <http://advancereplanning.org.au/advance-care-planning> (accessed 25 Mar 2017).
- 20 Detering K, Sutton E, Fraser S, *et al.* Feasibility and acceptability of advance care planning in elderly Italian and Greek speaking patients as compared to English-speaking patients: an Australian cross-sectional study. *BMJ Open* 2015;5:e008800.
- 21 Simpson-Young V, Fine M. Centre for research on social inclusion, macquarie university. case management in community aged care: mapping the landscape in NSW. <http://www.crsi.mq.edu.au/public/download.jsp?id=88> (accessed 27 Mar 2017).
- 22 Australian Institute of Health and Welfare. Residential aged care and aged care packages in the community 2012–13. 2014 <http://www.aihw.gov.au/aged-care/residential-and-community-2012-13/> (accessed 27 Mar 2017).
- 23 Fried TR, Redding CA, Robbins ML, *et al.* Stages of change for the component behaviors of advance care planning. *J Am Geriatr Soc* 2010;58:2329–36.
- 24 Fried TR, Redding CA, Robbins ML, *et al.* Promoting advance care planning as health behavior change: development of scales to assess decisional balance, medical and religious beliefs, and processes of change. *Patient Educ Couns* 2012;86:25–32.
- 25 Prochaska JO, Velicer WF. The transtheoretical model of health behavior change. *Am J Health Promot* 1997;12:38–48.
- 26 Rocque GB, Dionne-Odom JN, Sylvia Huang CH, *et al.* Implementation and impact of patient lay navigator-led advance care planning conversations. *J Pain Symptom Manage* 2017;53:682–92.
- 27 Sinclair C, Auret KA, Burgess A. The balancing point: understanding uptake of advance directive forms in a rural Australian community. *BMJ Support Palliat Care* 2013;3:358–65.
- 28 Simon J, Porterfield P, Bouchal SR, *et al.* 'Not yet' and 'Just ask': barriers and facilitators to advance care planning—a qualitative descriptive study of the perspectives of seriously ill, older patients and their families. *BMJ Support Palliat Care* 2015;5:54–62.
- 29 Supportive & Palliative Care Indicators Tool (SPICT™) (accessed 16th July 2017) <http://www.spict.org.uk>
- 30 Lynn J. Living long in fragile health: the new demographics shape end of life care. *Hastings Cent Rep* 2005;Spec No:s14–s18.
- 31 White B, Tilse C, Wilson J, *et al.* Prevalence and predictors of advance directives in Australia. *Intern Med J* 2014;44:975–80.