

Full Reference List of 184 Papers

Abarshi E, Echteld MA, Van den Block L, et al. Use of palliative care services and general practitioner visits at the end of life in The Netherlands and Belgium. *J Pain Symptom Manage*. 2011;41(2):436-48.

Collins K, Harding R. Improving HIV management in sub-Saharan Africa: how much palliative care is needed? *AIDS Care* 2007;19(10):1304-1306.

Albers G, Harding R, Pasma HR, et al. What is the methodological rigour of palliative care research in long-term care facilities in Europe? A systematic review. *Palliat Med* 2012;26(5):722-733.

Alexander CS, Memiah P, Henley YB, et al. Palliative care and support for persons with HIV/AIDS in 7 African countries: implementation experience and future priorities. *Am J Hosp Palliat Care* 2012;29(4):279-285.

Amery J. Mapping children's palliative care around the world: an online survey of children's palliative care services and professionals' educational needs. *J Palliat Med* 2012;15(6):646-652.

Amery JM, Rose CJ, Holmes, J, Nguyen J, Byarugaba C. The Beginnings of Children's Palliative Care in Africa: Evaluation of a Children's Palliative Care Service in Africa 2009;12(11):1015-1021.

Ando M, Morita T, Ahn SH, Marquez-Wong F, Ide S. International comparison study on the primary concerns of terminally ill cancer patients in short-term life review interviews among Japanese, Koreans, and Americans. *Palliat Support Care* 2009;7(3):349-355.

Anquinet L, Rietjens JAC, Seale C, et al. The Practice of Continuous Deep Sedation Until Death in Flanders (Belgium), The Netherlands, and the U.K.: A Comparative Study. *J Pain Symptom Manage*. 2012;44(1):33-43.

Bainbridge D, Brazil K, Krueger P, Ploeg J, Taniguchi A. A proposed systems approach to the evaluation of integrated palliative care. *BMC Palliat Care* 2010;9:8.

Bansal M, Patel FD, Mohanti BK, Sharma SC. Setting up a palliative care clinic within a radiotherapy department: a model for developing countries. *Support Care Cancer* 2003;11(6):343-347.

Barazzetti G, Borreani C, Miccinesi G, Toscani F. What "best practice" could be in Palliative Care: an analysis of statements on practice and ethics expressed by the main Health Organizations. *BMC Palliat Care*. 2010;9:1.

Bausewein C, Calanzani N, Daveson BA, et al. Burden to others' as a public concern in advanced cancer: a comparative survey in seven European countries. *BMC Cancer*. 2013;8:13.

Bentur N, Resnizky. Validation of the McGill Quality of Life Questionnaire in home hospice settings in Israel. *Palliat Med* 2005;19(7):538-544.

Bilsen J, Norup M, Deliens L, et al. Drugs used to alleviate symptoms with life shortening as a possible side effect: End-of-life care in six European countries. *J Pain Symptom Manage* 2006;31(2):111-121.

- Bingley A, Clark D. A Comparative Review of Palliative Care Development in Six Countries Represented by the Middle East Cancer Consortium (MECC). *J Pain Symptom Manage* 2009;37(3):287-296.
- Bishop K, Mele N, Koppmann MJ, Day S. A review of paediatric palliative care nursing education in Latin America. *Int J Palliat Nurs* 2009;15(8):377-382.
- Bito S, Matsumura S, Singer MK, et al. Acculturation and end-of-life decision making: comparison of Japanese and Japanese-American focus groups. *Bioethics* 2007;21(5):251-262.
- Bollini P, Venkateswaran C, Sureshkumar K. Palliative care in Kerala, India: A model for resource-poor settings. *Onkologie* 2004;27(2):138-142.
- Borasio GD, Shaw PJ, Hardiman O, et al. Standards of palliative care for patients with amyotrophic lateral sclerosis: results of a European survey. *Amyotroph Lateral Scler Other Motor Neuron Disord* 2001;2(3):159-164.
- Broad JB, Gott M, Kim H, et al. Where do people die? An international comparison of the percentage of deaths occurring in hospital and residential aged care settings in 45 populations, using published and available statistics. *Int J Public Health* 2013;58(2):257-67.
- Bruera E, Neuman CM, Mazzocato C, Stiefel F, Sala R. Attitudes and beliefs of palliative care physicians regarding communication with terminally ill cancer patients. *Palliat Med* 2000;14(4):287-298.
- Buiting HM, van Delden JJ, Rietjens JA, et al. Forgoing artificial nutrition or hydration in patients nearing death in six European countries. *J Pain Symptom Manage* 2007;34(3):305-314.
- Cameron BL, Salas AS, de Moissac D. Participatory knowledge exchange to support palliative care in Chile: Lessons learned through global health research. *Can J Nurs Res* 2011;43(3):16-37.
- Caraceni A, Martini C, Zecca E, et al. Breakthrough pain characteristics and syndromes in patients with cancer pain. An international survey. *Palliat Med* (2004);18(3):177-183.
- Cartwright C, Onwuteaka-Phillipsen BD, Williams G, et al. Physician discussions with terminally ill patients: a cross-national comparison. *Palliat Med* 2007;21(4):295-303.
- Miccinesi G, Fischer S, Paci E, et al. Physicians' attitudes towards end-of-life decisions: a comparison between seven countries. *Soc Sci Med* 2005;60(9):1961-1974.
- Carvajal A, Hribernik N, Duarte E, Sanz-Rubiales A, Centeno C. The Spanish Version of the Edmonton Symptom Assessment System-Revised (ESAS-r): First Psychometric Analysis Involving Patients With Advanced Cancer. *J Pain Symptom Manage* 2013;45(1):129-136.
- Centeno C, Noquera A, Lynch T, Clark D. Official certification of doctors working in Palliative Medicine in Europe: data from an EAPC study in 52 European countries. *Palliat Med* 2007;21(8):683-687.
- Centeno C, Clark D, Lynch T, et al. Facts and indicators on palliative care development in 52 countries of the WHO European region: results of an EAPC task force. *Palliat Med* 2007;21(6):463-471.

Charlton R, Dovey S, Mizushima Y, Ford E. Attitudes to death and dying in the UK, New Zealand and Japan. *J Palliat Care* 1995;1(1):42-47.

Chatuverdi SK. Ethical dilemmas in palliative care in traditional developing societies, with special reference to the Indian setting. *J Med Ethics* 2008;34(8):611-615.

Cherny NI, Catane R, European society of Medical Oncology Taskforce on Palliative and Supportive Care. Attitudes of medical oncologists toward palliative care for patients with advanced and incurable cancer - Report on a survey by the European Society of Medical Oncology Taskforce on Palliative and Supportive Care. *Cancer* 2003;98(11):2502-2510.

Cherny NI, Baselga J, de Conno F, Radbruch L. Formulary availability and regulatory barriers to accessibility of opioids for cancer pain in Europe: a report from the ESMO/EAPC Opioid Policy Initiative. *Ann Oncol* 2010;21(3):615-626.

Clark D, Wright M, Hunt J, Lynch T. Hospice and palliative care development in Africa: a multi-method review of services and experiences, *J Pain Symptom Manage* 2007;33(6):698-710.

Coenen A, Doorenbos AZ, Wilson SA. Nursing interventions to promote dignified dying in four countries. *Oncol Nurs Forum* 2007;34(6):1151-1156.

Cohen J, Bilsen J, Fischer S, et al. End-of-life decision-making in Belgium, Denmark, Sweden and Switzerland : does place of death make a difference? *J Epidemiol Community Health* 2007;61(12):1062-1068.

Cohen J, Bilsen J, Miccinesi G, et al. Using death certificate data to study place of death in 9 European countries: opportunities and weaknesses. *BMC Public Health* 2007;8:283.

Cohen J, Bilsen J, Addington-Hall, J, et al. Population-based study of dying in hospital in six European countries. *Palliat Med* 2008;22(6):702-710.

Cohen J, van Delden J, Mortier F, et al. Influence of physicians' life stances on attitudes to end-of-life decisions and actual end-of-life decision-making in six countries. *J Med Ethics* 2008;34(4):247-253.

Cohen LW, van der Sheen JT, Reed D, et al. Family perceptions of end-of-life care for long-term care residents with dementia: Differences between the United States and the Netherlands. *J Am Geriatr Soc* 2012;60(2):316-322.

Constantini M, Di Leo S, Beccaro M. Methodological issues in a before-after study design to evaluate the Liverpool Care Pathway for the Dying Patient in hospital. *Palliat Med* 2011;25(8):766-773.

Constantini M, Ottonelli S, Canavacci L, et al. The effectiveness of the Liverpool care pathway in improving end of life care for dying cancer patients in hospital. A cluster randomised trial. *BMC Health Serv Res* 2011;11:13.

Csikos A, Mastrojohn J3rd, Albanese T, et al. Physicians' Beliefs and Attitudes About End-of-Life Care: A Comparison of Selected Regions in Hungary and the United States. *J Pain Symptom Manage* 2010;39(1):76-87.

Curiale V. World-wide literature survey on geriatric palliative medicine. *EGM* 2(1):6-11.

De Cinque N, Monterosso L, Dadd G, Sidhu R, Lucas R. Bereavement support for families following the death of a child from cancer: practice characteristics of Australian and New Zealand paediatric oncology units. *J Paediatr Child Health* 2004;40(3):131-135.

Daveson BA, Bausewein C, Murtagh FE, et al. To be involved or not to be involved: A survey of public preferences for self-involvement in decision-making involving mental capacity (competency) within Europe. *Palliat Med* 2013;27(5):418-427.

Daveson BA, Bechinger-English D, Bausewein C, et al. Constructing understandings of end-of-life-care in Europe: A qualitative study involving cognitive interviewing with implications for cross-national surveys. *J Palliat Med* 2011;14(3):343-349.

Daveson BA, Simon ST, Benalia H, et al. Are we heading in the same direction? European and African doctors' and nurses' views and experiences regarding outcome measurement in palliative care. *Palliat Med* 2012;26(3):242-249.

Davies AN, Dickman A, Reid C, et al. The management of cancer-related breakthrough pain: Recommendations of a task group of the Science Committee of the Association for Palliative Medicine of Great Britain and Ireland. *Eur J Pain* 2009;13(4):331-338.

De Lima L. Key concepts in palliative care: the IAHPHC list of essential medicines in palliative care. *Eur J Hosp Pharm* 2012;19(1):34-37.

De Lima L, Krakauer EL, Lorenz K, et al. Ensuring palliative medicine availability: the development of the IAHPHC list of essential medicines for palliative care. *J Pain Symptom Manage* 2007;33(5):521-526.

De Lima, Bennett MI, Murray SA, et al. International Association for Hospice and Palliative Care (IAHPHC) List of Essential Practices in Palliative Care. *J Pain Palliat Care Pharmacother* 2012;26(2):118-122.

De Lima L, Sakowski JA, Stratton Hill C, Bruera E. Legislation analysis according to WHO and INCB criteria on opioid availability: a comparative study of 5 countries and the state of Texas. *Health Policy* 2001;56(2):99-110.

Decker SL, Higginson IJ. A tale of two cities: Factors affecting place of cancer death in London and New York. *Eur J Public Health* 2007;17(3):285-290.

Delgado E, Barfield RC, Baker JN, et al. Availability of palliative care services for children with cancer in economically diverse regions of the world. *Eur J Cancer* 2010;46(12):2260-2266.

Demmer E, Burghart G. Experiences of AIDS-related bereavement in the USA and South Africa: a comparative study. *Int Soc Work* 2008;51(3):360-370.

Di Leo S, Beccaro M, Finelli S, Borreani C, Constantini M. Expectations about and impact of the Liverpool Care Pathway for the dying patient in an Italian hospital. *Palliat Med* 2011;35(4):293-303.

Doorembos AZ, Abaquin C, Perrin ME, et al. Supporting dignified dying in the Philippines. *Int J Palliat Nurs* 2011;17(3):125-130.

- Doorenbos AZ, Wilson SA, Coenen A. A cross-cultural analysis of dignified dying. *J Nurs Scholarsh* 2006;38(4):352-357.
- Downing J, Jack BA. End-of-life care in rural areas: what is different? *Curr Opin Support Palliat Care* 2012;6(3):391-397.
- Downing J, Simon ST, Mwangi-Powell FN, et al. Outcomes 'Out of Africa': the selection and implementation of outcome measures for palliative care in Africa. *BMC Palliat Care* 2012;11:1.
- Cohen LM, Germain MJ, Poppel DM, et al. Dying well after discontinuing the life-support treatment of dialysis. *Arch Intern Med* 2000;160(16):2513-8.
- Harding R, Easterbrook P, Higginson IJ, et al. Access and equity in HIV/AIDS palliative care: a review of the evidence and responses. *Palliat Med* 2005;19(3):251-258.
- Ens CD, Chochinov HM, Bérard JL, et al. Pediatric palliative care online: the views of the health care professionals. *J Palliat Care* 2008;24(1):41-48.
- Ersek M, Sebego M, Bloom AM, et al. Development and evaluation of an international, interdisciplinary palliative care workshop in Botswana. *Int J Palliat Nurs* 2010;16(1):13-21.
- Evans N, Pasman HR, Vega Alonso T, et al. End-of-life decisions: A cross-national study of treatment preference discussions and surrogate decision-maker appointments. *PloS One* 2013; 8(3).
- Farnon C. A personal exploration of the German hospice system. *Am J Hosp Palliat Care* 1996;13(4):32-37.
- Fegg MJ, Kramer M, L'hoste S, Borasio GD. The Schedule for Meaning in Life Evaluation (SMiLE): Validation of a new instrument for meaning-in-life research. *J Pain Symptom Manage* 2008;35(4):356-364.
- Gerzevitz D, ferszt G, Vosit-Steller J, Mitrea N. Cross-cultural consultation on palliative care: the use of teleconferencing. *JHPN* 2009;11(4):239-244.
- Gilson AM, Maurer MA, Lebaron VT, Ryan KM, Cleary JF. Multivariate analysis of countries' government and health-care system influences on opioid availability for cancer pain relief and palliative care: More than a function of human development. *Palliat Med* 2013;27(2):105-114.
- Gomes B, Higginson IJ, Calanzani N, et al. Preferences for place of death if faced with advanced cancer: a population survey in England, Flanders, Germany, Italy, the Netherlands, Portugal and Spain. *Ann Oncol* 2012 (23(8):2006-2015.
- Gomez-Batiste X, Caja C, Espinosa J, et al. The Catalonia World Health Organization Demonstration Project for Palliative Care Implementation: Quantitative and Qualitative Results at 20 Years. *J Palliat Care* 2013;27(4):783-794.
- Gomez-Batiste X, Fontanals MD, Roca J, et al. Catalonia WHO demonstration project on palliative care implementation 1990-1995: Results in 1995. *J Pain Symptom Manage* 1996;12(2):73-78.

Gomez-Batiste X, Buisan M, Gonzalez MP, et al. The "La Caixa" Foundation and WHO Collaborating Center Spanish National Program for enhancing psychosocial and spiritual palliative care for patients with advanced diseases, and their families: Preliminary findings. *Palliat Support Care* 2011;9(3):239-239.

Gott M, Seymour J, Ingleton C, Gardiner C, Bellamy G. That's part of everybody's job': The perspectives of health care staff in England and New Zealand on the meaning and remit of palliative care. *Palliat Med* 2012;26(3):232-241.

Grond S, Zech D, Schug SA, Lynch J, Lehmann KA. Validation of World Health Organization guidelines for cancer pain relief during the last days and hours of life. *J Pain Symptom Manage* 1991;6(7):411-422.

Gurriere DN, Coyte PC. The Ambulatory and Home Care Record: A Methodological Framework for Economic Analyses in End-of-Life Care. *J Aging Res* 2011;374237-374237.

Gysels M, Pell C, Straus L, Pool R. End of life care in sub-Saharan Africa: a systematic review of the qualitative literature. *BMC Palliat Care* 2011;10.

Gysels M, Evans N, Menaca A, et al. Culture and End of Life Care: A Scoping Exercise in Seven European Countries. *PLS One* 2012;7(4).

Gysels M, Evans N, Menaca A, et al. Culture is a priority for research in end-of-life care in Europe: A research agenda. *J Pain Symptom Manage* 2012 44(2):285-294.

Hack, TF, McClement SE, Chochinov HM, et al. Learning from dying patients during their final days: life reflections gleaned from dignity therapy. *Palliat Med* 2010;24(7):715-723.

Hagen NA, Stiles C, Nikolaichuck C, et al. The Alberta Breakthrough Pain Assessment Tool for cancer patients: A validation study using a Delphi process and patient think-aloud interviews. *J Pain Symptom Manage* 2008;35(2):136-152.

Harding R, Higginson IJ, PRISMA. PRISMA: a pan-European co-ordinating action to advance the science in end-of-life cancer care. *Eur J Cancer* 2010;46(9):1493-1501.

Harding R, Stewart K, Marconi K, O'Neill JF, Higginson IJ. Current HIV/AIDS end-of-life care in sub-Saharan Africa: a survey of models, services, challenges and priorities. *BMC Public Health* 2003;3.

Harding R, Selman L, Aquipio G, et al. Prevalence, burden, and correlates of physical and psychological symptoms among HIV palliative care patients in sub-Saharan Africa: An international multicenter study. *J Pain Symptom Manage* 2012;44(1):1-9.

Harding R, Simon ST, Benalia H, et al. The PRISMA Symposium 1: Outcome Tool Use. Disharmony in European Outcomes Research for Palliative and Advanced Disease Care: Too Many Tools in Practice. *J Pain Symptom Manage* 2011 42(4):493-500.

Herrera E, Rocafort J, De Lima L, et al. Regional Palliative Care Program in Extremadura: An effective public health care model in a sparsely populated region. *J Pain Symptom Manage* 2007;33(5):591-598.

Herzler M, Franze T, Dietze F, Asadullah K. Dealing with the issue 'care of the dying' in medical education - results of a survey of 592 European physicians. *Med Educ* 2000;34(2):146-147.

Higginson IJ, Simon ST, Benalia H, et al. Republished: Which questions of two commonly used multidimensional palliative care patient reported outcome measures are most useful? Results from the European and African PRISMA survey. *Postgrad Med J* 2012;88(1042):451-457.

Hjermstad MJ, Gibbins J, Haugen DF. Pain assessment tools in palliative care: An urgent need for consensus. *Palliat Med* 2008;22(8):895-903.

Houttekier D, Cohen J, Bilsen J. Place of death in metropolitan regions: Metropolitan versus non-metropolitan variation in place of death in Belgium, The Netherlands and England. *Health Place* 2010;16(1):132-139.

Houttekier D, Cohen J, Bilsen J, et al. Place of death of older persons with dementia. A study in five European countries. *J Am Geriatr Soc* 2010;58(4):751-756.

Hudson PL, Zordan R, Trauer T. Research Priorities Associated with Family Caregivers in Palliative Care: International Perspectives. *J Palliat Med* 2011;14(4):397-401.

Iranmanesh S, Ghazanfari Z, Sävenstedt S, Häggström T. "Professional Development: Iranian and Swedish nurses' experiences of caring for dying people. *J Palliat Care* 202-209.

Jocham HR, Dassen T, Widdershoven G, Halfens R. Reliability and validity of the EORTC QLQ-C30 in palliative care cancer patients. *Cent Eur J Med* 2009;4(3):348-357.

Johnston DL, Nagel K, Friedman DL, et al. Availability and use of palliative care and end-of-life services for pediatric oncology patients. *J Clin Oncol* 2008;26(28):4646-4650.

Jünger S, Payne S, Brearley S, Ploenes V, Radbruch L. Consensus building in palliative care: a Europe-wide Delphi study on common understandings and conceptual differences. *J Pain Symptom Manage* 2012;44(2):192-205.

Kaasa S, Torvik K, Cherny K, Hanks G, de Conno F. Patient demographics and centre description in European palliative care units. A cross sectional survey of the European Association for Palliative Care (EAPC) Research Network. *Palliat Med* 2007;21(1):15-22.

Jo KH, Doorenbos AZ, Sung KW, et al. Nursing interventions to promote dignified dying in South Korea. *Int J Palliat Nurs* 2011;17(8):392-397.

Kim KU, Yoon SJ, Lee JL, et al. Validation of the Korean Version of the McMaster Quality of Life Scale in terminal cancer patients. *J Palliat Care* 2006;22(1):40-45.

Klinger CA, Connor SR. National associations survey: Advancing hospice and palliative care worldwide. *J Pain Symptom Manage* 2005;30(4):386-388.

Ko W, Beccaro M, Miccinesi G, et al. Awareness of General Practitioners concerning cancer patients' preferences for place of death: Evidence from four European countries. *Eur J Cancer* 2013;49(8):1967-1974.

- Krakauer EL, Wenk R, Buitrago R, Jenkins P, Scholten W. Opioid inaccessibility and its human consequences: reports from the field. *J Pain Palliat Pharmacother* 2010;24(3):239-243.
- Kyriaki M, Eleni T, Efi P, et al. The EORTC core quality of life questionnaire (QLQ-C30, version 3.0) in terminally ill cancer patients under palliative care: Validity and reliability in a Hellenic sample. *Int J Cancer* 2001;94(1):135-139.
- Larkin PJ, Dierckx de Casterlé B, Schotsmans P. Transition towards end of life in palliative care: an exploration of its meaning for advanced cancer patients in Europe. *J Palliat Care* 2007;23(2):69-79.
- Latour JM, Fulbrook P, Albarran JW. EfCCNa survey: European intensive care nurses' attitudes and beliefs towards end-of-life care. *Nurs Crit Care* 2009;14(3):110-121.
- Lee CY, Komatsu H, Zhang W, et al. Comparison of the Hospice Systems in the United States, Japan and Taiwan. *Asian Nurs Res (Korean Soc Nurs Sci)* 2010;4(4): 163-173.
- Leppert W, Majkowicz M, Ahmedzai SH. The adaptation of the Sheffield Profile for Assessment and Referral for Care (SPARC) to the Polish clinical setting for needs assessment of advanced cancer patients. *J Pain Symptom Manage* 2012;44(6):916-922.
- Lewis JM, Di Giacomo M, Currow D, Davidson PM. Dying in the margins: understanding palliative care and socioeconomic deprivation in the developed world. *J Pain Symptom Manage* 2011;42(1):105-118.
- Lin CC. Applying the American Pain Society's QA standards to evaluate the quality of pain management among surgical, oncology, and hospice inpatients in Taiwan. *Pain* 2000;87(1):43-49.
- Lindqvist O, Tishelman C, Hagelin CL, et al. Complexity in non-pharmacological caregiving activities at the end of life: an international qualitative study. *PLoS Med* 2012;9(2): e1001173-e1001173.
- Lindqvist O, Lundquist G, Dickman A, et al. Four Essential Drugs Needed for Quality Care of the Dying: A Delphi-Study Based International Expert Consensus Opinion. *J Palliat Med* 2012;16(1):38-43.
- Lloyd-Williams M, Dogra N, Morake R. A comparison of attitudes of medical students in England and in South Africa towards patients with life-limiting illness. *J Palliat Care* 2003;19(3):188-191.
- Lo RS, Woo J, Zhoc KC, et al. Cross-cultural validation of the McGill Quality of Life questionnaire in Hong Kong Chinese. *Palliat Med* 2001;15(5):387-397.
- Lo SH, Chan CY, Chan CH, et al. The implementation of an end-of-life integrated care pathway in a Chinese population. *Int J Palliat Nurs* 2009;15(8):384-388.
- Löfmark R, Nilstun T, Cartwright C, et al. Physicians' experiences with end-of-life decision-making: survey in 6 European countries and Australia. *BMC Med* 2008;6:4-4.
- Logie DE, Harding R. An evaluation of a morphine public health programme for cancer and AIDS pain relief in Sub-Saharan Africa. *BMC Public Health* 2005;5.
- Lorensen M, David AJ, Konishi E, Bunch EH. Ethical issues after the disclosure of a terminal illness: Danish and Norwegian hospice nurses' reflections. *Nurs Ethics* 2003;10(2):175-185.

Lovell DM, Hemmings G, Hill AB. Bereavement reactions of female Scots and Swazis: a preliminary comparison. *Br J Med Psychol* 1993;66(3):259-274.

Low J, Pattenden J, Candy B, Beattie JM, Jones L. Palliative Care in Advanced Heart Failure: An International Review of the Perspectives of Recipients and Health Professionals on Care Provision. *J Card Fail* 2011;17(3):231-252.

Luczak J, Kluziak M, Hunter GP. Education/training in supportive and palliative care in central and eastern Europe - Results of a questionnaire survey. *Support Care Cancer* 2002;10(4):292-302.

Lynch T, Clark D, Centeno C, et al. Barriers to the development of palliative care in the countries of central and eastern Europe and the commonwealth of independent States. *J Pain Symptom Manage* 2009;37(3):305-315.

Lynch T, Connor S, Clark D. Mapping levels of palliative care development: a global update. *J Pain Symptom Manage* 2012;45(6):1094-1106.

MacKenzie T, Innes J, Boyd M, et al. Evaluating the role and value of a national office to coordinate Liverpool Care Pathway implementation in New Zealand. *Int J Evid Based Healthc* 2011;9(3):252-260.

Marc Ho ZJ, Krishna LK, Goh C, Alethea Yee CP. The physician-patient relationship in treatment decision making at the end of life: a pilot study of cancer patients in a Southeast Asian society. *Palliat Support Care* 2013;11(1):13-19.

Marinangeli F, Ciccozzi A, Leonardis M, et al. Use of strong opioids in advanced cancer pain: a randomized trial. *J Pain Symptom Manage* 2004;27(5):409-416.

Marshall-Lucette S, Corbett K, Lartey N, Opio D, Bikaitwoha ME. Developing locally based research capacity in Uganda. *Int Nurs Rev* 2007;54(3):227-233.

Matsumura S, Bito S, Liu H, et al. Acculturation of attitudes toward end-of-life care - A cross-cultural survey of Japanese Americans and Japanese. *J Gen Intern Med* 2002;17(7):531-539.

McCaffery M, Ferrell BR. Nurses knowledge about cancer pain: a survey of 5 countries. *J Pain Symptom Manage* 1995;10(5):356-369.

Meeussen K, Van den Block L, Echteld M, et al. Advance Care Planning in Belgium and The Netherlands: A Nationwide Retrospective Study Via Sentinel Networks of General Practitioners. *J Pain Symptom Manage* 2011;42(4):565-577.

Minton O, Strasser F, Radbruch L, Stone P. Identification of factors associated with fatigue in advanced cancer: a subset analysis of the European palliative care research collaborative computerized symptom assessment data set. *J Pain Symptom Manage* 2012;43(2):226-235.

Mitera G, Fairchild A, DeAngelis C, et al. A multicenter assessment of the adequacy of cancer pain treatment using the Pain Management Index. *J Palliat Med* 2010;13(5):589-593.

Miyazaki K, Suzukamo Y, Shimozuma K, Nakayama T. Verification of the psychometric properties of the Japanese version of the European organization for research and treatment of cancer quality of life questionnaire core 15 palliative (EORTCQLQ-C15-PAL). *Qual Life Res* 21(2):335-340.

- Murray SA, Grant E, Grant A, Kendall M. Dying from cancer in developed and developing countries: lessons from two qualitative interview studies of patients and their carers. *BMJ*;326(7385):368.
- Mwangi-Powell FN, Powell RA, Harding R. Models of delivering palliative and end-of-life care in sub-Saharan Africa: a narrative review of the evidence. *Curr Opin Support Palliat Care* 2013;7(2):223-228.
- Namisango E, Katabira E, Karamagi C, Baguma P. Validation of the Missoula-Vitas Quality-of-Life Index among patients with advanced AIDS in Urban Kampala, Uganda. *J Pain Symptom Manage* 2007;33(2):189-202.
- Nicklasson M, Bergman B. Validity, reliability and clinical relevance of EORTC QLQ-C30 and LC13 in patients with chest malignancies in a palliative setting. *Qual Life Res* 2007;16(6):1019-1028.
- Nilstun T, Cartwright C, Löfmark R, et al. Access to death certificates: what should research ethics committees require for approval? *Ann Epidemiol* 2006;16(4):281-284.
- Onuoha FN, Munakata T, Serumaga-Zake PA, Nyonyintono RM, Bogere SM. Negative mental health factors in children orphaned by AIDS: Natural mentoring as a palliative care. *AIDS Behav* 2009;13(5):980-988.
- Onwuteaka-Philipsen BD, Fisher S, Cartwright C, et al. End-of-life decision making in Europe and Australia - A physician survey. *Arch Intern Med* 2006;166(8):921-929.
- Pahor M, Rasmussen BH. How does culture show? A case study of an international and interprofessional course in palliative care. *J Interprof Care* 2009;23(5):474-485.
- Paice JA, Ferrell B, Coyle N, Coyne P, Smith T. Living and Dying in East Africa: Implementing the End-of-Life Nursing Education Consortium Curriculum in Tanzania. *Clin J Oncol Nurs* 2010;14(2):161-166.
- Paice JA, Ferrell BR, Coyle N, Coyne P, Callaway M. Global efforts to improve palliative care: the International End-of-Life Nursing Education Consortium Training Programme. *J Adv Nurs* 2008;61(2):173-180.
- Pastrana T, De Lima L, Eisenchlas J, Wenk R. Palliative Care Research in Latin America and the Caribbean: From the Beginning to the Declaration of Venice and Beyond. *J Palliat Med* 2012;15(3):352-358.
- Pastrana T, Vallath N, Mastrojohn J, et al. Disparities in the contribution of low- and middle-income countries to palliative care research. *J Pain Symptom Manage* 2010;39(1):54-68.
- Knapp C, Woodworth L, Wright M, et al. Pediatric Palliative Care Provision Around the World: A Systematic Review. *Pediatr Blood Cancer* 2011;57(3):361-368.
- Pereira J, Bruera E, Quan H. Palliative care on the net: An online survey of health care professionals. *J Palliat Care* 17(1):41-45.
- Pousset G, Bilsen J, Cohen J. Deaths of children occurring at home in six European countries. *Child Care Health Dev* 2010;36(3):375-384.

Raijmakers NJ, van Zuylen L, Constantini M, et al. Issues and needs in end-of-life decision making: An international modified Delphi study. *Palliat Med* 2012;26(7):947-953.

Raijmakers NJ, van Zuylen L, Furst CJ, et al. Variation in medication use in cancer patients at the end of life: a cross-sectional analysis. *Support Care Cancer* 2013;21(4):1003-1011.

Rawlinson F, Finlay I. Assessing education in Palliative Medicine: development of a tool based on the Association for Palliative Medicine core curriculum. *Palliat Med* 2002;16(1):51-55.

Salisbury C, Bosanquet N, Wilkinson EK, et al. The impact of different models of specialist palliative care on patients' quality of life: A systematic literature review. *Palliat Med* 1999;13(1):3-17.

Schiessi C, Walshe M, Wildfeuer S, et al. Undergraduate Curricula in Palliative Medicine: A Systematic Analysis Based on the Palliative Education Assessment Tool. *J Palliat Med* 2013;16(1):20-30.

Schug SA, Zech D, Dörr U. Cancer pain management according to WHO analgesic guidelines. *J Pain Symptom Manage* 1990;5(1):27-32.

Selman LE, Higginson IJ, Aguiro G. Quality of life among patients receiving palliative care in South Africa and Uganda: a multi-centred study. *Health Qual Life Outcomes* 2011;9.

Selman L, Higginson IJ, Aguiro G, et al. Meeting information needs of patients with incurable progressive disease and their families in South Africa and Uganda: multicentre qualitative study. *BMJ* 2009;338.

Seymour JE, Janssens R, Broeckaert B. Relieving suffering at the end of life: Practitioners' perspectives on palliative sedation from three European countries. *Soc Sci Med* 2007;64(8):1679-1691.

Shin DW, Choi JE, Miyashita M, et al. Cross-Cultural Application of the Korean Version of the European Organization for Research and Treatment of Cancer Quality of Life Questionnaire-Core 15-Palliative Care. *J Pain Symptom Manage* 2011;41(2):478-484.

Shin DW, Choi JE, Miyashita M, et al. Measuring the Structure and Process of End-of-Life Care in Korea: Validation of the Korean Version of the Care Evaluation Scale (CES). *J Pain Symptom Manage* 2012;44(4):615-625.

Sigurdardottir KR, Haugen DF, van der Rijt CC, et al. Clinical priorities, barriers and solutions in end-of-life cancer care research across Europe. Report from a workshop. *Eur J Cancer* 2010;46(10):1815-1822.

Silbermann M. Current Trends in Opioid Consumption Globally and in Middle Eastern Countries. *J Pediatr Hematol Oncol* 2011;33(1):s1-s5.

Simoens S, Kutten B, Keirse B, et al. The costs of treating terminal patients. *J Pain Symptom Manage* 2010;40(3):436-448.

Sprung CL, Carmel S, Sjøkvist P, et al. Attitudes of European physicians, nurses, patients, and families regarding end-of-life decisions: the ETHICATT study. *Intensive Care Med* 2007;33(1):104-110.

Steel K, Ljunggren J, Topinková E, et al. The RAI-PC: an assessment instrument for palliative care in all settings. *Am J Hosp Palliat Care* 2003;20(3):211-219.

Stone P, Minton O. European Palliative Care Research collaborative pain guidelines. Central side-effects management: What is the evidence to support best practice in the management of sedation, cognitive impairment and myoclonus? *Palliat Med* 2011;25(5):431-441.

Suárez-del-Real Y1, Allende-Pérez S, Alférez-Mancera A, et al. Validation of the Mexican-Spanish version of the EORTC QLQ-C15-PAL questionnaire for the evaluation of health-related quality of life in patients on palliative care. *Psychooncology* 2011;20(8):889-896.

Tanneberger S, Pannuti F. The Bologna Hospital-at-Home: A model for cost-effective care of advanced cancer patients in developing countries. *Nat Med J India* 1998;11(5):231-235.

Thulesius HO, Scott H, Helgesson G, Lynöe N. De-tabooing dying control: a grounded theory study. *BMC Palliat Care* 2013;12.

Torres-Vigil I, Mendoza TR, Alonso-Babarro A, et al. Practice patterns and perceptions about parenteral hydration in the last weeks of life: a survey of palliative care physicians in Latin America. *J Pain Symptom Manage* 2012;43(1):47-58.

Vachon ML. Myths and realities in palliative/hospice care. *Hosp J* 1986;2(1):63-79.

Vainio A, Auvinen A. Prevalence of symptoms among patients with advanced cancer: an international collaborative study. *J Pain Symptom Manage* 1996;21(1):3-10.

Vermandere M, Lelepeire JD, Van Mechelen W, et al. Spirituality in palliative home care: a framework for the clinician. *Support Care Cancer* 2013;21(4):1061-1069.

Vignaroli E, Wenk R. The challenge of international consensus: defining an opioid essential prescription package. *Curr Opin Support Palliat Care* 2012;6(3):398-401.

Vignaroli E, Bennet MI, Nikolaichuk C, et al. Strategic pain management: The identification and development of the IAHPC Opioid Essential Prescription Package. *J Palliat Med* 2012;15(2):186-191.

Volz R, Akabayashi A, Reese C, Ohi G, Sass HM. Organization and patients' perception of palliative care: a crosscultural comparison. *Palliat Med* 1997;11(5):351-357.

Voorhees J, Rietjens J, Onwuteaka-Phillipsen B, et al. Discussing prognosis with terminally ill cancer patients and relatives: A survey of physicians' intentions in seven countries. *Patient Educ Couns* 2009;77(3):430-436.

Vosit-Steller J, Morse AB, Mitrea N. Evolution of an international collaboration: A unique experience across borders. *Clin J Oncol Nurs* 2011;15(5):564-566.

Warren JL, Barbera L, Bremner KE, et al. End-of-life care for lung cancer patients in the United States and Ontario. *J Natl Cancer Inst* 2011;103(11):853-862.

Wenk R, Diaz C, Echeverria M, et al. Argentina's WHO Cancer Pain Relief Program: a patient care model. *J Pain Symptom Manage* 1991;6(1):40-43.

Wenk R, De Lima L, Eisenchlas J. Palliative care research in Latin America: Results of a survey within the scope of the Declaration of Venice. *J Palliat Med* 2008;11(5):717-722.

Wiese CH, Lassen CL, Bartels UE, et al. International recommendations for outpatient palliative care and prehospital palliative emergencies - a prospective questionnaire-based investigation. *BMC Palliat Care* 2013;12.

Woitha K, Van Beek K, Ahmed N, et al. Development of a set of process and structure indicators for palliative care: the Europall project. *BMC Health Serv Res* 2012;12.

Shin DW, Choi J, Miyashita M, et al. Measuring comprehensive outcomes in palliative care: Validation of the Korean version of the Good Death Inventory. *J Pain Symptom Manage* 2011;42(4):632-642.

Wright M, Wood J, Lynch T, Clark D. Mapping levels of palliative care development: a global view. *J Pain Symptom Manage* 2008;35(5):469-485.