P-195

'PALLIATIVE CRITICAL CARE': TRANSFORMING THE RELATIONSHIP BETWEEN INTENSIVE CARE AND PALLIATIVE CARE

Helen Crispin, Carol Davis, Robert Chambers, Karen Haynes. *Southampton General Hospital, Southampton, UK*

10.1136/bmjspcare-2018-hospiceabs.220

Introduction Historically, joint-working between our General Intensive Care Unit (GICU) and Hospital Specialist Palliative Care Team (HSPCT) was restricted to occasional complex cases. In 2013 there was shared recognition that patients, families and staff might benefit from a broader, structured, collaborative approach.

In 2014, we created four hours dedicated intensivist time and enhanced joint working – 'Palliative Critical Care'. In 2016, a weekly combined ward round was established.

Here, we report the impact of this Quality Improvement Initiative on HSPCT activity and place of care.

Methods Retrospective review of HSPCT records between April 2013 and March 2018.

Results Over 300 patients have been seen under this collaborative model since April 2013. Initially, referrals increased from six in 2013/14 to 57 in 2014/15. Continuous development has resulted in sustained year on year increase in referrals; 107 in year 5 (2017/18).

Collaborative working has promoted recognition and treatment of symptoms, and provided an additional layer of support for patients, families and staff. Staff report increased confidence especially in symptom control of actively dying patients. The in-hospital mortality of this patient group is 46%–74% with an ICU mortality of 23%. Patient preferences have been supported; 61.5% of survivors were discharged home or to a hospice (2017/18). In selected cases discharge has been achieved direct from GICU; seven patients to a hospice and eight to home (2013/14 – 2016/17). Patient, family and staff feedback has been 'excellent'; one relative said '*I had no idea that palliative care and intensive care could work together especially for someone who doesn't have cancer and isn't dying!*'

Conclusion Through this Quality Improvement initiative the relationship between our GICU and HSPCT has been transformed from one of occasional interaction to a regular programme of collaborative working resulting in a sustained increase in referrals and supported patient preferences.

P-196

EXPLORING THE QUALITY OF THE DYING AND DEATH EXPERIENCE IN THE EMERGENCY DEPARTMENT: A REVIEW

^{1,2}Kay McCallum, ^{1,3}Debra Jackson, ¹Helen Walthall, ¹Helen Aveyard. ¹Oxford Brookes University, Oxford, UK; ²Oxford University Hospitals NHS Foundation Trust, Oxford, UK; ³University of Technology Sydney, Sydney, Australia

10.1136/bmjspcare-2018-hospiceabs.221

Background Patients attend the emergency department (ED) for a variety of reasons. Unfortunately some of these patients will not survive. Little is known about how staff and particularly carers experience being with a dying patient in this setting (Bailey, Murphy & Porock, 2011; Chan, 2011; Decker, Lee & Morphet, 2015). The aim of this review was to examine the quality of dying and death in the ED from the perspective of staff and carers.

Method A systematic search of MEDLINE, the Cumulative Index to Nursing and Allied Health Literature (CINAHL), AMED, Magonline (internurse), EMBASE and the Cochrane library was undertaken during February –August 2017. Fourteen articles fit the inclusion criteria (papers published in the English language between 1995–2017) for this review.

Findings and discussion The over-arching theme that emerged is that *care in the ED is focused on living not dying*. Because the ethos is about saving lives, those whose lives cannot be saved can be marginalised or side-lined, leading to a potentially poorer experience for the relatives and friends (carers) (Bailey, Murphy & Porock, 2011). Further themes included staff stress and distress, and the difficulty of meeting the needs of the carers in this environment. Carers felt the staff had no time for them and that communication was generally poor (Bailey, Murphy & Porock, 2011). However, there are minimal studies that asked the views of the carers and this is therefore an area in which further research is needed.

Implications for research and practice Findings suggest that there remains a gap in knowledge around the quality of dying and death in the ED from the perspective of carers of the dying patient. Supporting staff in the ED to care effectively for patients who are dying whether from acute or non-acute causes is paramount.

P-197

THE IMPACT OF A HOSPITAL BASED, VOLUNTEER-LED END OF LIFE SERVICE ON STAFF WELLBEING

Kim Bonnar, Julia Bearn. *Marie Curie, London, UK*

10.1136/bmjspcare-2018-hospiceabs.222

Background In 2014, a hospital in the South West of England and an end of life care charity introduced a service of specially trained volunteers to provide emotional and practical support to dying patients in hospital, and their friends and family.

Aims Alongside providing support to patients and families, the service was developed in acknowledgment of increasing pressure on staff. In particular, staff can have limited time to provide one-to-one support at the end of life (Nursing Standard/Marie Curie survey, 2018).

Method To identify and understand any potential impacts of the service on NHS staff, nine semi-structured interviews were conducted. The sample included consultants, nurses, receptionists and chaplaincy staff to allow for the exploration of a range of different perspectives. A thematic analysis was conducted using the interview transcripts.

Results Three main staff benefit themes were identified: relieving feelings of professional guilt, being able to offer patients and families options, and the service's responsiveness and reliability. Staff spoke of challenges in giving patients their time on busy hospital wards. Interviewees spoke of the professional guilt experienced when they could not provide one-to-one support. The introduction of a reliable, well regarded volunteer service however allowed them to offer options for patients, alongside giving families opportunities for respite. In turn, staff described how the service helped to relieve feelings of professional guilt.

Conclusion This work suggests the service has a positive impact on staff, particularly in terms of relieving feelings of professional guilt. This is acutely relevant in a time of high

levels of workforce pressure (Nursing Standard/Marie Curie survey, 2018) This work also highlights the importance of considering impacts of volunteer services beyond immediate beneficiaries.

P-198 REDUCING AVOIDABLE READMISSIONS FOR PEOPLE WITH COMPLEX NEEDS ASSOCIATED WITH FRAILTY

Kerry Bareham. *St Barnabas Lincolnshire Hospice, Lincoln, Lincolnshire*

10.1136/bmjspcare-2018-hospiceabs.223

Older people are the most frequent users of services and when admitted to hospitals are at greater risk of complications leading to longer lengths of stay and readmission rates. The current design of secondary care services fail to identify the complex underlying needs of older people associated with frailty, including palliative care needs. This results in poor discharge planning and subsequent avoidable future admissions. However, for those people who are admitted to community palliative care services their risk of a readmission to hospital within 28 days is halved.

This presentation aims to share the findings of a service transformation project implemented by the author as part of their MSc dissertation. This was achieved with the application of an Experienced Based Design approach with other stakeholders, along with the principles of Six Sigma to understand the root cause, and was supported by a systematic review of the literature. Which informed the case for the role of an Anticipatory Care Nurse based in A and E of an acute trust to educate front door clinicians how to identify people in their last year of life and develop person centred care planning.

An action research and rapid appraisal technique was applied through the use of the PDSA cycle was implemented to evaluate the service transformation project.

A 45% increase in the number of referrals to the Specialist Palliative Care Team and subsequent increase in referrals to community palliative care services was demonstrated. This was achieved by improved identification of need by front-line clinicians, resulting in widening access to hospice and palliative and end of life care and subsequent person-centred care planning and well-being. Thus, offering a solution to the STP aspiration to reduce avoidable admissions to hospital by 30%. Lincoln University are keen to support further research through expansion of this project.

P-199 CAN WE ENABLE MORE PEOPLE TO LIVE WELL AND DIE IN A PLACE AND MANNER OF THEIR CHOOSING?

Keri Thomas. *The National GSF Centre for End of Life Care, London, UK*

10.1136/bmjspcare-2018-hospiceabs.224

Aims A third of hospital patients are in their last year of life, yet approximately 12% are seen by specialist palliative care leaving generalists to care for 88% of patients. Generalists need to be enabled to provide good palliative care for their patients. The transition from reactive to proactive care by earlier recognition and assessment of personal needs ensures patients receive better coordinated care in a place and manner of their choosing.

Gold Standards Framework (GSF) is a quality improvement programme delivering proactive person-centred care for people

in the last years of life in all settings. We report on the introduction and impact of the GSF Acute Hospitals training Programme, with findings from the 12 whole hospital programmes and accredited wards.

Method The structured two year GSF Hospital training programme includes training, tools, support and evaluation at organisational, patient and staff level, including measures of progress and attainment. The programme is delivered to the whole hospital (commissioned) or individual wards (open). GSF encourages teams to identify patients earlier (aiming for the 30% figure where appropriate), assess their needs (offering all initial advance care planning discussions) and proactively planning care aligned to preferences, enabling more to die at home if that is their wish.

Results Evaluations indicate significant progress in several key areas in many hospitals and a 'culture change' on the wards. GSF Accredited wards show embedded good practice including earlier identification (average 34%), more proactive needs-based approach, more offered advance care planning discussions (76%–87%), better communication with GPs, reduced hospitalisation and improved more appropriate referrals to palliative care specialists.

Conclusions Findings from the GSF trained hospitals and accredited wards show significant changes, with proactive, better coordinated, person-centred care. A key factor is earlier recognition and accredited wards are able to demonstrate they are identifying 30% of patients.

P-200 QUALITY IMPROVEMENT IN ACTION – IMPROVING CARE FOR PATIENTS ADMITTED TO HOSPITAL IN AN EMERGENCY

Julia Grant, Karen Murray, Valerie Oliviant, Lizzie Hyde, Kate Russ. *University Hospitals Leicester, Leicester, UK*

10.1136/bmjspcare-2018-hospiceabs.225

Background University Hospitals Leicester (UHL) joined a quality improvement collaborative (ECLHIP) to improve care for patients who maybe in their last three months of life who attend or are admitted to hospital in an emergency. Hospitals are an important provider of care for this group. UHL is one of the largest Trusts in the English NHS, caring for over 2000 patients in their last days of life each year. An emergency admission may indicate underlying clinical decline. Quality of care varies and there may be unwanted inpatient stays and/or treatments contributing to overall experience.

Method The collaborative begins with diagnosis (activity data, qualitative casefile review, patient experience walkthrough, review of reports and plans). A driver diagram illustrates the 'theory of change' based on findings to plan improvements. The model for improvement provides the framework for engaged frontline clinicians to conduct tests of change. The approach includes informative evaluation, patient and relative experience and planning for sustainability.

Results 62% (n=1991) emergency, adult patients who died in UHL had an outpatient appointment, emergency admission, attendance and/or planned admission in the 90 days preceding their final admission. Casefile review: 7/10 patients had likely missed opportunities to plan ahead. During the final admission, uncertain recovery and/or dying was recognised but this was often late and/or did not translate into desired action. A survey of A and E staff indicated gaps in knowledge on how to access SPC (24% confident) and communication around