ACP for frail elderly and patients with complex chronic health problems as well as evidence for ACP interventions in this population. ACP education of Dutch elderly care physicians will be highlighted. The specific challenge for elderly care physicians in ACP for persons with decision-making disabilities will be discussed. Finally, the collaboration between elderly care physicians and general practitioners will be illustrated.

Content Introduction ACP in elderly care medicine – by dr. Daisy J.A. Janssen

Specific challenges for elderly care physicians: ACP and decision-making disabilities – by prof. Cees Hertogh

Collaboration between elderly care physician and general practitioner – by general practitioner invited by dr. Annicka van der Plas

OP77

THE ADVANCE PROJECT: AN AUSTRALIAN NATIONAL PROGRAM TO SUPPORT NURSES TO INITIATE ADVANCE CARE PLANNING IN GENERAL PRACTICE

¹J Clayton*, ¹S Nagarajan, ²V Lewis, ³K Detering, ⁴E Halcomb, ⁵J Tieman, ⁶G Mitchell, ⁷J Phillips, ⁸R Morton, ⁹D Costa, ⁹J Gavin, ¹J Stokes, ⁸A Livingstone, ⁸T Shaw, ¹J Rhee. ¹HammondCare, Greenwich, New South Wales, Australia; ²LaTrobe University, Melbourne, Australia; ³Austin Health, Melbourne, Australia; ⁴University of Wollongong, Wollongong, Australia; ⁵Flinders University, South Australia, Australia; ⁶University of Queensland, Queensland, Australia; ⁷University of Technology, Sydney, Australia; ⁸University of Sydney, Sydney, Australia; ⁹Queensland University of Technology, Australia

10.1136/spcare-2019-ACPICONGRESSABS.77

Background Primary care has been advocated as an ideal setting to initiate advance care planning (ACP). Few studies have examined the role of general practice nurses (GPNs) in promoting/initiating ACP. The Advance Project evaluated initiatives to address this gap.

Methods This Australian Government-funded program aims to increase GPNs' confidence in initiating conversations with patients/carers about ACP during routine health assessments with older and/or chronically ill patients using a structured interview. This is part of a broader program enabling GPNs to identify patients at risk of deteriorating and dying and to assess these patients' palliative/supportive care needs. Identified needs are then addressed in consultation with General Practitioners. The program includes a suite of resources and multi-component training (online, face-to-face and individual tele-mentoring). Pre/post/follow-up surveys and qualitative interviews collected GPNs' perspectives about the training/resources and barriers to implementation in clinical practice.

Results As of 31 December 2017, 823 GPNs enrolled in training and 536 completed one or more training components. 27 workshops were held across Australia, including 182 regional/rural participants. 585 pre-training, 384 post-training, and 125 follow-up surveys were received. 20 GPNs were interviewed. There were significant improvements in GPNs' confidence, comfort, knowledge and attitudes towards initiating ACP post-training that was sustained at follow-up. Participants were significantly more likely to have had ACP discussions with their patients at follow-up (81%) compared to baseline (55%, $p < 0.001$).

Discussion/Conclusion GPNs can have an important role in initiating ACP. The evaluation informed refinement/expansion of the resources/training to support team-based initiation of ACP in general practice <http://www.theadvanceproject.com.au>

OP78

REGIONAL (CENTRAL) VERSUS INSTITUTIONAL: COMPETING STRATEGIES FOR NATIONWIDE ACP IMPLEMENTATION

¹J In der Schmitzen*, ²G Marckmann. ¹Medical Faculty, Heinrich-Heine-University, Düsseldorf, Germany; ²Medical Faculty, LMU, Munich, Germany

10.1136/spcare-2019-ACPICONGRESSABS.78

Background Implementing ACP in nursing homes is often essentially done by educating selected n/h staff as ACP facilitators. Recent German legislation covers ACP offered to nursing home residents, and offers an alternative strategy for implementation, i.e. cooperation of participating n/hs with a regional (central) partner that employs a team of facilitators. Which of these two strategies should be preferred?

Methods 1. Follow-up of facilitator training effectivity in Germany in 2015–2017, 2. review of the literature, 3. theoretical analysis of the competing rationales.

Results Of some 270 facilitator trainees attending our ACP courses, only few report ongoing practice as an ACP facilitator. A number of important publications describe facilitators and barriers, or essential elements, of successful ACP implementation, but few if any compare regional versus institutional implementation strategies of ACP yet. Similarly, while regional ACP coordination is described as an important precondition for sustainable ACP implementation, it requires significant resources on top of institutional implementation. A comparative analysis yields a number of strong reasons why regional may well beat institutional implementation strategies, referring to staff aptitude, team building, regional coordination, economic efficiency, and both sustainability and expandability. Arguments that have been raised against qualifying external staff can be shown not to consider sufficiently the potential of creating regional (central) facilitator teams.

Conclusion Regional implementation of ACP, characterised by regional (central) facilitator teams cooperating with nursing homes and other institutions, has yet rarely been described, but poses a substantial potential when compared to conventional institutional implementation strategies that deserves scientific evaluation.

OP79

CREATING MOMENTUM AND CONSISTENCY WITH A NATIONAL FIVE YEAR STRATEGY IN NEW ZEALAND

L Manson*, L Price. Health Quality Safety Commission, Wellington, New Zealand

10.1136/spcare-2019-ACPICONGRESSABS.79

Background The development of ACP in New Zealand was driven by the ACP Cooperative, a grass roots organisation of clinicians. The Cooperative's aim was to drive consistency and address the barriers to ACP implementation. Over time, the lack of an official mandate and funding threatened the sustainability of the work of the Cooperative.

Method In 2017 the Cooperative partnered with the Health Quality & Safety Commission – a crown agency tasked with supporting the public health sector to improve the quality and safety of services. Together they presented a business case to the district health boards (DHBs) to agree to a national programme with a clear mandate and funding. The DHBs agreed to a five-year strategy and roadmap of national and local actions aimed at increasing ACP activity and addressing sustainability. The key strategy workstreams and their aims are:

- promotion: that future health care planning and end of life discussions are normalised in society
- resources: ACP is accessible to all regardless of language, literacy level or cultural beliefs
- education and training: we have workforce and community prepared to have conversations and use ACPs
- monitoring and evaluation: we know care is based on what matters to consumers
- implementation: we are maximising value for DHBs

Result a national mandate, strategy and specified actions have resulted in an increase in ACP activity; increased governance with decision-making being supported by a representative Steering Group; wider national stakeholder engagement and buy in from agencies in and outside of the health sector.

OP80

SHARED CARE PLANNING: A NEW MODEL TO INTEGRATE ADVANCE CARE PLANNING INTO COMMUNITY. THE BASQUE COUNTRY EXPERIENCE

I Saralegui*. *Organización Sanitaria Integrada Araba, Vitoria-Gasteiz, Spain*

10.1136/spcare-2019-ACPICONGRESSABS.80

In the Basque Country, a project has been implemented based on community education, the training of health and social workers. It tries to elicit the individual's preferences and encourages them to take part in planning their care. Writing down advance directives is not the main aim.

The target population is chronic patients, elderly people and anyone at the end of life. The project also includes everyone who wants to think about the process of dying and needs to have a conversation related to this topic.

The project started in 2014 as a bottom-up project and it has grown into a top-down project performed in the whole Basque Health Service.

Results More than one hundred conferences and debates have been taking place in neighborhood associations, cultural centers, libraries or educational centers. More than five thousand people have attended these activities. One thousand workers have attended a basic training course, more than 500 workers have been trained as SCP facilitators and more than 700 doctors and nurses have participated in conversations with patients and families helped by an SCP facilitator. In 75% of cases, the citizens asked to be included after attending a conference; only 25% of participants were included because of a doctor or nurse's invitation. Many support documents have been created in order to explain the project and to make it easier to understand.

The keystone is the training of health and social workers in order to answer citizens' requests and to integrate the conversations into everyday care.

OP81

IMPLEMENTATION OF ADVANCE CARE PLANNING IN AGED CARE: WHAT DO YOU WANT AND HOW CAN WE HELP?

¹H Kelly, ¹L Nolte, ²M Fearn, ²F Batchelor, ²B Haralambous, ²P Mackell, ²K Hwang, ¹K Detering. ¹*Advance Care Planning Australia, Melbourne, Australia*; ²*National Ageing Research Institute Inc, Melbourne, Australia*

10.1136/spcare-2019-ACPICONGRESSABS.81

Introduction The goal of advance care planning (acp) is to ensure that individuals receive future care consistent with their expressed preferences. Benefits of acp include: increased adherence to a person's preferences, higher staff satisfaction, reductions in unwanted hospitalisations and medical treatments, and reduced stress and anxiety for family. Despite clear benefits, uptake of acp has been limited in Australian aged care settings. This research explored barriers and enablers to implementing acp in residential and community aged care settings, from the perspectives of aged care staff and older people.

Method Focus groups and interviews with aged care staff and older people were conducted in four Australian states. Sixty staff from 15 residential and community aged care organisations and 24 older people participated. A thematic analysis was undertaken to identify facilitators and barriers.

Results Factors identified as impacting on the experiences and implementation of acp included: knowledge and understanding access to education/training understanding of relevant legislation having defined roles/responsibilities cognitive capacity of the older person timing of initiating the conversation availability of clear policies/procedures engaging older people and their families diversity within the workforce and older people.

Conclusions Aged care staff indicated there are a range of factors that can impact on acp implementation. All these factors may need to be considered when implementing acp within Australian aged care organisations. The findings informed the development of a new national resource titled *Advance care planning in aged care: a guide to support implementation in community and residential settings*.

OP82

IDENTIFICATION, IMPLEMENTATION AND EVALUATION OF INDICATORS TO MONITOR SUCCESSFUL UPTAKE OF ADVANCE CARE PLANNING IN ALBERTA

^{1,2}J Xiao*, ³J SimOn, ⁴T Lynn Wityk Martin, ⁴S Iversen, ¹M Douglas, ¹A Potapov, ¹M Nesari, ³P Biondo, ³A Kanter, ^{1,2}K Fassbender. ¹*University of Alberta, Edmonton, Canada*; ²*Covenant Health Palliative Institute, Edmonton, AB, Canada*; ³*University of Calgary, Calgary, Canada*; ⁴*Alberta Health Services, Edmonton, AB, Canada*

10.1136/spcare-2019-ACPICONGRESSABS.82

Background In 2014, a province-wide policy for advance care planning (ACP) and Goals of Care Designation (GCD) was implemented in Alberta, Canada; nevertheless, few quality indicators have been rigorously developed or evaluated for measuring the uptake of ACP/GCD.

Methods In phase I, we performed a systematic literature review and environmental scan to identify potential ACP/GCD indicators. A Delphi consensus-based approach, consisting of 3 rounds of face-to-face meetings and/or online surveys, was used to develop a short list of indicators. In phase II, the panelists met face-to-face to operationalize and implement the indicators. In phase III, two validated questionnaires and semi-structured interviews of 60 individuals (stratified by manager/practitioner and physician/nurse) are being used to evaluate the usability and acceptability of the implemented indicators on a dashboard interface.

Results A total of 132 potential indicators were identified in phase I. The indicators were reduced and refined to 18 after 3 Delphi rounds. Phase II resulted in 9 valid and feasible indicators in a measurable format (i.e. numerator, denominator, data source defined). The Phase III protocol is under