OPEN ACCESS

Persistent inequalities in Hospice at Home provision

Jackie Buck,¹ Liz Webb,² Lorraine Moth,² Lynn Morgan,² Stephen Barclay³

¹Faculty of Medicine and Health Sciences, University of East Anglia, Norwich, UK

²Arthur Rank Hospice Charity, Cambridge, UK

³Primary Care Unit, Department of Public Health and Primary Care, Institute of Public Health, University of Cambridge, Cambridge, UK

Correspondence to

Dr Jackie Buck, Faculty of Medicine and Health Sciences, University of East Anglia, Norwich, NR4 7TJ, UK; J.Buck@uea.ac.uk

Received 18 April 2017

Revised 20 December 2017

Accepted 3 January 2018

ABSTRACT

Objective To describe the nature and scope of a new Hospice at Home (H@H) service and to identify its equality of provision.

Methods Case note review of patients supported by a H@H service for 1 year from September 2012 to August 2013 (n=321). Descriptive analysis to report frequencies and proportions of quantitative data extracted from service logs, referral forms and care records; thematic analysis of qualitative data from care record free text.

Results Demand outstripped supply. Twice as many night care episodes were requested (n=1237) as were provided (n=613). Inequalities in access to the service related to underlying diagnosis and socioeconomic status. 75% of patients using the service had cancer (221/293 with documented diagnosis). Of those who died at home in the areas surrounding the hospice, 53% (163/311) of people with cancer and 11% (49/431) of those without cancer received H@H support. People who received H@H care were often more affluent than the population average for the area within which they lived. Roles of the service identified included: care planning/implementation, specialist end-of-life care assessment and advice, 'holding' complex patients until hospice beds become available and clinical nursing care.

Conclusion There is significant unmet need and potentially large latent demand for the H@H service. People without cancer or of lower socioeconomic status are less likely to access the service. Action is needed to ensure greater and more equitable service provision in this and similar services nationally and internationally.

HOW THIS FITS IN

Hospice at Home services are being implemented nationally and internationally to support the growing numbers of people wishing to die at home. With considerable variation in practice, little is known about how these services function. This evaluation of one service found demand

far outstripped supply and revealed inequality of access, patients with cancer being five times more likely to receive Hospice and Home care than those dying from non-malignant disease.

INTRODUCTION

UK healthcare policy has seen an increasing focus on palliative and end-of-life care over recent years, stimulated by the 2008 National Health Service (NHS) End of Life Care Strategy, that sought to ensure that people approaching the end of their lives receive the best care possible and are as involved in decisions regarding their care as they wish.¹

One such decision concerns where individuals would prefer to be cared for towards the end of their lives. While for many the stated preference is to be at home, this varies across clinical conditions²⁻⁴ and may decrease with older age as people fear experiencing greater dependency due to chronic disease and are concerned about becoming a burden to family members.⁵⁻⁹ Recent reviews have highlighted how many studies reporting preferred place of death included high proportions of respondents with no recorded or stated preference regarding where they would like to die.¹⁰⁻¹¹ In light of the most prevalent place of death in many countries being hospital,¹²⁻¹⁶ the UK hospice movement has responded with the development of 'hospice at home' (H@H) services. These provide more intensive nursing care in the home than is usually available¹⁷ seeking to enable patients with advanced illness considered to be approaching the end of life, to be cared for and to die at home, if that is their preference. H@H aims to reduce inpatient admissions, facilitate discharge from inpatient care, provide assistance during crises and to provide support during the final days of life at home.

To cite: Buck J, Webb L, Moth L, et al. *BMJ Supportive & Palliative Care* Epub ahead of print: [please include Day Month Year]. doi:10.1136/bmjspcare-2017-001367

Several H@H services embrace the fundamental paradigm of end-of-life care, with an acceptance of death as a natural end of life to be enabled at home if possible, rather than a defeat of modern medicine to be fought in hospital.¹⁸ H@H services give considerable importance to autonomy in enabling people to die at home if so wished while providing optimum comfort (beneficence) and avoiding unnecessary investigations and admissions (non-maleficence). They are believed to be more cost effective than inpatient care (justice).¹⁹

The international literature identifies several factors that are known to adversely impact on the likelihood of dying at home, including limited availability and use of home care services to support patients and their informal carers,²⁰ non-cancer diagnosis, greater socioeconomic deprivation, older age, non-Caucasian ethnicity and female gender^{2 16 21–24} though the effect of some of these factors varies between countries. Fear of the dying process and among patients, their families and healthcare professionals also plays a role in preventing people from dying at home.²⁵ Accurate prognostication and timing of palliative care service introduction is challenging for frail older people with multiple comorbid conditions.^{26 27}

While there is some evidence that community-based palliative care teams may reduce acute care use at the end of life and hospital deaths,²⁸ international studies of the effectiveness of H@H services are limited by the diverse models of care, which have often been developed in response to varying local needs.^{17 29} The small body of evidence suggests bereaved lay carers and healthcare professionals have broadly favourable opinions and experiences of H@H.^{30–33} Death at home with community support is associated with lower health resource costs compared with death in hospital or hospice.^{34 35}

H@H and similar home-based palliative services continue to be developed across the UK and in other countries, yet few studies have described in any detail the care provided. This paper describes for one service:

1. Service recipients
2. Sources and appropriateness of referrals
3. Service supply in relation to demand
4. Equity of service provision
5. Service roles and functions

METHODS

Setting and subjects

Based at the Arthur Rank Hospice in Cambridge, UK, the Cambridgeshire H@H service covers an urban and large rural area, with a population of 620 000 and low levels of ethnic diversity (84.5% White British). While the population of Cambridge city is young (mean age 35.6 years), the mean age of the rest of the county (39.5 years) more closely mirrors the age of the general population (40.0 years).³⁶

Most episodes of H@H care are delivered by a team member staying with a patient overnight from

22:00 to 07:00. For the purposes of this evaluation, an episode of care was defined as a period of night care. Referrals are triaged by the senior nurse during the day and assigned a red, amber or green rating based on the severity of clinical need of the patient and family with care prioritised for those assessed as red priority. The level of symptom control required and the emotional welfare of the patient, family or carers is taken in to account if documented on the referral forms or during discussions with referring healthcare professionals. Those requiring a higher level of physical or psychological care are assigned to registered nurses, those with less complex needs to healthcare assistants. All team members have had advanced education in communications skills and end-of-life care. During the course of the evaluation the team expanded from 1 Clinical Nurse Specialist manager and 1 qualified nurse to 6 nurses, 2 bank nurses, 14 healthcare assistants, a complementary therapist and 2 part-time administrative assistants. The service assessed referrals to be appropriate if patients were estimated to be in the last 2 weeks of life.

Data collection

The study population comprised all patients accepted for care by the Cambridgeshire H@H service between 1 September 2012 and 30 August 2013. Individual patient level data were obtained from referral forms, clinical records and study-specific 'night care' documents (table 1). These data were extracted by research administrators and entered onto a study spreadsheet with a study identification number for each patient.

Data concerning place of death for all adults within Cambridgeshire were obtained from the local Public Health Intelligence Team. The patient's GP practice was used to assign patients to an area of the County.

Socioeconomic status was defined using the Index of Multiple Deprivation 2010 based on patient post-codes, which were collected and held separately from the main dataset in order to maintain anonymity. Diagnosis was established from certified underlying cause of death; 'living alone' indicated usually lived alone with no resident carer; 'living at home' indicated living in own home or care home.

Analysis

Descriptive quantitative analysis of frequencies and percentages was undertaken within MS Excel 2010. Free text data extracted from the night care documents were analysed using thematic analysis, which aims to bring order and structure to data by identifying themes and assessing whether there is any relationship or variation between them.³⁷ Qualitative data management was facilitated by NVivo10. All analysis was undertaken by JB who was a nurse researcher at the University of Cambridge.

Table 1 Sources of data

Data source	Completed by	Data items
H@H provision of service log (completed each day of the evaluation)	H@H lead nurse	Number of: episodes of care provided, episodes of care requested but not fulfilled, episodes of care declined per day*
Referral forms	Person referring patient to H@H service	Reasons for referral (tick boxes) Referrer role Patient's General Practice
Hospice-based patient communication/nursing notes	H@H nurses and healthcare assistants	Age, diagnosis, residential status, medications and treatments on referral, medications and treatments and service use during H@H service input, input of other services, telephone calls made and received, evidence of discussions relating to preferred place of care and death and 'do not resuscitate' status, carer information, cause of death
Night care documents	H@H nurses and healthcare assistants after provision of episode of overnight care	Overnight actions including: care given, 'just in case' medications administered, contact with out of hours services, plus comments transcribed verbatim from the free text comments box for in which H@H staff recorded any other information they wanted to include

*Episode of care=care overnight.

H@H, Hospice at Home.

RESULTS

During the 12-month study period, 321 patients received care from the H@H service: 1122 episodes of night care were provided (median 2 per patient, range 0–48). Tables summarising results below report varying numbers in analysis due to missing data.

Referrals, patient characteristics and place of death

Patient characteristics are described in table 2 and referrer information in table 3. The majority of patients who received care from the H@H service were 78 years old or older, three quarters had a cancer diagnosis, just under half were women. One in five patients lived alone. Patient's ethnicity was recorded by the clinical team in less than 15% of cases and so is not reported here. Most patients died within 1 week of referral (55%, n=152): 6% (n=17) died within 24 hours and 13% (n=37) were referred more than a month before death (range 1–204 days between referral and death). The predominant reasons for referral were psychological support for patient and family (54%, n=233) and symptom control (52%, n=118) (multiple reasons given).

Of those with a recorded preferred place of death (n=283) 91% (n=259) died in their preferred place, see table 4.

Table 2 Patient demographics

Characteristic	Number for analysis*	Median	Range
Age	293	78	(22–104)
Distance from hospice base (km)	290	12	(<1–74)
		%	Number
Female	293	46	(n=135)
Lives alone	290	20	(n=58)
Had cancer diagnosis	281	75	(n=221)

*Missing data: age and gender=28, living arrangement and distance from hospice n=31, diagnosis n=40.

Supply and demand for H@H service

Both supply (episodes of care provided) and demand (episodes of care requested) rose steadily throughout the evaluation (figure 1). Supply rose almost three-fold over the 12 months, a time when the service was expanding considerably, with 178 episodes of care provided in August 2013 compared with 64 in September 2012. Unmet need, based on episodes of care requested but not provided, also rose during the study period, with some fluctuation. H@H provided at least one episode of care for 27% (200/742) of all deaths at home in the three areas closest to the hospice.

Equity of service provision

In the two areas most often served by H@H (South Cambridgeshire and Cambridge City), there was a trend for those receiving the service to be less socially deprived than the populations within which they lived (table 5). Very few patients in Fenland, the most rural and deprived area in the county, received the service.

H@H provided care for 11% (49/431) of those who died at home with conditions other than cancer and 53% (163/311) of those who died at home with cancer in the three areas closest to the hospice (Cambridge

Table 3 Referrer characteristics (n=287)

Source of referral*	%	n	Most frequent referrer group from each source	Days from referral to death (median)
Community†	45	(129)	Community Nurses	7
Hospice‡	35	(101)	Palliative Care Clinical Nurse Specialists	7
Hospital§	20	(57)	Discharge Planning Nurse Specialists	10

*Missing data: referrer not documented for 34 patients.

†Community included nurse and general practitioners.

‡Hospice included clinical nurse specialists and hospice doctors.

§Hospital included discharge planning nurse specialists and all other referrals from within the hospital.

Table 4 Preferred and actual place of death of patients cared for by H@H service (n=283)*

Preferred place of death (n=283)	Place of death									
	Home (n=223)		Hospice (n=43)		Hospital (n=4)		Nursing home (n=5)		Unknown (n=18)	
	%	n	%	n	%	n	%	n	%	n
Home	86	(227)	91	(207)	3	(6)	1	(2)	<1	(1)
Hospice	8	(24)	4	(1)	87	(21)	–	–	4	(1)
Either home or hospice	10	(29)	41	(12)	55	(16)	7	(1)	–	–
Hospital	<1	(1)	–	–	–	–	100	(1)	–	–
Nursing home	1	(2)	–	–	–	–	–	–	100	(2)

*Preferred place of death not recorded for 38 patients.

City, South Cambs. and East Cambs., data not available for Huntingdonshire and Fenland).

Roles and functions

Data from 736 night care documents (65% completion rate) and 1519 telephone call records involving H@H team members revealed a wide range of roles performed by the H@H team (table 6). At times staff reported having ‘gone in cold’ to a patient’s home with little information: this was particularly challenging when they encountered a home environment that was unsuited for end-of-life care.

Organisational and specialist roles

The H@H service played an important role in communication between care professionals regarding

the planning and delivery of care, as documented in telephone call data. Most calls were from H@H to family and carers (35%, n=531) or healthcare professionals (35%, n=526). Many conversations were with the community nurses responsible for care during the day, including reports on the patient’s clinical condition, discussion of syringe driver drugs or anticipatory medication. Twenty-one per cent (n=312) of calls were from healthcare professionals to H@H, while 7% (n=105) were from family and carers. The H@H service was at times used to support patients overnight who were awaiting hospice admission.

The team’s palliative care Clinical Nurse Specialists also visited patients at home during the day to assess care needs and provided specialist advice for healthcare

Figure 1 Supply and demand for Hospice at Home Service

Table 5 Index of multiple deprivation scores (n=253)

Zone	Number in sample (n)	Sample mean	SD of sample mean	95% CIs of sample mean		Population mean
				Lower limit	Upper limit	
South Cambs.	107	6.1	3.5	5.5	6.8	6.9
Huntingdonshire	8	11.7	8.6	5.7	17.6	10.6
East Cambs.	49	11.5	5.2	10	12.9	10.7
Cambridge City	78	13.2	6.5	11.7	14.6	16.4
Fenland	11	20.2	7.1	15.9	24.4	22.2

professionals concerning symptom management and clinical care issues.

DISCUSSION

Summary

This study examined the role of one H@H service in one area of the UK. Care planning and coordination of care across services during the day and at night was a key element, along with overnight nursing care provision, providing complex clinical care in sometimes challenging home situations. People who died from cancer at home were five times more likely to be supported by H@H than those with non-cancer diagnoses: this may be due to patterns of referral to the service, the referral criterion that people should be within 2 weeks of death or other reasons. In addition, the most deprived and rural area of the county was poorly served by H@H: only 11 people there received the service. Action to ensure that H@H is available for patients with conditions other than cancer and in more deprived areas is urgently needed.

Strengths and limitations

While limited to a single H@H service, the paper provides adds to the currently limited knowledge-base available for other service development. We are unable to determine the effectiveness of H@H within the descriptive study design. We were also unable to quantify or examine the day time assessment visits. Our data do not allow us to determine the number of deaths

in care homes which were supported, as 'home' was defined as the usual place of residence, which included private domestic dwellings as well as care homes. Despite it being a mandatory requirement to record patient's ethnicity in patient's medical records in UK, this data item was rarely completed in the routine clinical documents accessed as part of this evaluation. The lack of available data regarding patients' ethnicity is a notable limitation, as being from an ethnic minority is a known barrier to accessing specialise end-of-life care^{38 39}

Comparison with existing literature

H@H services vary widely across the UK, from daytime-only services provided by Clinical Nurse Specialists and Palliative Care doctors to 24 hours services staffed by healthcare assistants. Some also provide postbereavement and lymphoedema care.^{40–43} Few studies have investigated the roles that H@H services fulfil.⁴⁴

In 1971 Julian Tudor Hart wrote of the 'Inverse Care Law', an observation that 'the availability of good medical care tends to vary inversely with the need for the population served',⁴⁵ contrary to the principle of equity which is at the heart of the National Health Service. Subsequent research has repeatedly confirmed that those living in areas of higher social deprivation, with greater healthcare needs, have poorer services which they have greater difficulty accessing^{46–48} and

Table 6 Night time nursing roles of Hospice at Home team

Role	Examples
Liaison with community-based staff at night	► Liaising with out of hours nurses and GPs in order to obtain appropriate symptom relief during the night (repositioning, prescription of medications, administration of medications)
Pharmacological symptom management	► Administering 'as required' and 'just in case medicines' such as midazolam, diamorphine, lorazepam, glycopyrronium (qualified nurses only)
Specialist assessment	► 'Going in cold' to homes of new patients with very little prior knowledge in order to assess end-of-life care needs and report back to all concerned. Difficult as environment usually not suited to end-of-life care.
After death care	► Liaising with family, GPs and funeral homes to initiate post death proceedings ► Remaining with and comforting carers until body removed from home
Advocacy	► Safeguarding, passing on to higher authorities any abuse of patients and carers detected during night time conversations ► Acting as buffers between large families and patients who needed to rest ► Challenging ineffective drug regimes
Fundamental nursing care	► Pressure area care, general hygiene, comfort and reassurance
Psychological support for patient and family	► Offering counselling to distressed patients and family members

are further disadvantaged by greater multimorbidity at an earlier age.⁴⁹ The present study is consistent with this literature and with a study of a previous form of the Cambridgeshire H@H which found patients with cancer in lower socioeconomic areas were less likely to be referred.^{21 50} The Manchester H@H service has reported lower referral rates for those with greater social deprivation.²² The inverse care law is still in operation in the 21st century.

The great majority of patients supported by H@H had cancer. While this compares favourably with a H@H service piloted in North West England in which 92% of patients had cancer,⁵¹ our finding that only 11% of people who died at home with conditions other than cancer received H@H support, while 54% of people who died at home with cancer were cared for by H@H, is continuing evidence of diagnostic inequality of H@H provision.^{52–54} General practitioners in Scotland have been found to identify palliative care needs for 75% of people who died from cancer but only 20% of those who died from other conditions.⁵⁵ A study in Wales found that 46.2% of people who died of cancer had recorded palliative care needs or had received specialist palliative care, compared with only 5.3% of people who died from other causes.⁵⁶

In part this may be due to greater prognostic uncertainty concerning illness trajectories in conditions other than cancer.⁵⁷ It may also be due to organisational and institutional cultures which traditionally have had a cancer focus in palliative and end-of-life care. It is widely acknowledged that it is of urgent importance to redress this inequality in access palliative care services such as H@H: end-of-life care for those without cancer is of lower quality, a deficit that might be mitigated by a H@H service.⁵⁸

Implications for research and practice

In a society where death has become increasingly hidden and medicalised, many people who wish to support loved ones to die at home may be doing so with little or no prior experience of death and dying to draw on.⁵⁹ With ageing populations and rising multimorbidity later in life, the need for palliative care is increasing.^{60 61} As the trend towards fewer people dying in hospital continues in many, though not all countries,^{61–64} services such as H@H, which exist to support carers and patients emotionally and physically to enable a natural acceptance and dignified end of life, are becoming an increasingly vital resource to allow people to die at home.

This study has identified the important roles that a H@H service plays in the care of people approaching the end of life and in a detail that has not been reported before. It has also highlighted a dismaying gap between demand for such a service and the available supply. Robust research into alternative models of care is a priority to ensure equity of access for all people and families who would benefit from palliative

care services such as H@H, particularly those from less advantaged backgrounds and with non-cancer diagnoses. We would urge services to measure and demonstrate their effectiveness and equality of care provision.

Acknowledgements We would like to acknowledge the contributions of Dr Zoe Morris to the study design, Katie York, Andrea Newman and Andy Cowan to data collection and entry and Dr Jane Fleming for very helpful comments on the final draft of this paper.

Contributors LMot, LMor, LW and SB conceived the study. All authors contributed to the study planning and design. JB contributed to data collection. JB analysed and interpreted the data with input from SB. JB wrote the manuscript with input and feedback from SB, LMot, LMor and LW.

Funding The study was jointly funded by the Arthur Rank Hospice Charity and the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care Cambridgeshire and Peterborough (CLAHRC CP) Programme, which was commissioned as NIHR CLAHRC East of England in 2014.

Disclaimer The opinions expressed are those of the authors and not those of the NHS, the NIHR or Department of Health.

Competing interests This study was undertaken by the Palliative and End of Life Care Research Group of the University of Cambridge. It was part-funded by the Arthur Rank House Hospice Charity who part-fund the H@H service. LW, Mot and LMor are employed by the Arthur Rank Hospice Charity.

Ethics approval This study was approved by the Chair of a Local Research Ethics Committee (LREC) to be service evaluation and not to require LREC approval. The research team followed standard procedures for the ethical conduct of research involving the use of patient-identifiable information.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Department of Health. End of life care strategy. 2008 <https://www.gov.uk/government/publications/end-of-life-care-strategy-promoting-high-quality-care-for-adults-at-the-end-of-their-life>
- 2 Murtagh FEM BC, Petkova H, Sleeman KE, *et al.* *Understanding place of death for patients with non malignant conditions: a systematic review*. London: National Institute for Health Research, Queens Publisher, 2012.
- 3 Higginson IJ, Sen-Gupta GJ. Place of care in advanced cancer: a qualitative systematic literature review of patient preferences. *J Palliat Med* 2000;3:287–300.
- 4 Brogaard T, Neergaard MA, Sokolowski I, *et al.* Congruence between preferred and actual place of care and death among Danish cancer patients. *Palliat Med* 2013;27:155–64.

- 5 Gomes B, Calanzani N, Gysels M, *et al.* Heterogeneity and changes in preferences for dying at home: a systematic review. *BMC Palliat Care* 2013;12:7.
- 6 Fried TR, van Doorn C, O'Leary JR, *et al.* Older persons' preferences for site of terminal care. *Ann Intern Med* 1999;131:109–12.
- 7 Gott M, Seymour J, Bellamy G, *et al.* Older people's views about home as a place of care at the end of life. *Palliat Med* 2004;18:460–7.
- 8 Evans R, Finucane A, Vanhegan L, *et al.* Do place-of-death preferences for patients receiving specialist palliative care change over time? *Int J Palliat Nurs* 2014;20:579–83.
- 9 Fleming J, Farquhar M, Brayne C, *et al.* Death and the oldest old: attitudes and preferences for end-of-life care—qualitative research within a population-based cohort study. *PLoS One* 2016;11:e0150686.
- 10 Hoare S, Morris ZS, Kelly MP, *et al.* Do patients want to die at home? A systematic review of the UK literature, focused on missing preferences for place of death. *PLoS One* 2015;10:e0142723.
- 11 Pollock K. Is home always the best and preferred place of death? *BMJ* 2015;351:h4855.
- 12 Gomes B, Calanzani N, Higginson IJ. Reversal of the British trends in place of death: time series analysis 2004–2010. *Palliat Med* 2012;26:102–7.
- 13 De Roo ML, Miccinesi G, Onwuteaka-Philipsen BD, *et al.* Actual and preferred place of death of home-dwelling patients in four European countries: making sense of quality indicators. *PLoS One* 2014;9:e93762.
- 14 Loucka M, Payne SA, Brearley SG. Place of death in the Czech Republic and Slovakia: a population based comparative study using death certificates data. *BMC Palliat Care* 2014;13:13.
- 15 Gágyor I, Himmel W, Pierau A, *et al.* Dying at home or in the hospital? An observational study in German general practice. *Eur J Gen Pract* 2016;22:9–15.
- 16 Pivodic L, Pardon K, Morin L, *et al.* Place of death in the population dying from diseases indicative of palliative care need: a cross-national population-level study in 14 countries. *J Epidemiol Community Health* 2016;70:17–24.
- 17 National Association for Hospice at Home. What is hospice at home? 2014 <http://www.nahh.org.uk/about-hospice-care/what-is-hospice-at-home> (accessed Feb 2017).
- 18 Gellie A, Mills A, Levinson M, *et al.* Death: a foe to be conquered? Questioning the paradigm. *Age Ageing* 2015;44:7–10.
- 19 Beauchamp TL, Childress JF. *Principles of biomedical ethics*. USA: Oxford University Press, 2001.
- 20 Gomes B, Higginson IJ. Factors influencing death at home in terminally ill patients with cancer: systematic review. *BMJ* 2006;332:515–21.
- 21 Grande GE, McKerral A, Todd CJ. Which cancer patients are referred to Hospital at Home for palliative care? *Palliat Med* 2002;16:115–23.
- 22 Campbell M, Grande G, Wilson C, *et al.* Exploring differences in referrals to a hospice at home service in two socio-economically distinct areas of Manchester, UK. *Palliat Med* 2010;24:403–9.
- 23 National End of Life Care Intelligence Network. Deprivation and death: variation in place and cause of death. 2012 http://www.endoflifecare-intelligence.org.uk/resources/publications/deprivation_and_death (accessed Feb 2017).
- 24 Black H, Waugh C, Munoz-Arroyo R, *et al.* Predictors of place of death in South West Scotland 2000–2010: Retrospective cohort study. *Palliat Med* 2016;30:764–71.
- 25 Sayma M, Saleh D, Kerwat D, *et al.* A qualitative inquiry into the barriers and facilitators to achieving home death. *BMJ Support Palliat Care* 2017;1:bmjpcare-2016-001260.
- 26 Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006;3:e442.
- 27 Lunney JR, Lynn J, Foley DJ, *et al.* Patterns of functional decline at the end of life. *JAMA* 2003;289:2387–92.
- 28 Seow H, Brazil K, Sussman J, *et al.* Impact of community based, specialist palliative care teams on hospitalisations and emergency department visits late in life and hospital deaths: a pooled analysis. *BMJ* 2014;348:g3496.
- 29 Bell G, Greene K, Hunt J, *et al.* Developing national standards for hospice-at-home Services. *End of Life Journal* 2013;3:1–5.
- 30 McKay EA, Taylor AE, Armstrong C. "What she told us made the world of difference": carers' perspectives on a hospice at home service. *J Palliat Care* 2013;29:170–7.
- 31 McLaughlin D, Sullivan K, Hasson F. Hospice at home service: the carer's perspective. *Support Care Cancer* 2007;15:163–70.
- 32 Jack BA, Baldry CR, Groves KE, *et al.* Supporting home care for the dying: an evaluation of healthcare professionals' perspectives of an individually tailored hospice at home service. *J Clin Nurs* 2013;22:2778–86.
- 33 Lucas B, Small N, Greasley P, *et al.* A reassuring presence: an evaluation of bradford district hospice at home service. *BMC Palliat Care* 2008;7:9.
- 34 McBride T, Morton A, Nichols A, *et al.* Comparing the costs of alternative models of end-of-life care. *J Palliat Care* 2011;27:126–33.
- 35 Georgiou T, Bardsley M. *Exploring the cost of care at the end of life*. London: Nuffield Trust Research Report, 2014.
- 36 Council County Council. 2011 Census QA Pack - Cambridge. <https://www.cambridge.gov.uk/sites/default/files/docs/census-2011-cambridge-data.pdf> (accessed Feb 2017).
- 37 Polit DF. *Beck CT nursing research: generating and assessing evidence for nursing practice*. 10th Edn. Philadelphia: Wolters Kluwer, 2017.
- 38 Koffman J, Ho YK, Davies J, *et al.* Does ethnicity affect where people with cancer die? A population-based 10 year study. *PLoS One* 2014;9:e95052.
- 39 Evans N, Meñaca A, Koffman J, *et al.* Cultural competence in end-of-life care: terms, definitions, and conceptual models from the British literature. *J Palliat Med* 2012;15:812–20.
- 40 St John's Hospice. St John's hospice service information. <http://www.stjohnshospice.org.uk/our-care/our-services/hospice-home?phpMyAdmin=66784a0607be50a367d248d1c690334e> (accessed Feb 2017).
- 41 Hospice at Home West Cumbria. Hospice at home west cumbria service information. <http://hospiceathomewestcumbria.org.uk/> (accessed Feb 2017).
- 42 Carlisle and North Lakeland Hospice. Home service information. <http://www.hospiceathome.co.uk/> (accessed Feb 2017).
- 43 St Peter's Hospice. Service information. <http://www.stpetershospice.org.uk/patients-and-families/hospice-at-home-service/> (accessed Feb 2017).
- 44 Davis C. End of life care: a rapid response hospice at home service. *Nurs Older People* 2010;22:22–4.
- 45 Tudor Hart J. The inverse care law. *The Lancet* 1971;297:405–12.
- 46 Stirling AM, Wilson P, McConnachie A. Deprivation, psychological distress, and consultation length in general practice. *Br J Gen Pract* 2001;51:456–60.
- 47 Mercer SW, Watt GC. The inverse care law: clinical primary care encounters in deprived and affluent areas of Scotland. *Ann Fam Med* 2007;5:503–10.
- 48 Salisbury C, Johnson L, Purdy S, *et al.* Epidemiology and impact of multimorbidity in primary care: a retrospective cohort study. *Br J Gen Pract* 2011;61:12–21.
- 49 Barnett K, Mercer SW, Norbury M, *et al.* Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012;380:37–43.

- 50 Grande GE, Addington-Hall JM, Todd CJ. Place of death and access to home care services: are certain patient groups at a disadvantage? *Soc Sci Med* 1998;47:565–79.
- 51 Baldry C, Jack B, Groves K, *et al.* 'Made to fit': filling critical gaps with a hospice at home service. *BMJ Support Palliat Care* 2011;1:236.
- 52 Marie Curie. Triggers for palliative care: improving access to care for people with diseases other than cancer. <https://www.mariecurie.org.uk/globalassets/media/documents/policy/policy-publications/june-2015/triggers-for-palliative-care-full-report.pdf> (accessed Jan 2018).
- 53 Dixon JKD, Matosevic T, Clark M, *et al.* Equity of provision of palliative care in the UK. 2015 <https://www.mariecurie.org.uk/globalassets/media/documents/policy/campaigns/equity-palliative-care-uk-report-full-lse.pdf> (accessed Feb 2017).
- 54 The National Council for Palliative Care. National survey of patient activity data for specialist palliative care services: MDS full report for the year 2012-2013. http://www.endoflifecare-intelligence.org.uk/resources/publications/mds_report (accessed Jan 2018).
- 55 Zheng L, Finucane A, Oxenham D, *et al.* How good is primary care at identifying patients who need palliative care? A mixed methods study. *Euro J Palliat Care* 2013;20:216–22.
- 56 Marie Curie. Death and dying in Wales: an analysis of inconsistencies in access to specialist palliative care and hospital activity in the last year of life. <https://www.mariecurie.org.uk/globalassets/media/documents/policy/policy-publications/december-2014/death-dying-wales-full-report.pdf> (accessed Jan 2018).
- 57 Small N, Gardiner C, Barnes S, *et al.* Using a prediction of death in the next 12 months as a prompt for referral to palliative care acts to the detriment of patients with heart failure and chronic obstructive pulmonary disease. *Palliat Med* 2010;24:740–1.
- 58 Bajwah S, Higginson IJ, Ross JR, *et al.* The palliative care needs for fibrotic interstitial lung disease: a qualitative study of patients, informal caregivers and health professionals. *Palliat Med* 2013;27:869–76.
- 59 Watts JH. *Death, dying and bereavement: Issues for practice*: Dunedin Academic, 2010.
- 60 Downing J, Grant L, Leng M, *et al.* Understanding models of palliative care delivery in sub-saharan Africa: learning from programs in Kenya and Malawi. *J Pain Symptom Manage* 2015;50:362–70.
- 61 Sarmiento VP, Higginson IJ, Ferreira PL, *et al.* Past trends and projections of hospital deaths to inform the integration of palliative care in one of the most ageing countries in the world. *Palliat Med* 2016;30:363–73.
- 62 Houttekier D, Cohen J, Surkyn J, *et al.* Study of recent and future trends in place of death in Belgium using death certificate data: a shift from hospitals to care homes. *BMC Public Health* 2011;11:228.
- 63 Broad JB, Gott M, Kim H, *et al.* Where do people die? An international comparison of the percentage of deaths occurring in hospital and residential aged care settings in 45 populations, using published and available statistics. *Int J Public Health* 2013;58:257–67.
- 64 Simon ST, Gomes B, Koeskeroglu P, *et al.* Population, mortality and place of death in Germany (1950-2050) - implications for end-of-life care in the future. *Public Health* 2012;126:937–46.